

PDF version of www.hcvguidelines.org

AMERICAN ASSOCIATION FOR
THE STUDY OF LIVER DISEASES



Recommendations for Testing, Managing, and Treating Hepatitis C

Downloaded from <http://www.hcvguidelines.org>

Visit the HCV Guidance website to access the most up-to-date version

Revised Date: December 19, 2014

Complete revision made to [Initial](#), [Retreatment](#), [Monitoring](#), and *Unique Populations* ([HIV/HCV Coinfection](#), [Cirrhosis](#), [Post-Liver Transplantation](#), and [Renal Impairment](#)) sections on December 19, 2014.

INTRODUCTION

NOTICE: Guidance for hepatitis C treatment in adults is changing constantly with the advent of new therapies and other developments. A static version of this guidance, such as printouts of this website material, booklets, slides, and other materials, may be outdated by the time you read this. We urge you to review this guidance on this website (www.hcvguidelines.org) for the latest recommendations.

The landscape of treatment for hepatitis C virus (HCV) infection has evolved substantially since the introduction of highly effective HCV protease inhibitor therapies in 2011. The pace of change is expected to increase rapidly, as numerous new drugs with different mechanisms of action will likely become available over the next few years. To provide healthcare professionals with timely guidance as new therapies are available and integrated into HCV regimens, the Infectious Diseases Society of America (IDSA) and American Association for the Study of Liver Diseases (AASLD), in collaboration with the International Antiviral Society–USA (IAS–USA), have developed a web-based process for the rapid formulation and dissemination of evidence-based, expert-developed recommendations for hepatitis C management. The IAS–USA provides the structure and assistance to sustain the process that represents the work of leading authorities in hepatitis C prevention, diagnosis, and treatment in adults.

The AASLD/IDSA/IAS–USA hepatitis C Guidance addresses management issues ranging from testing and linkage to care, the crucial first steps toward improving health outcomes for HCV-infected persons, to the optimal treatment regimen in particular patient situations. Recommendations are based on evidence and are rapidly updated as new data from peer-reviewed evidence become available. For each treatment option, recommendations reflect the best possible management for a given patient and a given point of disease progression. Recommendations are graded with regard to the level of the evidence and strength of the recommendation. The AASLD/IDSA/IAS–USA hepatitis C Guidance is supported by the membership-based societies and not by pharmaceutical companies or other commercial interests. The Boards of Directors of AASLD and IDSA have appointed an oversight panel of 5 co-chairs and have selected panel members from the 2 societies and the collaborating partner based on their expertise in hepatitis C research and care. Likewise, the Guidance development process is generally consistent with that used by the IAS–USA (<https://www.iasusa.org/about/program-development-policy>).

This Guidance should be considered a "living document" in that the Guidance will be updated frequently as new information and treatments become available. This continually evolving report provides guidance on FDA-approved regimens. At times, it may also recommend off-label use of certain drugs or tests or provide guidance for regimens not yet approved by FDA. Readers should consult prescribing information and other resources for further information. Of note, the choice of treatment may, in the future, be further guided by data from cost-effectiveness studies.

Changes made on this page on September 25, 2014.

METHODS

The Guidance was developed by a panel of HCV experts in the fields of hepatology and infectious diseases, using an evidence-based review of information that is largely available to healthcare practitioners. The process and detailed methods for developing the Guidance are detailed in [Methods Table 1](#). Recommendations were graded according to the strength of the recommendation and quality of the supporting evidence (see [Methods Table 2](#)). Commonly used abbreviations are expanded in [Methods Table 3](#).

Methods Table 1. Summary of the Process and Methods for the Guidance Development

Topic	Description
Statement of Need	Increased awareness of the rising number of complications of hepatitis C virus (HCV) infection, the recent screening initiatives by the Centers for Disease Control and Prevention (CDC) and US Preventive Services Task Force (USPSTF), and the rapid evolution of highly effective antiviral therapy for HCV infection have driven a need for timely guidance on how new developments change practice for health care professionals.
Goal of the Guidance	The goal of the Guidance is to provide up-to-date recommendations to health care practitioners on the optimal screening, management, and treatment for adults with HCV infection in the United States, considering the best available evidence. The Guidance is updated regularly, as new data, information, and tools and treatments become available.
Panel members	Panel members are chosen based on their expertise in the diagnosis, management, and treatment of HCV infection. Members from the fields of hepatology and infectious diseases are included, as well as HCV community representatives. Members were appointed by the respective Sponsor Societies after vetting by an appointed Sponsor Society committee. The Panel chairs are appointed by the Society boards, 2 each from the Sponsor Societies and 1 representing the Collaborating Partner. All Panel chairs and members serve as volunteers (not compensated) for defined terms (2-3 years), which may be renewed based on panel needs.
Conflict of interest management	<p>The panel was established with the goal of having no personal (ie, direct payment to the individual) financial conflicts of interest among its chairs and among fewer than half of its panel members. All potential panel members are asked to disclose any personal relationship with a pharmaceutical, biotechnology, medical device, or health-related company or venture that may result in financial benefit. Disclosures are obtained prior to the panel member appointments and for 1 year prior to the initiation of the work of the panel. Full transparency of potential financial conflicts is an important goal for the guidance that best ensures the credibility of the processed and the recommendations.</p> <p>Individuals are also asked to disclose funding of HCV-related research activities to their institutional division, department, or practice group.</p> <p>Disclosures are reviewed by the HCV Guidance Chairs, who make assessments based on the conflict-of-interest policies of the sponsoring organizations (AASLD and IDSA) and the collaborative partner (IAS–USA). Personal and institutional financial relationships with commercial entities that have products in the field of hepatitis C are assessed.</p> <p>The following relationships are prohibited during membership on the guidance panel and are grounds for exclusion from the panel:</p>

- Employment with any commercial company with products in the field of hepatitis C.
- An ownership interest in a commercial entity that produces hepatitis C products.
- Participation in/payment for promotional or marketing activities sponsored by companies with HCV-related products including non-CME educational activities or speakers bureaus for audiences outside of the company.

The following relationships or activities are reportable but were not deemed to merit exclusion:

- Commercial support of research that is paid to an organization or practice group. Due to the rapidly evolving nature of the subject matter, having individuals with expertise in the particular clinical topic is crucial to developing the highest-quality and most-informed recommendations. To that end, research support from commercial entities is not considered grounds for panel exclusion (an unresolvable conflict) if the funding of the research was paid to the institution or practice group, as opposed to the individual. In the instance of someone conducting clinical research in a community practice, research funds to the group practice were acceptable.
- Participation on commercial company scientific advisory boards. Participation in advisory boards, data safety monitoring boards, or in consultancies sponsored by the research arm of a company (eg, study design or data safety monitoring board) is considered a potential personal conflict but is not considered a criterion for exclusion.

The HCV Guidance Chairs achieved a majority of panel members with no personal financial interests.

Panel members are asked to inform the group of any changes to their disclosure status and are given the opportunity to recuse themselves (or be recused) from the discussion where a perceived conflict of interest that cannot be resolved exists.

Financial disclosures for each Panel member can be accessed [here](#).

Intended Audience Sponsors, funding, and collaborating partner

Medical practitioners especially those who provide care to or manage patients with hepatitis C.

The American Association for the Study of Liver Diseases (AASLD) and the Infectious Diseases Society of America (IDSA) are the sponsors of the Guidance and provide ongoing financial support. The International Antiviral Society-USA (IAS-USA) is the Collaborating Partner responsible for managing the Panel and the Guidance development process.

Grant support was sought and obtained from the Centers for Disease Control and Prevention (CDC) for the initial gathering and review of evidence related to hepatitis C screening and testing recommendations and interventions to

implement HCV screening in clinical settings.

Evidence identification and collection

The Guidance is developed using an evidence-based review of information that is largely available to health care practitioners. Data from the following sources are considered by Panel members when making recommendations: research published in the peer-reviewed literature or presented at major national or international scientific conferences; safety warnings from the US Food and Drug Administration (FDA) or other regulatory agencies or from manufacturers; drug interaction data; prescribing information from FDA-approved products; and registration data for new products under FDA review. Press releases, unpublished reports, and personal communications are generally not considered.

Literature searches are conducted regularly and before each major revision to ensure that the Panel addresses all relevant published data. Medical subject headings and free text terms are combined to maximize retrieval of relevant citations from the PubMed, Scopus, EMBASE, and Web of Science databases. To be considered for inclusion, articles were required to have been published in English from 2010 to the present. Data from abstracts presented at national or international scientific conferences are also considered

Grading of the evidence and RECOMMENDATIONS

The Guidance is presented in the form of RECOMMENDATIONS. Each RECOMMENDATION is rated in terms of the level of the evidence and strength of the recommendation, using a modification of the scale adapted from the American College of Cardiology and the American Heart Association Practice Guidelines. ([American Heart Association, 2014](#)); ([Shiffman, 2003](#)) A summary of the supporting (and conflicting) evidence follows each RECOMMENDATION or set of RECOMMENDATIONS.

Data review and synthesis and preparation of RECOMMENDATIONS and supporting information

Draft RECOMMENDATIONS are developed by subgroups of the full Panel with interest and expertise in particular sections of the Guidance. Following development of supporting text and references, the sections are reviewed by the full Panel and Chairs. A penultimate draft is submitted to the AASLD and IDSA Governing Boards for final review and approval before posting online on the website, www.hcvguidelines.org.

Subgroups of the Panel meet regularly by conference call as needed to update RECOMMENDATIONS and supporting evidence. Updates may be prompted by new publications or presentations at major national or international scientific conferences, new drug approvals (or new indications, dosing formulations, or frequency of dosing), new safety warnings, or other information that may have a substantial impact on the clinical care of patients. Updates and changes in the Guidance are indicated by highlighted text on the online site and a notice of update is posted on the Home Page.

Abbreviations

Commonly used abbreviations in the text with their expansions are listed in [Methods Table 3](#).

Opportunity for Comments

Evidence-based comments may be submitted to the Panel by email to hcvguidelines@iasusa.org, or by clicking on the "Send a comment to the Panel" button on www.hcvguidelines.org/contact-us. The Panel considers evidence-based comments about the RECOMMENDATIONS, grades, and evidence summary but should not be contacted for individual patient

management questions.

Changes on this page made on November 20, 2014.

Methods Table 2. Grading System Used to Rate the Level of the Evidence and Strength of the Recommendation for Each Recommendation

Recommendations are based on scientific evidence and expert opinion. Each recommended statement includes a Roman numeral (I, II, or III) that represents the level of the evidence that supports the recommendation, and a letter (A, B, or C) that represents the strength of the recommendation.

Classification	Description
Class I	Conditions for which there is evidence and/or general agreement that a given diagnostic evaluation, procedure, or treatment is beneficial, useful, and effective
Class II	Conditions for which there is conflicting evidence and/or a divergence of opinion about the usefulness and efficacy of a diagnostic evaluation, procedure, or treatment
Class IIa	Weight of evidence and/or opinion is in favor of usefulness and efficacy
Class IIb	Usefulness and efficacy are less well established by evidence and/or opinion
Class III	Conditions for which there is evidence and/or general agreement that a diagnostic evaluation, procedure, or treatment is not useful and effective or if it in some cases may be harmful
Level of Evidence	Description
Level A*	Data derived from multiple randomized clinical trials, meta-analyses, or equivalent
Level B*	Data derived from a single randomized trial, nonrandomized studies, or equivalent
Level C	Consensus opinion of experts, case studies, or standard of care

Adapted from the American College of Cardiology and the American Heart Association Practice Guidelines. ([American Heart Association, 2011](#)); ([Shiffman, 2003](#))

*In some situations, such as for IFN-sparing HCV treatments, randomized clinical trials with an existing standard-of-care arm cannot ethically or practicably be conducted. The US Food and Drug Administration (FDA) has suggested alternative study designs, including historical controls or immediate versus deferred, placebo-controlled trials. For additional examples and definitions see FDA link:

<http://www.fda.gov/downloads/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/UCM225333.pdf>. In those instances for which there was a single pre-determined, FDA-approved equivalency established, panel members considered the evidence as equivalent to a randomized controlled trial for levels A or B.

Methods Table 3. Commonly Used Abbreviations and Their Expansions

Abbreviation	Expansion or Notes
<i>These terms are not expanded in text</i>	
HCV	hepatitis C virus. In this Guidance "hepatitis C virus" and HCV refer to the virus. Hepatitis C and HCV infection or HCV disease refer to the resulting disease.
IFN	interferon alfa
PEG	peginterferon alfa
<i>These terms are expanded at first mention in text</i>	
ALT	alanine aminotransferase
AST	aspartate aminotransferase
BOC	boceprevir
CBC	complete blood cell (eg, complete blood cell count)
CrCl	creatinine clearance
CTP	Child Turcotte Pugh (see below)
DAA	direct-acting antiviral
ESRD	end-stage renal disease
GFR	glomerular filtration rate
HBsAg	hepatitis B virus surface antigen
HBV	hepatitis B virus
HCC	hepatocellular carcinoma
IDU	injection drug use or user
INR	international normalized ratio
MELD	model for end-stage liver disease
MSM	men who have sex with men
OATP	organic anion-transporting polypeptide
P-gp	p-glycoprotein
RAV	resistance-associated variant
RBC	red blood cell (eg, red blood cell count)
RBV	ribavirin
RGT	response-guided therapy
RVR	rapid virologic response
sAg	surface antigen
SMV	simeprevir; used for the treatment of those with genotype 1 of hepatitis C virus (HCV) who have compensated liver disease, including cirrhosis
SOF	sofosbuvir; a nucleoside analogue used in combination with other drugs for the treatment of HCV infection
SVR12 (or 24 or 48, etc)	sustained virologic response at 12 weeks (or at 24 weeks, or at 48 weeks, etc)
TSH	thyroid-stimulating hormone
TVR	telaprevir; an antiviral agent to treat hepatitis C
US FDA	US Food and Drug Administration
Definition of Terms	
Child Turcotte Pugh	Class A Class B Class C

(CTP) classification of the severity of cirrhosis	Total points	5–6	7–9	10–15
	Factor	1 Point	2 Points	3 Points
	Total bilirubin (µmol/L)	<34	34–50	>50
	Serum albumin (g/L)	>35	28–35	<28
	Prothrombin time/international normalized ratio	<1.7	1.71–2.30	>2.30
	Ascites	None	Mild	Moderate to Severe
	Hepatic encephalopathy	None	Grade I–II (or suppressed with medication)	Grade III–IV (or refractory)
IFN ineligible	IFN ineligible is defined as one or more of the below: <ul style="list-style-type: none"> • Intolerance to IFN • Autoimmune hepatitis and other autoimmune disorders • Hypersensitivity to PEG or any of its components • Decompensated hepatic disease • Major uncontrolled depressive illness • A baseline neutrophil count below 1500/µL, a baseline platelet count below 90,000/µL or baseline hemoglobin below 10 g/dL • A history of preexisting cardiac disease 			
Relapser	a person who has achieved an undetectable level of virus during a prior treatment course of PEG/RBV and relapsed after treatment was stopped			

HCV TESTING AND LINKAGE TO CARE

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

A summary of recommendations for Testing and Linkage to Care is found in the [BOX](#).

One-time HCV testing is recommended for persons born between 1945 and 1965*, without prior ascertainment of risk.

Rating: Class I, Level B

Other persons should be screened for risk factors for HCV infection, and 1-time testing should be performed for all persons with behaviors, exposures, and conditions associated with an increased risk of HCV infection.

1. Risk behaviors

Injection-drug use (current or ever, including those who injected once)

Intranasal illicit drug use

2. Risk exposures

Long-term hemodialysis (ever)

Getting a tattoo in an unregulated setting

Healthcare, emergency medical, and public safety workers after needlesticks, sharps, or mucosal exposures to HCV-infected blood

Children born to HCV-infected women

Prior recipients of transfusions or organ transplants, including persons who:

- were notified that they received blood from a donor who later tested positive for HCV infection
- received a transfusion of blood or blood components, or underwent an organ transplant before July 1992
- received clotting factor concentrates produced before 1987

Persons who were ever incarcerated

3. Other

HIV infection

Unexplained chronic liver disease and chronic hepatitis including elevated alanine aminotransferase levels

Solid organ donors (deceased and living)

Rating: Class I, Level B

*Regardless of country of birth

Of the estimated 2.2 to 3.2 million persons (2003 to 2010 National Health and Nutrition Examination Survey of the US noninstitutionalized civilian population) ([Denniston, 2014](#)) chronically infected with HCV in the United States, approximately 50 % are unaware that they are infected. ([Denniston, 2012](#)) Identification of those with active infection is the first step toward improving health outcomes among persons with HCV infection and preventing transmission. ([Smith, 2012](#)); ([US Preventive Services Task Force, 2013](#)); ([Centers for Disease Control and Prevention, 1998](#))

HCV testing is recommended in select populations based on demography, prior exposures, high-risk behaviors, and medical conditions. Recommendations for testing are based on HCV prevalence in these populations, proven benefits of care and treatment in reducing the risk of hepatocellular carcinoma and all-cause mortality, and the potential public health benefit of reducing transmission through early treatment, viral clearance, and reduced risk behaviors. ([Smith, 2012](#)); ([US Preventive Services Task Force, 2013](#)); ([Centers for Disease Control and Prevention, 1998](#))

HCV is primarily transmitted through percutaneous exposure to blood. Other modes of transmission include mother-to-infant and contaminated devices shared for noninjection drug use; sexual transmission also occurs but generally seems to be inefficient except among HIV-infected men who have unprotected sex with men. ([Schmidt, 2014](#)) The most important risk for HCV infection is injection drug use, accounting for at least 60% of acute HCV infections in the United States. Health care exposures are important sources of transmission, including the receipt of blood products before 1992 (after which routine screening of blood supply was implemented), receipt of clotting factor concentrates before 1987, long-term hemodialysis, needlestick injuries among healthcare workers, and patient-to-patient transmission resulting from poor infection control practices. Other risk factors include having been born to an HCV-infected mother, having been incarcerated, and having received a tattoo in an unregulated setting. The importance of these risk factors might differ based on geographic location and population. ([US Preventive Services Task Force, 2013](#)); ([Centers for Disease Control and Prevention, 1998](#)). An estimated 29% of incarcerated persons in North America are anti-HCV positive, supporting the recommendation to test this population for HCV. ([Larney, 2013](#)) Because of shared transmission modes, persons with HIV infection are at risk for HCV; sexual transmission is a particular risk for HIV-infected men who have unprotected sex with men. ([Hosein,](#)

[2013](#)); ([van de Laar, 2010](#)) Recent data also support testing in all deceased and living solid-organ donors because of the risk of HCV infection posed to the recipient. ([Seem, 2013](#)); ([Lai, 2013](#)) Although Centers for Disease Control and Prevention (CDC) and US Preventive Services Task Force hepatitis C testing guidelines do not specifically recommend testing immigrants from countries with a high prevalence (eg, Egypt or Pakistan) of hepatitis C virus infection, such persons should be tested if they were born from 1945 through 1965 or if they have risk factors (listed in [Summary Box](#)) for infection.

In 2012, CDC expanded its guidelines originally issued in 1998 ([Centers for Disease Control and Prevention, 1998](#)) for risk-based HCV testing with a recommendation to offer a 1-time (see [Summary Box](#)) HCV test to all persons born from 1945 through 1965, without prior ascertainment of HCV risk-factors. This recommendation was supported by evidence demonstrating that a risk-based strategy alone failed to identify more than 50% of HCV infections in part due to patient underreporting of their risk and provider limitations in ascertaining risk-factor information. Furthermore, persons in the 1945 to 1965 birth cohort accounted for nearly three-fourths of all HCV infections, with a 5-times higher prevalence (3.25%) than other persons, reflecting a higher incidence of HCV infections in the 1970s and 1980s (peaking at 230,000, compared with 15,000 in 2009). A recent retrospective review showed that 68% of persons with HCV infection would have been identified through a birth cohort testing strategy, whereas only 27% would have been screened with the risk-based approach. ([Mahajan, 2013](#)) The cost-effectiveness of 1-time birth cohort testing is comparable to that of current risk-based screening strategies. ([Smith, 2012](#))

CDC and the US Preventive Services Task Force (USPSTF) both recommend a 1-time HCV test in asymptomatic persons belonging to the 1945 to 1965 birth cohort and other persons based on exposures, behaviors, and conditions that increase risk for HCV infection.

Annual HCV testing is recommended for persons who inject drugs and for HIV-seropositive men who have unprotected sex with men. Periodic testing should be offered to other persons with ongoing risk factors for exposure to HCV.

Rating: Class IIA, Level C

Evidence regarding the frequency of testing in persons at risk for ongoing exposure to HCV is lacking; therefore, clinicians should determine the periodicity of testing based on the risk of reinfection. Because of the high incidence of HCV infection among persons who inject drugs and among HIV-infected MSM who have unprotected sex ([Aberg, 2013](#)); ([Linan, 2012](#)); ([Wandeler, 2012](#)); ([Witt, 2013](#)); ([Bravo, 2012](#)); ([Williams, 2011](#)), at least annual HCV testing is recommended in these subgroups.

Implementation of clinical decision support tools or prompts for HCV testing in electronic health records could facilitate reminding clinicians of HCV testing when indicated. ([Hsu, 2013](#)); ([Litwin, 2012](#)); (<http://nvhr.org/EMR>)

An anti-HCV test is recommended for HCV testing, and if the result is positive, current infection should be confirmed by a sensitive HCV RNA test.

Rating: Class I, Level A

Among persons with a negative anti-HCV test who are suspected of having liver disease, testing for HCV RNA or follow-up testing for HCV antibody is recommended if exposure to HCV occurred within the past 6 months; testing for HCV RNA can also be considered in persons who are immunocompromised.

Rating: Class I, Level C

Among persons at risk of reinfection after previous spontaneous or treatment-related viral clearance, initial HCV-RNA testing is recommended because an anti-HCV test is expected to be positive.

Rating: Class I, Level C

Quantitative HCV RNA testing is recommended prior to the initiation of antiviral therapy to document the baseline level of viremia (ie, baseline viral load).

Rating: Class I, Level A

Testing for HCV genotype is recommended to guide selection of the most appropriate antiviral regimen.

Rating: Class I, Level A

If found to have positive results for anti-HCV test and negative results for HCV RNA by polymerase chain reaction (PCR), persons should be informed that they do not have evidence of current (active) HCV infection.

Rating: Class I, Level A

All persons recommended for HCV testing should first be tested for HCV antibody (anti-HCV) ([Centers for Disease Control and Prevention \[CDC\], 2013](#)); ([Alter, 2003](#)) using an FDA-approved test. FDA-approved tests include laboratory-based assays and a point-of-care assay (ie, OraQuick HCV Rapid Antibody Test [OraSure Technologies]). ([Lee, 2011](#)) The latter is an indirect immunoassay with a sensitivity and specificity similar to those of FDA-approved laboratory-based HCV antibody assays.

A positive test result for anti-HCV indicates either current (active) HCV infection (acute or chronic), past infection that has resolved, or a false-positive test result. ([Pawlotsky, 2002](#)) Therefore, an HCV nucleic acid test (NAT) to detect viremia is necessary to confirm current (active) HCV infection and guide clinical management, including initiation of HCV treatment. HCV RNA testing should also be performed in persons with a negative anti-HCV test who are either immunocompromised (eg, persons receiving chronic hemodialysis) ([KDIGO, 2008](#)) or who might have been exposed to HCV within the last 6 months because

these persons may be anti-HCV negative. An HCV RNA test is also needed to detect reinfection in anti-HCV–positive persons after previous spontaneous or treatment-related viral clearance.

An FDA-approved quantitative or qualitative NAT with a detection level of 25 IU/mL or lower should be used to detect HCV RNA. [Testing and Linkage to Care Table 1](#) lists FDA-approved, commercially available anti-HCV screening assays. [Testing and Linkage to Care Figure 1](#) shows the CDC-recommended testing algorithm.

Persons who have positive results for an anti-HCV test and negative results for HCV RNA by polymerase chain reaction (PCR) should be informed that they do not have laboratory evidence of current (active) HCV infection. Additional HCV testing is typically unnecessary. The HCV RNA test can be repeated when there is a high index of suspicion for recent infection or in patients with ongoing risk factors for HCV infection.

Practitioners or persons may seek additional testing to learn if the HCV antibody test represents a remote HCV infection that has resolved or a false-positive result. For patients with no apparent risk for HCV infection, the likelihood of a false-positive HCV antibody test is directly related to the HCV prevalence in the tested population; false-positive test results for anti-HCV are most common for populations with a low prevalence of HCV infection. ([Alter, 2003](#)) If further testing is desired to distinguish between true positivity and biologic false positivity for HCV antibody, testing may be done with a second FDA-approved HCV antibody assay that is different from the assay used for initial antibody testing. A biologic false result should not occur with 2 different tests. ([Vermeersch, 2008](#)); ([Centers for Disease Control and Prevention \[CDC\], 2013](#)) Prior to the initiation of HCV therapy, quantitative HCV RNA testing may be used to determine the baseline level of viremia (ie, viral load) in order to define the duration of treatment for certain regimens. The degree of viral load decline after initiation of treatment is less predictive of sustained virologic response in the era of direct-acting antiviral therapy (**see section on [Pretreatment and On-Treatment Monitoring](#)**). Testing for HCV genotype helps to guide selection of the most appropriate treatment regimen.

Persons with current (active) HCV infection should receive education and interventions aimed at reducing progression of liver disease and preventing transmission of HCV.

Rating: Class IIa, Level B

1. *Abstinence from alcohol and, when appropriate, interventions to facilitate cessation of alcohol consumption should be advised for all persons with HCV infection.*

Rating: Class IIa, Level B

2. *Evaluation for other conditions that may accelerate liver fibrosis, including HBV and HIV infections, is recommended for all persons with HCV infection.*

Rating: Class IIb, Level B

3. *Evaluation for advanced fibrosis using liver biopsy, imaging, or noninvasive markers is recommended for all persons with HCV infection, to facilitate an*

appropriate decision regarding HCV treatment strategy and to determine the need for initiating additional measures for the management of cirrhosis (eg, hepatocellular carcinoma screening).

Rating: Class I, Level B

4. *Vaccination against hepatitis A and hepatitis B is recommended for all susceptible persons with HCV infection.*

Rating: Class IIa, Level C

5. *All persons with HCV infection should be provided education on how to avoid HCV transmission to others.*

Rating: Class I, Level C

In addition to receiving therapy, HCV-infected persons should be educated about how to prevent further damage to their liver. Most important is prevention of the potential deleterious effect of alcohol. Numerous studies have found a strong association between the use of excess alcohol and the development or progression of liver fibrosis and the development of hepatocellular carcinoma. ([Poynard, 1997](#)); ([Harris, 2001](#)); ([Wiley, 1998](#)); ([Corrao, 1998](#)); ([Bellentani, 1999](#)); ([Noda, 1996](#)); ([Safdar, 2004](#))

The daily consumption of more than 50 grams of alcohol has a high likelihood of worsening fibrosis. Some studies indicate that daily consumption of smaller amounts of alcohol also has a deleterious effect on the liver; however, these data are controversial. ([Westin, 2002](#)) Excess alcohol intake may also cause steatohepatitis. Alcohol screening and brief interventions such as those outlined by the National Institute of Alcohol Abuse and Alcoholism (http://pubs.niaaa.nih.gov/publications/Practitioner/CliniciansGuide2005/clinicians_guide.htm) have been demonstrated to reduce alcohol consumption and episodes of binge drinking in the general population and among HCV-infected persons who consume alcohol heavily. ([Whitlock, 2004](#)); ([Dieperink, 2010](#)); ([Proeschold-Bell, 2012](#)) Persons identified as abusing alcohol and having alcohol dependence require treatment and consideration for referral to an addiction specialist.

HBV and HIV coinfection have been associated with poorer prognosis of HCV in cohort studies. ([Thein, 2008a](#)); ([Zarski, 1998](#)) Owing to overlapping risk factors for these infections and additional benefits of their identification and treatment, persons with HCV should be tested for HIV antibody and hepatitis B surface antigen (HBsAg) using standard assays for screening ([Moyer, 2013](#)); ([Centers for Disease Control and Prevention, 2008](#)) (<http://www.aafp.org/afp/2008/0315/p819.html>) and (<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5708a1.htm>) and counseled on how to reduce their risk of acquiring these infections, including through HBV vaccination (see below).

Patients with obesity and metabolic syndrome having underlying insulin resistance are more prone to have nonalcoholic fatty liver disease, which is a risk factor for fibrosis progression in HCV-infected persons. ([Hourigan, 1999](#)); ([Ortiz, 2002](#)) Therefore, HCV-infected persons who are overweight or obese

(defined by a body mass index 25 kg/m² or higher or 30 kg/m² or higher, respectively) should be counseled regarding strategies to reduce weight and improve insulin resistance via diet, exercise, and medical therapies. (Musso, 2010); (Shaw, 2006) Patients with HCV infection and hyperlipidemia or cardiovascular comorbidities may also benefit from various hypolipidemic drugs. Prospective studies have demonstrated the safety and efficacy of statins in patients with chronic HCV and others with compensated chronic liver disease. (Lewis, 2007) Therefore, these agents should not be withheld in HCV-infected patients.

The severity of liver disease associated with chronic HCV infection is a key factor in determining the initial and follow-up evaluation of patients. Although patients with more advanced disease may have a lower response to HCV therapy, they are also most likely to derive the greatest survival benefit. (Ghany, 2011) A liver biopsy can provide objective, semiquantitative information regarding the amount and pattern of collagen or scar tissue in the liver that can assist with treatment and monitoring plans. The Metavir fibrosis score (F0-F4) and Ishak fibrosis score (0-6) are commonly used to score the amount of hepatic collagen. A liver biopsy can also help assess the severity of liver inflammation, or of hepatic steatosis, and help exclude competing causes of liver injury. (Kleiner, 2005) However, the procedure has a low but real risk of complications, and sampling artifact makes its serial use in most patients less desirable. (Regev, 2002) Noninvasive methods frequently used to estimate liver disease severity include a liver-directed physical exam (normal in most patients), routine blood tests (eg, serum alanine aminotransferase [ALT] and aspartate aminotransferase [AST], albumin, bilirubin, international normalized ratio levels, and complete blood cell counts with platelets), serum fibrosis marker panels, liver imaging (eg, ultrasound, computed tomography scan), and transient elastography. Simple blood tests (eg, serum AST-to-platelet ratio index [APRI]) (Wai, 2003) (<http://www.hepatitisc.uw.edu/page/clinical-calculators/apri>) and assessment of liver surface nodularity and spleen size by liver ultrasound or other cross-sectional imaging modalities can help determine if patients with HCV have occult portal hypertension, which is associated with a greater likelihood of developing future hepatic complications in untreated patients. (Chou, 2013); (Rockey, 2006) Liver elastography can provide instant information regarding liver stiffness at the point of care and can reliably distinguish patients with a high versus low likelihood of cirrhosis. (Castera, 2012); (Bonder, 2014) Because persons with known or suspected bridging fibrosis and cirrhosis are at increased risk of developing complications of advanced liver disease, they require more frequent follow-up; these persons should also avoid hepatotoxic drugs (eg, excessive acetaminophen [ie, ≥2 g/d] or certain herbal supplements) or nephrotoxic drugs (eg, nonsteroidal antiinflammatory drugs) and receive ongoing imaging surveillance for liver cancer and gastroesophageal varices. (Sangiovanni, 2006); (Fontana, 2010)

Exposure to infected blood is the primary mode of HCV transmission. HCV-infected persons must be informed of the precautions needed to avoid exposing others to infected blood. This is particularly important for persons who use injection drugs, given that HCV transmission in this population primarily results from the sharing of needles and other infected implements. Recently, epidemics of acute HCV due to sexual transmission in HIV-infected men who have sex with men have also been described. (van de Laar, 2009); (Urbanus, 2009); (Fierer, 2008) **Testing and Linkage to Care Table 2** outlines measures to avoid HCV transmission. HCV is not spread by sneezing, hugging, holding hands, coughing, or sharing eating utensils or drinking glasses, nor is it transmitted through food or water.

Evaluation by a practitioner who is prepared to provide comprehensive management, including consideration of antiviral therapy, is recommended for all persons with current (active) HCV infection.

Rating: Class IIa, Level C

The definition of evaluation is: *Patient has attended a medical care visit with a practitioner able to complete a full assessment, the pros and cons of antiviral therapy have been discussed, and the patient has been transitioned into treatment, if appropriate.*

Improvement in identification of current (active) HCV infection and advances in treatment regimens will have limited impact on HCV-related morbidity and mortality without concomitant improvement in linkage to care. All patients with current HCV infection and a positive HCV RNA test result should be evaluated by a practitioner with expertise in assessment of liver disease severity and HCV treatment. Subspecialty care and consultation are required for persons with HCV infection who have advanced fibrosis or cirrhosis (stage F3 or above on Metavir scale), including possible referral for consideration of liver transplantation. In the United States, only an estimated 13% to 18% of persons chronically infected with HCV receive treatment. ([Holmberg, 2013](#)) Lack of appropriate practitioner assessment and delays in linkage to care can result in negative health outcomes. Further, patients who are lost to follow-up fail to benefit from evolving evaluation and treatment options.

Commonly cited patient-related barriers to treatment initiation include contraindications to treatment (eg, medical or psychiatric comorbidities), lack of acceptance of treatment (eg, asymptomatic nature of disease, competing priorities, low treatment efficacy, and long treatment duration and adverse effects), and lack of access to treatment (eg, cost and distance to specialist). ([Khokhar, 2007](#)); ([Arora, 2011](#)); ([Clark, 2012](#)) Common practitioner-related barriers include perceived patient-related barriers (eg, fear of adverse effects, treatment duration, cost, and effectiveness), lack of expertise in HCV treatment, lack of specialty referral resources, resistance to treating persons currently using illicit drugs or alcohol, and concern about cost of HCV treatment. ([Morrill, 2005](#)); ([Reilley, 2013](#)); ([McGowan, 2013](#)) Some possible strategies to address these barriers are listed in [Testing and Linkage to Care Table 3](#). One strategy that addresses several barriers is colocalization of HCV screening, evaluation, and treatment with other medical or social services. Colocalization has already been applied to settings with a high prevalence of HCV infection (eg, correctional facilities and programs providing needle exchange, substance abuse treatment, and methadone maintenance) but is not uniformly available. ([Islam, 2012](#)); ([Stein, 2012](#)); ([Bruggmann, 2013](#))

A strategy that addresses lack of access to specialists (a primary barrier to hepatitis C care) is participation in models involving close collaboration between primary care practitioners and subspecialists. ([Arora, 2011](#)); ([Rossaro, 2013](#)); ([Miller, 2012](#)) Such collaborations have used telemedicine and knowledge networks to overcome geographic distances to specialists. ([Arora, 2011](#)); ([Rossaro, 2013](#)) For example, Project ECHO (Extension for Community Healthcare Outcomes [<http://www.echohcvexperts.com>]) uses videoconferencing to enhance primary care practitioner capacity in rendering HCV care and treatment to New Mexico's large rural and underserved population. ([Arora, 2011](#)) Through case-based learning and real-time feedback from a multidisciplinary team of specialists (ie, gastroenterology, infectious diseases, pharmacology, and psychiatry practitioners), Project ECHO has expanded access to HCV infection treatment in populations that might have otherwise remained untreated.

Additional strategies for enhancing linkage to care could be adapted from other fields, such as tuberculosis and HIV, but remain to be evaluated for HCV infection. For example, use of directly observed therapy has enhanced adherence to tuberculosis treatment, and use of case managers and patient navigators has reduced loss of follow-up in HIV care. ([Govindasamy, 2012](#)) An assessment of efficacy and comparative effectiveness of these strategies is a crucial area of future research for patients with HCV infection. Replication and expansion of best practices and new models for linkage to HCV care will also be crucial to

maximize the public health impact of newer treatment paradigms.

Changes made on January 28, 2015.

Testing and Linkage to Care Box: Summary of Recommendations for Testing and Linkage to Care

Testing and Linkage To Care Box. Summary of Recommendations for Testing and Linkage to Care

One-time HCV testing is recommended at for persons born between 1945 and 1965*, without prior ascertainment of risk.

Rating: Class I, Level B

Other persons should be screened for risk factors for HCV infection, and 1-time testing should be performed for all persons with behaviors, exposures, and conditions associated with an increased risk of HCV infection.

1. Risk behaviors

Injection-drug use (current or ever, including those who injected once)

Intranasal illicit drug use

2. Risk exposures

Long-term hemodialysis (ever)

Getting a tattoo in an unregulated setting

Healthcare, emergency medical, and public safety workers after needlesticks, sharps, or mucosal exposures to HCV-infected blood

Children born to HCV-infected women

Prior recipients of transfusions or organ transplants, including persons who:

- were notified that they received blood from a donor who later tested positive for HCV infection
- received a transfusion of blood or blood components, or underwent an organ transplant before July 1992

- received clotting factor concentrates produced before 1987

Persons who were ever incarcerated

3. Other

HIV infection

Unexplained chronic liver disease and chronic hepatitis including elevated alanine aminotransferase levels

Solid organ donors (deceased and living)

Rating: Class I, Level B

*Regardless of country of birth

Annual HCV testing is recommended for persons who inject drugs and for HIV-seropositive men who have unprotected sex with men. Periodic testing should be offered to other persons with ongoing risk factors for exposure to HCV.

Rating: Class IIA, Level C

An anti-HCV test is recommended for HCV testing, and if the result is positive, current infection should be confirmed by a sensitive HCV RNA test.

Rating: Class I, Level A

Among persons with a negative anti-HCV test who are suspected of having liver disease, testing for HCV RNA or follow-up testing for HCV antibody is recommended if exposure to HCV occurred within the past 6 months; testing for HCV RNA can also be considered in persons who are immunocompromised.

Rating: Class I, Level C

Among persons at risk of reinfection after previous spontaneous or treatment-related viral clearance, initial HCV-RNA testing is recommended because an anti-HCV test is expected to be positive.

Rating: Class I, Level C

Quantitative HCV RNA testing is recommended prior to the initiation of antiviral therapy to document the baseline level of viremia (ie, baseline viral load).

Rating: Class I, Level A

Testing for HCV genotype is recommended to guide selection of the most appropriate antiviral regimen.

Rating: Class I, Level A

If found to have positive results for anti-HCV test and negative results for HCV RNA by polymerase chain reaction, persons should be informed that they do not have evidence of current (active) HCV infection.

Rating: Class I, Level A

Persons with current (active) HCV infection should receive education and interventions aimed at reducing progression of liver disease and preventing transmission of HCV.

Rating: Class IIa, Level B

- 1. Abstinence from alcohol and, when appropriate, interventions to facilitate cessation of alcohol consumption should be advised for all persons with HCV infection.*

Rating: Class IIa, Level B

- 2. Evaluation for other conditions that may accelerate liver fibrosis, including HBV and HIV infections, is recommended for all persons with HCV infection.*

Rating: Class IIb, Level B

- 3. Evaluation for advanced fibrosis using liver biopsy, imaging, or noninvasive markers is recommended for all persons with HCV infection to facilitate an appropriate decision regarding HCV treatment strategy and to determine the need for initiating additional measures for the management of cirrhosis (eg, hepatocellular carcinoma screening).*

Rating: Class I, Level B

- 4. Vaccination against hepatitis A and hepatitis B is recommended for all susceptible persons with HCV infection.*

Rating: Class IIa, Level C

5. *All persons with HCV infection should be provided education on how to avoid HCV transmission to others.*

Rating: Class I, Level C

Evaluation by a practitioner who is prepared to provide comprehensive management, including consideration of antiviral therapy, is recommended for all persons with current (active) HCV infection.

Rating: Class IIa, Level C

Testing and Linkage to Care Table 1. FDA-approved, Commercially Available Anti-HCV Screening Assays

Assay	Manufacturer	Format
Abbott HCV EIA 2.0	Abbott	EIA (Manual)
Advia Centaur HCV	Siemens	CIA (Automated)
ARCHITECT Anti-HCV	Abbott	CMIA (Automated)
AxSYM Anti-HCV	Abbott	MEIA (Automated)
OraQuick HCV Rapid Antibody Test	OraSure	Immunochromatographic (Manual)
Ortho HCV Version 3.0 EIA	Ortho	EIA (Manual)
VITROS Anti-HCV	Ortho	CIA (Automated)
Anti-HCV = HCV antibody; EIA = enzyme immunoassay; CIA = chemiluminescent immunoassay; MEIA = microparticle enzyme immunoassay; CMIA = chemiluminescent microparticle immunoassay		
Table prepared by Saleem Kamili, PhD, Centers for Disease Control and Prevention.		

Testing and Linkage to Care Table 2. Measures to Prevent Transmission of HCV

Persons with HCV infection should be counseled to avoid sharing toothbrushes and dental or shaving equipment, and be cautioned to cover any bleeding wound to prevent the possibility of others coming into contact with their blood.

Persons should be counseled to stop using illicit drugs and enter substance abuse treatment. Those who continue to inject drugs should be counseled to avoid reusing or sharing syringes, needles, water, cotton, and other drug preparation equipment; use new sterile syringes and filters and disinfected cookers; clean the injection site with a new alcohol swab; and dispose of syringes and needles after one use in a safe, puncture-proof container.

Persons with HCV infection should be advised not to donate blood and to discuss HCV serostatus prior to donation of body organs, other tissue, or semen.

Persons with HIV infection and those with multiple sexual partners or sexually transmitted infections should be encouraged to use barrier precautions to prevent sexual transmission. Other persons with HCV infection should be counseled that the risk of sexual transmission is low and may not warrant barrier protection.

Household surfaces and implements contaminated with visible blood from an HCV-infected person should be cleaned using a dilution of 1 part household bleach to 9 parts water. Gloves should be worn when cleaning up blood spills.

Testing and Linkage to Care Table 3: Common Barriers to HCV Treatment and Potential Strategies

Barrier	Strategy
<p>Contraindications to treatment (eg, comorbidities, substance abuse, and psychiatric disorders)</p>	<p>Counseling and education</p> <p>Referral to services (eg, psychiatry and opioid substitution therapy)</p> <p>Optimize treatment with simpler and less toxic regimens</p>
<p>Competing priority and loss to follow-up</p>	<p>Conduct counseling and education</p> <p>Engage case managers and patient navigators (HIV model)</p> <p>Co-localize services (eg, primary care, medical homes, and drug treatment)</p>
<p>Long treatment duration and adverse effects</p>	<p>Optimize treatment with simpler and better tolerated regimens</p> <p>Education and monitoring</p> <p>Directly observed therapy (tuberculosis model)</p>
<p>Lack of access to treatment (high cost, lack of insurance, geographic distance, and lack of availability of specialists)</p>	<p>Leverage expansion of coverage through the Patient Protection and Affordable Care Act</p> <p>Participate in models of care involving close collaboration between primary care practitioners and specialists</p>

Pharmaceutical patient assistance programs

Co-localize services (primary care, medical homes, drug treatment)


Lack of practitioner expertise

Collaboration with specialists (eg, via Project ECHO-like models and telemedicine)

Develop accessible and clear HCV treatment guidelines

Develop electronic health record performance measures and clinical decision support tools (eg, pop-up reminders and standing orders)

Testing and Linkage to Care Figure 1. CDC Recommended Testing Sequence for Identifying Current HCV Infection



* For persons who might have been exposed to HCV within the past 6 months, testing for HCV RNA or follow-up testing for HCV antibody should be performed. For persons who are immunocompromised, testing for HCV RNA should be performed.

† To differentiate past, resolved HCV infection from biologic false positivity for HCV antibody, testing with another HCV antibody assay can be considered. Repeat HCV RNA testing if the person tested is suspected to have had HCV exposure within the past 6 months or has clinical evidence of HCV disease, or if there is concern regarding the handling or storage of the test specimen.

Adapted from Centers for Disease Control and Prevention (CDC), 2013. ([Centers for Disease Control and Prevention \[CDC\], 2013](#))

WHEN AND IN WHOM TO INITIATE HCV THERAPY

Successful hepatitis C treatment results in sustained virologic response (SVR), which is tantamount to virologic cure, and as such, is expected to benefit nearly all chronically infected persons. Evidence clearly supports treatment in all HCV-infected persons, except those with limited life expectancy (less than 12 months) due to non–liver-related comorbid conditions (See sections on [HIV/HCV coinfection](#), [cirrhosis](#), [liver transplantation](#), and [renal impairment](#)). Urgent initiation of treatment is recommended for some patients, such as those with advanced fibrosis or compensated cirrhosis (see [Table 1](#)).

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

A summary of recommendations for When and in Whom to Initiate HCV Therapy is found in the [BOX](#).

Goal of treatment

The goal of treatment of HCV-infected persons is to reduce all-cause mortality and liver-related health adverse consequences, including end-stage liver disease and hepatocellular carcinoma, by the achievement of virologic cure as evidenced by an SVR.

Rating: Class I, Level A

Clinical Benefit of Cure

The proximate goal of HCV therapy is SVR (virologic cure), defined as the continued absence of detectable HCV RNA at least 12 weeks after completion of therapy. SVR is a marker for cure of HCV infection and has been shown to be durable in large prospective studies in more than 99% of patients followed up for 5 years or more. ([Swain, 2010](#)); ([Manns, 2013](#)) Patients in whom an SVR is achieved have HCV antibodies, but no longer have detectable HCV RNA serum, liver tissue, or mononuclear cells, and achieve substantial liver histology improvement. ([Marcellin, 1997](#)); ([Coppola, 2013](#)); ([Garcia-Bengochea, 1999](#)) Assessment of viral response, including documentation of SVR, requires use of US Food and Drug Administration (FDA)-approved quantitative or qualitative nucleic acid test (NAT) with a detection level of 25 IU/mL or lower.

Patients who are cured of their HCV infection experience numerous health benefits, including a decrease in liver inflammation as reflected by improved aminotransferase (ie, alanine aminotransferase [ALT], aspartate aminotransferase[AST]) levels and a reduction in the rate of progression of liver fibrosis. ([Poynard, 2002b](#)) Of 3010 treatment-naïve HCV-infected patients with pretreatment and posttreatment biopsies from 4 randomized trials of 10 different IFN-based regimens (biopsies separated by a mean of 20 months), 39% to 73% of patients achieving an SVR had improvement in liver fibrosis and necrosis ([Poynard, 2002b](#)) and cirrhosis resolved in half of the cases. Portal hypertension, splenomegaly, and other clinical manifestations of advanced liver disease also improved. Among HCV-infected persons, SVR is

associated with a more than 70% reduction in the risk of liver cancer (hepatocellular carcinoma) and a 90% reduction in the risk of liver-related mortality and liver transplantation. ([Morgan, 2013](#)); ([van der Meer, 2012](#)); ([Veldt, 2007](#))

Cure of HCV infection also reduces symptoms and mortality from severe extrahepatic manifestations, including cryoglobulinemic vasculitis, a condition affecting 10% to 15% of HCV-infected patients. ([Fabrizi, 2013](#)); ([Landau, 2010](#)) HCV-infected persons with non-Hodgkin lymphoma and other lymphoproliferative disorders achieve complete or partial remission in up to 75% of cases following successful antiviral therapy for HCV infection. ([Gisbert, 2005](#)); ([Takahashi, 2012](#)); ([Svoboda, 2005](#)); ([Mazzaro, 2002](#)); ([Hermine, 2002](#)) These reductions in disease severity contribute to dramatic reductions in all-cause mortality. ([van der Meer, 2012](#)); ([Backus, 2011](#)) Lastly, patients achieving SVR have substantially improved quality of life, which includes physical, emotional, and social health. ([Neary, 1999](#)); ([Younossi, 2013](#)) Because of the myriad benefits associated with successful HCV treatment, clinicians should treat HCV-infected patients with antiviral therapy with the goal of achieving an SVR, preferably early in the course of their chronic HCV infection before the development of severe liver disease and other complications.

Recommendations for when and in whom to initiate treatment

Treatment is recommended for patients with chronic HCV infection.

Rating: Class I, Level A

Immediate treatment is assigned the highest priority for those patients with advanced fibrosis (Metavir F3), those with compensated cirrhosis (Metavir F4), liver transplant recipients, and patients with severe extrahepatic hepatitis C ([Table 1](#)).

Based on available resources, **immediate** treatment should be prioritized as necessary so that patients at high risk for liver-related complications and severe extrahepatic hepatitis C complications are given high priority ([Table 1](#)).

Ratings: See tables

The most immediate and high-impact benefits of SVR will be realized by populations that are at the highest risk for liver-related complications due to progressive liver disease (Metavir F3 or F4) and transplant recipients or those with clinically severe extrahepatic manifestations ([Table 1](#)).

Other populations at high risk for liver disease progression (Metavir F2) or with substantial extrahepatic manifestations ([Table 1](#)) are also expected to garner appreciable benefits, although the time course for realizing these benefits may be more protracted.

SVR will also remove the risk of further transmission. Treatment of individuals at high risk to transmit HCV to others ([Table 2](#)) may yield long-term future benefits from decreased transmission and a potential decrease in HCV disease prevalence.

When and in Whom to Initiate HCV Therapy Table 1. Settings of Liver-Related Complications and Extrahepatic Disease in Which HCV Treatment is Most Likely to Provide the Most Immediate and Impactful Benefits

Highest Priority for Treatment Owing to Highest Risk for Severe Complications

Advanced fibrosis (Metavir F3) or compensated cirrhosis (Metavir F4)

Rating: Class I, Level A

Organ transplant

Rating: Class I, Level B

Type 2 or 3 essential mixed cryoglobulinemia with end-organ manifestations (eg, vasculitis)

Rating: Class I, Level B

Proteinuria, nephrotic syndrome, or membranoproliferative glomerulonephritis

Rating: Class IIa, Level B

High Priority for Treatment Owing to High Risk for Complications

Fibrosis (Metavir F2)

Rating: Class I, level B

HIV-1 coinfection

Rating: Class I, Level B

Hepatitis B virus (HBV) coinfection

Rating: Class IIa, Level C

Other coexistent liver disease (eg, [NASH])

Rating: Class IIa, Level C

Debilitating fatigue

Rating: Class IIa, Level B

Type 2 Diabetes mellitus (insulin resistant)

Rating: Class IIa, Level B

Porphyria cutanea tarda

Rating: Class IIb, Level C

Ratings refer to the strength and level of evidence with regard to benefits of treatment in these settings.

When and in Whom to Initiate HCV Therapy Table 2. Persons **At Elevated Risk of HCV Transmission* and in Whom HCV Treatment May Yield Transmission Reduction Benefits**

Men who have sex with men (MSM) with high-risk sexual practices

Active injection drug users

Incarcerated persons

Persons on long-term hemodialysis

HCV-infected women of child-bearing potential wishing to get pregnant

Rating: Class IIa, Level C

*Patients at **substantial** risk of transmitting HCV should be counseled on ways to decrease transmission and minimize the risk of reinfection.

Persons with Advanced Liver Disease

For persons with advanced liver disease (Metavir stage F3 or F4), the risk of developing complications of liver disease such as hepatic decompensation ([child Turcotte Pugh \[CTP\] Class B or C \[Methods Table 3\]](#)) or hepatocellular carcinoma is substantial and may occur in a relatively short timeframe. A large prospective study of patients with cirrhosis resulting from HCV infection examined the risk of decompensation, including hepatocellular carcinoma, ascites, jaundice, bleeding, and encephalopathy, and found that the overall annual incidence rate was 3.9%. ([Sangiovanni, 2006](#)) The National Institutes of Health (NIH)-sponsored HALT-C study included a group of 220 patients with cirrhosis resulting from HCV infection who were observed for approximately 8 years. A primary outcome of death, hepatic decompensation, hepatocellular carcinoma, or advance in CTP score of 2 or higher occurred at a rate of 7.5% per year. ([Everson, 2006](#)); ([Di Bisceglie, 2008](#)) Patients with a CTP score of 7 or higher experienced a death rate of 10% per year.

Numerous studies have demonstrated that hepatitis C therapy and the achievement of an SVR in this population results in dramatic decreases in hepatic decompensation events, hepatocellular carcinoma, and liver-related mortality. ([Morgan, 2013](#)); ([van der Meer, 2012](#)); ([Backus, 2011](#)); ([Dienstag, 2011](#)); ([Berenguer, 2009](#)); ([Mira, 2013](#)) In the HALT-C study, patients with advanced fibrosis secondary to HCV infection who achieved an SVR, compared with patients with similarly advanced liver fibrosis who did not achieve an SVR, had a decreased need for liver transplantation (hazard ratio [HR], .17, 95% confidence interval [CI], .06–.46), development of liver-related morbidity and mortality (HR, .15, 95% CI, .06–.38) and hepatocellular carcinoma (HR, .19, 95% CI, .04–.80). ([Dienstag, 2011](#)) Based on these considerations, prompt HCV treatment is recommended for persons with advanced liver disease unless contraindicated (eg, hypersensitivity) or substantial nonhepatic life-limiting comorbidities are present. Importantly, persons with advanced liver disease also require long-term follow-up and hepatocellular carcinoma (HCC) surveillance regardless of treatment outcome (see [Monitoring Section](#)).

Given the clinical complexity and the need for close monitoring, patients with advanced liver disease that has already decompensated ([GTR Class B or C \(Method Table 1\)](#)) should, in general, be treated by physicians with experience in treating HCV in conjunction with a liver transplant center, if possible.

Persons Who Have Undergone Liver Transplantation

In HCV-infected individuals, HCV infection of the allograft occurs universally in patients in whom viral replication is ongoing at the time of transplantation. Histologic features of hepatitis develop in about 75% of recipients in the first 6 months following liver transplantation. ([Neumann, 2004](#)) By the fifth postoperative year, untreated, up to 30% have progressed to cirrhosis. ([Neumann, 2004](#)); ([Charlton, 1998](#)) A small proportion of patients (4%-7%) develop an accelerated course of liver injury (cholestatic hepatitis C, associated with very high levels of viremia) with subsequent rapid allograft failure. Recurrence of HCV infection post-transplantation has led to shorter period of graft survival for recipients with HCV infection than for recipients who undergo liver transplantation for other indications. ([Forman, 2002](#))

Effective antiviral therapy pre-transplantation resulting in a SVR (virologic cure) prevents HCV recurrence post-transplantation. ([Everson, 2003](#)) In addition, complete HCV viral suppression prior to transplantation prevents recurrent HCV infection of the graft in the majority of cases. ([Forns, 2004](#)); ([Everson, 2005](#)) Preliminary data from a study of patients with complications of cirrhosis secondary to HCV infection who are wait-listed for liver transplantation that included patients with MELD scores up to 14 and CPT scores up to 8 found that treatment with sofosbuvir and weight-based ribavirin (RBV) for up to 48 weeks was well tolerated and was associated with an overall post-transplant SVR of 69%. ([Curry, 2013b](#)) Post-transplant SVR was near ubiquitous among patients who had undetectable HCV RNA for 28 days or longer prior to transplantation.

Treatment of established HCV infection post-transplantation also yields substantial improvements in patient and graft survival. ([Berenguer, 2008](#)); ([Picciotto, 2007](#)) The availability of effective IFN-free HCV treatment regimens has addressed the major hurdles to treating HCV recurrence post-transplantation: poor tolerability and efficacy. In a multicenter, open-label study evaluating the ability of sofosbuvir plus RBV to induce virologic suppression in 40 post-liver-transplant patients with compensated recurrence of HCV infection, daily sofosbuvir and RBV for 24 weeks achieved an SVR at 12 weeks (SVR12) in 70%. ([Charlton, 2014b](#)) No deaths, graft losses, or episodes of rejection occurred. Six patients had serious adverse events, all of which were considered unrelated to study treatment. There were no drug interactions with sofosbuvir and any of the concomitant immunosuppressive agents reported. In contrast, treatment with sofosbuvir plus RBV with or without PEG in 64 patients with severe, decompensated cirrhosis resulting from recurrence of HCV infection following liver transplantation was associated with an overall SVR12 of 57% and a mortality rate of 25%. ([Forns, 2013c](#)) On an intent-to-treat basis, treatment was associated with clinical improvement in 64% and stable disease in 11% of patients.

Persons at Greater Risk of Rapidly Progressive Fibrosis and Cirrhosis

Fibrosis progression is variable across different patient populations as well as within the same individual over time. Many of the components that determine fibrosis progression and development of cirrhosis in an individual are unknown. However, certain factors, such as coinfection with HIV or hepatitis B virus (HBV) and prevalent coexistent liver diseases (eg, nonalcoholic steatohepatitis [NASH]), are well-recognized contributors to accelerated fibrosis progression. Patients with these conditions should be prioritized for hepatitis C therapy.

HIV coinfection. HIV coinfection accelerates fibrosis progression among HCV-infected persons, ([Benhamou, 1999](#)); ([Macias, 2009](#)); ([Konerman, 2014](#)) although control of HIV replication and restoration

of CD4+ cell counts may mitigate this to some extent. ([Benhamou, 2001](#)); ([Bräu, 2006](#)) In the largest paired biopsy study, 282 HIV/HCV-coinfected patients with 435 paired biopsies were prospectively evaluated; ([Konerman, 2014](#)) one-third of patients showed fibrosis progression of at least 1 Metavir stage at a median of 2.5 years. Importantly, 45% of patients with no fibrosis on initial biopsy had progression. Finally, a more rapid progression to death following decompensation, combined with a lack of widespread access to liver transplantation and poor outcomes following transplantation, argue for treatment prioritization in this population, regardless of the current fibrosis stage. ([Pineda, 2005](#)); ([Merchante, 2006](#)); ([Terrault, 2012](#))

HBV coinfection and other coexistent liver diseases. The prevalence of HBV/HCV coinfection is estimated at 1.4% in the United States and 5% to 10% globally. ([Tyson, 2013](#)); ([Chu, 2008](#)) Persons with HBV/HCV coinfection and detectable viremia of both viruses are at increased risk for disease progression, decompensated liver disease, and the development of hepatocellular carcinoma.

HBV/HCV coinfecting individuals are susceptible to a process called viral interference wherein one virus may interfere with the replication of the other virus. Thus, when treating one or both viruses with antiviral drugs, it is prudent to periodically retest HBV DNA and HCV RNA levels during and after therapy, particularly if only one of the viruses is being treated at a time. Treatment of HCV in such cases utilizes the same genotype-specific regimens as are recommended for HCV mono-infection (**see [Treatment Section](#)**). HBV infections in such cases should be treated as recommended for HBV mono-infection. ([Lok, 2009](#))

Persons with other chronic liver diseases who have coincident chronic HCV infection should be considered for hepatitis C therapy, given the potential for rapid progression of liver disease. An IFN-free regimen is generally preferred for immune-mediated liver diseases such as autoimmune hepatitis because of the potential for IFN to exacerbate them.

Persons with Severe Extrahepatic Manifestations of Chronic HCV Infection

Chronic hepatitis C is associated with a syndrome of cryoglobulinemia and an immune complex and lymphoproliferative disorder that produces arthralgias, fatigue, palpable purpura, renal disease (eg, membranoproliferative glomerulonephritis), neurologic disease (eg, peripheral neuropathy, central nervous system vasculitis), and reduced complement levels. ([Agnello, 1992](#)) Because patients with chronic hepatitis C frequently have laboratory evidence of cryoglobulins (more than 50% in some series), antiviral treatment should be prioritized for those with the syndrome of cryoglobulinemia and symptoms or objective evidence of end-organ manifestations. IFN-based regimens can produce clinical remission; however, the adverse effects of IFN may mimic manifestations of cryoglobulinemia. ([Saadoun, 2014](#)) Although clinical data are not yet available, the use of IFN-free direct-acting antiviral (DAA) regimens is an attractive alternative for these patients. Organ-threatening disease (eg, severe neuropathy, renal failure, digital ischemia), in addition to the antiviral HCV therapy, should also be treated more acutely with immunosuppressive agents or plasmapheresis to clear immune complexes.

Glomerular disease results from deposition of HCV-related immune complexes in the glomeruli. ([Johnson, 1993](#)) Successful treatment of HCV using IFN-based regimens can reverse proteinuria and the nephrotic syndrome, but usually does not fully ameliorate azotemia. ([Johnson, 1994](#)) No clinical trial data are yet available using IFN-free regimens, but the high rates of SVR (virologic cure) using antiviral therapy support their use in management of hepatitis C-related renal disease and cryoglobulinemia.

The relationship between chronic hepatitis C and diabetes (most notably type 2 diabetes and insulin resistance) is complex and incompletely understood. The prevalence and incidence of diabetes is increased in the context of hepatitis C. ([White, 2008](#)) In the United States, type 2 diabetes occurs more

frequently in HCV-infected patients with a more than 3-fold greater risk in persons over 40 years of age. ([Mehta, 2000](#)) The positive correlation between quantity of plasma HCV RNA and established markers of insulin resistance confirms this relationship. ([Yoneda, 2007](#)) Insulin resistance and type 2 diabetes are independent predictors of a more rapid progression of liver fibrosis and impaired response to IFN-based therapy. ([Petta, 2008](#)) Patients with type 2 diabetes and insulin resistance also are at increased risk for hepatocellular carcinoma. ([Hung, 2010](#))

Successful antiviral treatment has been associated with improved markers of insulin resistance and greatly reduced incidence of new onset of type 2 diabetes and insulin resistance in HCV-infected patients. ([Arase, 2009](#)) Most recently, antiviral therapy for HCV infection has been shown to improve clinical outcomes related to diabetes. In a large prospective cohort from Taiwan, the incidence of ESRD, ischemic stroke, and acute coronary syndrome were greatly reduced in HCV-infected patients with diabetes who received antiviral therapy compared with untreated, matched controls. ([Hsu, 2014](#)) Therefore, antiviral therapy in patients with pre-diabetes who have hepatitis C may prevent progression to diabetes and reduce renal and cardiovascular complications in hepatitis C patients with established diabetes.

Persons with Other Extrahepatic Manifestations of HCV Infection

In patients with chronic hepatitis C, fatigue is the most frequently reported symptom and has a major effect on quality of life and activity level evidenced by numerous measures of impaired quality of life. ([Foster, 1998](#)) The presence and severity of fatigue appears to correlate poorly with disease activity, although it may be more common and severe in HCV-infected individuals with cirrhosis. ([Poynard, 2002a](#)) Despite difficulties in separating fatigue symptoms associated with hepatitis C from those associated with other concurrent conditions (eg, anemia, depression), numerous studies have reported a reduction in fatigue after cure of HCV. ([Bonkovsky, 2007](#)) In the Virahep-C study, 401 HCV patients were evaluated for fatigue prior to treatment and after therapy using validated scales assessing presence and severity of fatigue. ([Sarkar, 2012](#)) At baseline, 52% of patients reported having fatigue, which was more frequent and severe in patients with cirrhosis than in those without cirrhosis. Achieving an SVR was associated with a substantial decrease in frequency and severity of fatigue. A recent analysis of 413 patients who achieved an SVR12 from the NEUTRINO and FUSION trials treated with a sofosbuvir-containing regimen demonstrated improvement in patient fatigue (present in 12%) from the pretreatment level. ([Younossi, 2014](#)) After achieving an SVR12, participants had marked improvement in fatigue over their pretreatment scores using 3 separate validated questionnaires.

The reported prevalence of HCV infection in patients with porphyria cutanea tarda approximates 50% and occurs disproportionately in those with cirrhosis. ([Gisbert, 2003](#)) The treatment of choice for active porphyria cutanea tarda is iron reduction by phlebotomy and maintenance of a mildly iron-reduced state without anemia. However, although improvement of porphyria cutanea tarda during HCV treatment with IFN has frequently been described ([Takikawa, 1995](#)), there are currently insufficient data to determine whether treating HCV infection with DAAs and achieving an SVR improves porphyria cutanea tarda.

Lichen planus is characterized by pruritic papules involving mucous membranes, hair, and nails. Antibodies to HCV are present in 10% to 40% of patients with lichen planus, but the causal link with chronic infection is not established. Resolution of lichen planus has been reported with IFN-based regimens, but there have also been reports of exacerbation of lichen planus with these treatments. Although it is unknown whether DAAs will have more success against lichen planus, treatment with IFN-free regimens would appear to be a more advisable approach to addressing this disorder. ([Gumber, 1995](#))

Persons **With Elevated Risk of Transmitting HCV**

Persons who have successfully achieved an SVR (virologic cure) no longer transmit the virus to others. As such, successful treatment benefits public health. Several health models have shown that even modest increases in successful treatment of HCV infection among persons who inject drugs can decrease prevalence and incidence. ([Martin, 2013a](#)); ([Durier, 2012](#)); ([Martin, 2013b](#)); ([Hellard, 2012](#)) Models developed to estimate the impact of HCV testing and treatment on the burden of hepatitis C at a country level reveal that large decreases in HCV prevalence and incidence are possible as more persons are successfully treated. ([Wedemeyer, 2014](#)) There are also benefits between couples and among families to eradicating HCV infection and thus eliminating the perception that an individual might be contagious. In addition, mother-to-child transmission of HCV does not occur if the woman is not viremic, providing an additional benefit of curing a woman before she becomes pregnant. ([Thomas, 1998](#)) However, the safety and efficacy of treating women who are already pregnant to prevent transmission to the fetus have not yet been established, and thus treatment is not recommended for pregnant women.

Successful treatment of HCV-infected persons at greatest risk for transmission represents a formidable tool to help stop HCV transmission in those who continue to engage in high-risk behaviors. To guide implementation of hepatitis C treatment as a prevention strategy, studies are needed to define the best candidates for treatment to stop transmission; the additional interventions needed to maximize the benefits of HCV treatment (eg, preventing reinfection), and the cost effectiveness of the strategies when used in the target populations.

Persons who inject drugs. Injection drug use is the most common risk factor for HCV infection in the United States and Europe with an HCV seroprevalence from 10% to 70%; ([Amon, 2008](#)); ([Nelson, 2011](#)) injection drug use also accounts for the majority of new infections (approximately 70%) and is the key driving force in perpetuation of the epidemic. Given these facts, and the absence of an effective vaccine against HCV, testing and linkage to care combined with treatment of HCV infection using potent IFN-free regimens has the potential to dramatically decrease HCV incidence and prevalence. ([Martin, 2013b](#)) However, treatment-based strategies to prevent HCV transmission have yet to be studied, including how to integrate hepatitis C treatment with other risk reduction strategies including opiate substitution therapy and needle and syringe exchange programs. ([Martin, 2013a](#))

Studies of IFN-containing treatments in persons who inject drugs have shown comparable adherence and efficacy rates to patients who do not use injection drugs. A recent meta-analysis of treatment in active or recent injection drug users with PEG with or without RBV showed SVR rates of 37% and 67% for genotype 1 or 4 and 2 or 3, respectively. ([Aspinall, 2013](#)) As shorter, better-tolerated, more-efficacious IFN-free therapies are introduced, these SVR rates are expected to improve. Importantly, the rate of reinfection in this population is lower (2.4/100 person years of observation) than incident infection in the injection drug user population in general (6.1-27.2/100 person years); though reinfection increases with active or ongoing injection drug use (6.44/100 person years) and available data are limited in follow-up duration. ([Aspinall, 2013](#)); ([Grady, 2013](#))

Ideally, treatment of HCV-infected persons who inject drugs should be delivered in a multidisciplinary care setting with services to reduce the risk of reinfection and for management of the common social and psychiatric comorbidities in this population. Regardless of the treatment setting, recent and active injection drug use should not be seen as an absolute contraindication to HCV therapy. Scale up of HCV treatment in persons who inject drugs is necessary to positively impact the HCV epidemic in the United States and globally.

HIV-infected men who have sex with men (MSM) with high-risk sexual practices. Over the past

decade a dramatic increase in incident HCV infections has been demonstrated in several US cities among HIV-infected MSM who did not report injection drug use as a risk factor. ([van de Laar, 2010](#)) Recognition and treatment of HCV (including acute infection) in this population may represent an important step in preventing subsequent infections. As with persons who inject drugs, HIV-infected MSM with ongoing high-risk sexual practices should be treated for their HCV infection in conjunction with continued education on risk reduction strategies. In particular, safer sex strategies should be emphasized given the high rates of reinfection after SVR, which may approach 30% over 2 years, in HIV-infected MSM with acute HCV infection. ([Lambers, 2011](#))

Incarcerated persons. Among incarcerated individuals, the HCV seroprevalence ranges from 30% to 60% ([Post, 2013](#)) and an acute infection rate of about 1%. ([Larney, 2013](#)) Screening for HCV is relatively uncommon in state prison systems. Treatment uptake has been limited in part because of the toxic effects and long treatment duration of older IFN-based therapies as well as concerns about cost. ([Spaulding, 2006](#)) In particular, truncation of treatment owing to release from prison during therapy has been cited as a major limitation to widespread, effective HCV treatment in correctional facilities. ([Post, 2013](#)); ([Chew, 2009](#)) Shorter (12-week to 24-week) HCV therapies reduce duration of stay-related barriers to HCV treatment in prisons. Likewise, the improved safety of newer, all-oral regimens diminishes toxicity concerns. Coordinated treatment efforts within prison systems would likely rapidly decrease the prevalence of HCV infection in this at-risk population, although research is needed in this area.

Persons on hemodialysis. The prevalence of HCV infection is markedly elevated in persons on hemodialysis and ranged from 2.6 to 22.9% in a large multinational study. ([Fissell, 2004](#)) Studies in the United States found a similarly elevated prevalence of 7.8% to 8.9%. ([Centers for Disease Control and Prevention, 2001](#)); ([Finelli, 2005](#)) Importantly, the seroprevalence of HCV was found to increase with time on dialysis suggesting that nosocomial transmission, among other risk factors, plays a role in HCV acquisition in these patients. ([Fissell, 2004](#)) Improved education and strict adherence to universal precautions can drastically reduce nosocomial HCV transmission risks for persons on hemodialysis, ([Jadoul, 1998](#)) but clearance of HCV viremia through treatment-induced SVR eliminates the potential for transmission.

HCV-infected persons on hemodialysis have a decreased quality of life and increased mortality compared with persons on hemodialysis without HCV infection. ([Fabrizi, 2002](#)); ([Fabrizi, 2007](#)); ([Fabrizi, 2009](#)) HCV infection in this population also has a deleterious impact on kidney transplantation outcomes with decreased patient and graft survival. ([Fabrizi, 2014](#)) The increased risk for nosocomial transmission combined with the substantial clinical impact of HCV infection in those on hemodialysis suggest that this group should also be prioritized for HCV therapy as effective antiviral regimens that can be used in advanced renal failure become available.

For all these populations, the decision to treat should be based on a favorable risk-benefit ratio taking into account the anticipated reduction in transmission(s) versus the likelihood of reinfection.

Populations Unlikely to Benefit from HCV Treatment

Patients with limited life expectancy for whom HCV therapy would not improve symptoms or prognosis do not require treatment. Chronic hepatitis C is associated with a wide range of comorbid conditions. ([Butt, 2011](#)); ([Louie, 2012](#)) Little evidence exists to support initiation of HCV treatment in patients with limited life expectancy (less than 12 months) due to non-liver-related comorbid conditions. For these patients, the benefits of HCV treatment are unlikely to be realized, and palliative care strategies should take precedence. ([Holmes, 2006](#)); ([Maddison, 2011](#))

Recommendations for pretreatment assessment

An assessment of the degree of hepatic fibrosis, using noninvasive testing or liver biopsy, is recommended.

Rating: Class I, Level A

An accurate assessment of fibrosis is vital in assessing the urgency for treatment. The degree of hepatic fibrosis is one of the most robust prognostic factors used to predict disease progression and clinical outcomes. ([Everhart, 2010](#)) Those with substantial fibrosis (defined as Metavir F2 or higher) should be given priority for therapy in an effort to decrease the risk of clinical consequences such as cirrhosis, liver failure, and hepatocellular cancer. However, those with severe fibrosis (Metavir stage F3 and F4) are most in need of immediate therapy. In addition to urgency for antiviral therapy, individuals with severe fibrosis require surveillance monitoring for liver cancer, esophageal varices, and hepatic function. ([Garcia-Tsao, 2007](#)); ([Bruix, 2011](#))

Although liver biopsy is the diagnostic standard, sampling error and observer variability limit test performance, particularly when inadequate sampling occurs. Up to one-third of bilobar biopsies had a difference of at least 1 stage between the lobes. ([Bedossa, 2003](#)) In addition, the test is invasive and minor complications are common, limiting patient and practitioner acceptance. Serious complications such as bleeding, although rare, are well recognized.

Noninvasive tests to stage the degree of fibrosis in patients with chronic HCV infection include models incorporating indirect serum biomarkers (routine tests), direct serum biomarkers (components of the extracellular matrix produced by activated hepatic stellate cells), and vibration-controlled transient liver elastography. No single method is recognized to have high accuracy alone and each test must be interpreted carefully. A recent publication of the Agency for Healthcare Research and Quality found evidence in support of a number of blood tests; however, at best they are only moderately useful for identifying clinically significant fibrosis or cirrhosis. ([Selph, 2014](#))

Vibration-controlled transient liver elastography is a noninvasive way to measure liver stiffness and correlates well with measurement of substantial fibrosis or cirrhosis in patients with chronic HCV infection. A cutoff value of 8.7 kPa correlates with Metavir F2 or higher fibrosis stage; greater than 9.5 kPa with F3; and 14.5 or higher kPa with F4 or cirrhosis. The measurement range does overlap between stages. ([Ziol, 2005](#))

The most efficient approach to fibrosis assessment is to combine direct biomarkers and vibration-controlled transient liver elastography. ([Boursier, 2012](#)) A biopsy should be considered for any patient who has discordant results between the 2 modalities that would affect clinical decision making. For example, 1 shows cirrhosis and the other does not. The need for liver biopsy with this approach is markedly reduced.

Alternatively, if direct biomarkers or vibration-controlled transient liver elastography are not available, the aspartate aminotransferase-to-platelet ratio index (APRI) or fibrosis-4 index (FIB-4) can help identify those most likely to have F3 or F4 fibrosis stage. ([Sebastiani, 2009](#)); ([Castera, 2010](#)); ([Chou, 2013b](#)) An APRI above 2.0 or FIB-4 above 3.25 has a high specificity for advanced fibrosis or cirrhosis, although neither test

is sensitive enough to rule out substantial fibrosis if values are below these thresholds. ([Chou, 2013b](#)) Biopsy should be considered in those in whom more accurate fibrosis staging would impact treatment decisions.

Individuals with clinically evident cirrhosis do not require additional staging (biopsy or noninvasive assessment).

Recommendation for repeat liver disease assessment

Ongoing assessment of liver disease is recommended for persons in whom therapy is deferred.

Rating: Class I, Level C

When therapy is deferred, it is especially important to monitor liver disease in these patients. Among individuals with less-advanced stages of fibrosis, fibrosis progression over time will help determine the urgency of subsequent antiviral therapy. Fibrosis progression varies markedly between individuals based on host, environmental, and viral factors ([Table 3](#)). ([Feld, 2006](#)) Fibrosis may not progress linearly. Some individuals (often those who are aged 50 years or older) may progress slowly for many years followed by an acceleration of fibrosis progression. Others may never develop substantial liver fibrosis despite longstanding infection. The presence of existing fibrosis is a strong risk factor for future fibrosis progression. Fibrosis results from chronic hepatic necroinflammation and thus a higher activity grade on liver biopsy and higher serum transaminase values is associated with more rapid fibrosis progression. ([Ghany, 2003](#)) However, even patients with normal alanine aminotransferase (ALT) levels may develop substantial liver fibrosis over time. ([Pradat, 2002](#)); ([Nutt, 2000](#))

Host factors associated with more rapid fibrosis progression include male sex, longer duration of infection, and older age at the time of infection. ([Poynard, 2001](#)) Many patients have concomitant nonalcoholic fatty liver disease, and the presence of hepatic steatosis with or without steatohepatitis on liver biopsy as well as elevated body mass index and insulin resistance are associated with fibrosis progression, as is iron overload. ([Konerman, 2014](#)); ([Everhart, 2009](#)) Chronic alcohol use is an important risk factor because alcohol consumption has been associated with more rapid fibrosis progression. ([Feld, 2006](#)) A safe amount of alcohol consumption has not been established. Cigarette smoking may also lead to more rapid fibrosis progression.

Immunosuppression leads to more rapid fibrosis progression, particularly HIV coinfection and solid organ transplantation. ([Macias, 2009](#)); ([Konerman, 2014](#)); ([Berenguer, 2013](#)) Therefore, immunocompromised patients should be prioritized for antiviral therapy even if they have mild liver fibrosis at presentation.

The level of virus in the serum (HCV RNA) is not highly correlated with the stage of disease (degree of inflammation or fibrosis). Available data suggest that fibrosis progression occurs most rapidly in patients with genotype 3 HCV infection. ([Kanwal, 2014](#)) ([Bochud, 2009](#)) Aside from coinfections with HBV or HIV, no other viral factors are consistently associated with disease progression.

Although an ideal interval for assessment has not been established, annual evaluation is appropriate to

discuss modifiable risk factors and update testing for hepatic function and markers for disease progression. For all individuals with advanced fibrosis, liver cancer screening dictates a minimum of every 6 months evaluation.

When and in Whom to Initiate HCV Therapy Table 3. Factors Associated with Accelerated Fibrosis Progression

Host	Viral
<p>Non-Modifiable Fibrosis stage Inflammation grade Older age at time of infection Male sex Organ transplant</p> <p>Modifiable Alcohol consumption Nonalcoholic fatty liver disease Obesity Insulin resistance</p>	<p>Genotype 3 Coinfection with hepatitis B virus (HBV) or HIV</p>

When and in Whom to Initiate HCV Therapy Box. Summary of Recommendations for When and in Whom to Initiate HCV Therapy

When and in Whom to Initiate HCV Therapy Box. Summary of Recommendations for When and in Whom to Initiate HCV Therapy

Goal of treatment

The goal of treatment of HCV-infected persons is to reduce all-cause mortality and liver-related health adverse consequences, including end-stage liver disease and hepatocellular carcinoma, by the achievement of virologic cure as evidenced by an SVR.

Rating: Class I, Level A

Recommendations for when and in whom to initiate treatment

Treatment is recommended for patients with chronic HCV infection.

Rating: Class I, Level A

Immediate treatment is assigned the highest priority for those patients with advanced fibrosis (Metavir F3), those with compensated cirrhosis (Metavir F4), liver transplant recipients, and patients with severe extrahepatic hepatitis C ([Table 1](#)).

Based on available resources, **immediate** treatment should be prioritized as necessary so that patients at high risk for liver-related complications and severe extrahepatic hepatitis C complications are given high priority ([Table 1](#)).

Ratings: See tables

Recommendations for pretreatment assessment

An assessment of the degree of hepatic fibrosis, using noninvasive testing or liver

biopsy, is recommended.

Rating: Class I, Level A

Recommendation for repeat liver disease assessment

Ongoing assessment of liver disease is recommended for persons in whom therapy is deferred.

Rating: Class I, Level C

Table 1. Settings of Liver-Related Complications and Extrahepatic Disease in Which HCV Treatment is Most Likely to Provide the Most Immediate and Impactful Benefits

Highest Priority for Treatment Owing to Highest Risk for Severe Complications

Advanced fibrosis (Metavir F3) or compensated cirrhosis (Metavir F4)

Rating: Class I, Level A

Organ transplant

Rating: Class I, Level B

Type 2 or 3 essential mixed cryoglobulinemia with end-organ manifestations (eg, vasculitis)

Rating: Class I, Level B

Proteinuria, nephrotic syndrome, or membranoproliferative glomerulonephritis

Rating: Class IIa, Level B

High Priority for Treatment Owing to High Risk for Complications

Fibrosis (Metavir F2)

Rating: Class I, level B

HIV-1 coinfection

Rating: Class I, Level B

HBV coinfection

Rating: Class IIa, Level C

Other coexistent liver disease (eg, NASH)

Rating: Class IIa, Level C

Debilitating fatigue

Rating: Class IIa, Level B

Type 2 Diabetes mellitus (insulin resistant)

Rating: Class IIa, Level B

Porphyria cutanea tarda

Rating: Class IIb, Level C

Ratings refer to the strength and level of evidence with regard to benefits of treatment in these settings.

Table 2. Persons At Elevated Risk of HCV Transmission* and in Whom HCV Treatment May Yield Transmission Reduction Benefits

Men who have sex with men (MSM) with high-risk sexual practices

Active injection drug users

Incarcerated persons

Persons on long-term hemodialysis

HCV-infected women of child-bearing potential wishing to get pregnant

Rating: Class IIa, Level C

*Patients at substantial risk of transmitting HCV should be counseled on ways to decrease transmission and minimize the risk of reinfection.

Table 3. Factors Associated with Accelerated Fibrosis Progression

Host	Viral
Non-Modifiable Fibrosis stage Inflammation grade Older age at time of infection Male sex Organ transplant	Genotype 3 Coinfection with HBV or HIV
Modifiable Alcohol consumption Nonalcoholic fatty liver disease Obesity Insulin resistance	

INITIAL TREATMENT OF HCV INFECTION

/*-->*/

Initial Treatment of HCV Infection includes patients with chronic hepatitis C infection who have not been previously treated with IFN, PEG-IFN, RBV, or any HCV direct-acting antiviral (DAA) agent, whether experimental, investigational, or US Food and Drug Administration (FDA) approved.

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

A summary of recommendations for initial treatment is found in the [BOX](#).

The level of evidence available to inform the best treatment decision for each patient varies, as does the strength of the recommendation, and is rated accordingly (see [Methods Table 2](#)). In addition, when treatment differs for a particular group, such as those infected with specific HCV genotype or subtype, specific recommendations are given. A regimen is classified as "Recommended" when it is favored for most patients and "Alternative" when optimal in a particular subset of patients in that category. When a treatment is clearly inferior or is deemed harmful, it is classified as "Not Recommended." Unless otherwise indicated, such regimens should not be administered to patients with HCV infection. **Specific considerations of persons with [HIV/HCV coinfection](#), [decompensated cirrhosis \(moderate or severe hepatic impairment; Child Turcotte Pugh \[CTP\] class B or C\)](#), [post-liver transplant HCV infection](#), and those with severe [renal impairment](#) or end-stage renal disease are addressed in other sections of the Guidance.**

When several regimens are offered at the same recommendation level they are listed in alphabetic order. In this case consideration of choice of regimen should be determined based on patient-specific data, including drug interactions. As always, patients receiving antiviral therapy require careful pretreatment assessment for comorbidities that may influence treatment response. All patients should have careful monitoring during treatment, particularly for anemia if RBV is included in the regimen. (See [Monitoring Section](#))

I. Genotype 1

Three highly potent DAA oral combination regimens are recommended for HCV genotype 1–infected patients, although there are differences in the recommended regimens based on the HCV subtype. Patients with HCV genotype 1a tend to have higher relapse rates than patients with HCV genotype 1b with certain regimens. Genotype 1 HCV infection that cannot be subtyped should be treated as genotype 1a infection.

The introduction of DAAs into HCV treatment regimens increased the risk of drug interactions with other concomitant medications used by patients, and now with combinations of DAAs, attention to drug interactions is all the more important (see [Drug Interactions Table](#)). The product prescribing information and other resources (eg, <http://www.hep-druginteractions.org/>) should be referenced regularly to ensure safety when prescribing DAA regimens. In particular, the daily fixed-dose combination of ledipasvir (90 mg) and sofosbuvir (400 mg) (hereafter ledipasvir/sofosbuvir) has a potential interaction with acid-suppressing medications, for example proton pump inhibitors, which may result in decreased absorption of ledipasvir

and lower exposures. Because of over-the-counter access to acid-suppressing medications, a comprehensive assessment of all prescribed and over-the-counter medications is recommended prior to initiating treatment. If possible, acid-suppressing medications should be held prior to and during the HCV treatment period to optimize ledipasvir exposure. For patients in whom interruption of acid suppression is not possible, dosing of acid suppressants is recommended per the prescribing information.

Similarly, the daily fixed-dose combination of paritaprevir (150 mg), ritonavir (100 mg), and ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) (hereafter paritaprevir/ritonavir/ombitasvir plus dasabuvir) has a substantial interaction with the long-acting inhaled beta-adrenoceptor agonist salmeterol, and concurrent administration is not recommended owing to an increased risk of cardiovascular adverse events including QT segment prolongation.

A. Genotype 1a

Three options with similar efficacy in general are recommended for treatment-naive patients with HCV genotype 1a infection (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks* is recommended for treatment-naive patients with HCV genotype 1a infection.

***See text for further detail on [length of treatment](#)**

Rating: Class I, Level A

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks (no cirrhosis) or 24 weeks (cirrhosis) is recommended for treatment-naive patients with HCV genotype 1a infection.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks (no cirrhosis) or 24 weeks (cirrhosis) is recommended for treatment-naive patients with HCV genotype 1a infection.

Rating: Class IIa, Level B

For HCV genotype 1a–infected, treatment-naive patients, there are 3 regimens of comparable efficacy, as outlined above.

Ledipasvir/sofosbuvir was approved by the FDA for the treatment of HCV genotype 1 infection in treatment-naive patients based on 2 registration trials: ION-1 (865 treatment-naive patients; those with cirrhosis were included) and ION-3 (647 treatment-naive patients; those with cirrhosis were excluded). ION-1 investigated length of treatment (12 weeks vs 24 weeks) and the need for RBV. ([Afdhal, 2014a](#)) Sustained virologic response at 12 weeks (SVR12) was 97% to 99% across all arms, with no difference in SVR based on length of treatment, use of RBV, or HCV genotype 1 subtype. Sixteen percent of subjects enrolled were classified as having cirrhosis. There was no difference in SVR12 in those with cirrhosis (97%) versus those without cirrhosis (98%). ION-3 excluded patients with cirrhosis and investigated shortening therapy from 12 weeks to 8 weeks (with or without RBV). ([Kowdley, 2014](#)) SVR12 was 93% to 95% across all arms, with no difference in SVR in the intention-to-treat analysis. However, relapse rates were higher in the 8-week arms (20 of 431) regardless of RBV use compared with the 12-week arm (3 of 216). Post hoc analyses of the 2 RBV-free arms assessed baseline predictors of relapse and identified lower relapse rates in patients receiving 8 weeks of ledipasvir/sofosbuvir who had baseline HCV RNA levels below 6 million IU/mL (2%; 2 of 123), and was the same for patients with similar baseline HCV RNA levels who received 12 weeks (2%; 2 of 131). This analysis was not controlled and thus substantially limits the generalizability of this approach to clinical practice. Shortening treatment to less than 12 weeks should be done with caution and performed at the discretion of the practitioner.

Paritaprevir/ritonavir/ombitasvir plus dasabuvir and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) was approved by the FDA for the treatment of HCV genotype 1a infection in treatment-naive patients based on 3 registration trials: SAPPHERE-I (322 treatment-naive patients with genotype 1a HCV infection; those with cirrhosis were excluded), PEARL-IV (305 treatment-naive patients with genotype 1a; those with cirrhosis were excluded), and TURQUOISE-II (261 treatment-naive and -experienced patients with genotype 1a; those with cirrhosis were included). The SAPPHERE-I trial reported a high SVR12 rate (95.3%) with 12 weeks of paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV. ([Feld, 2014](#)) Overall, virologic failure was higher for patients with HCV genotype 1a (7 of 8) than patients with HCV genotype 1b (1 virologic failure). PEARL-IV was specifically designed to determine the role of paritaprevir/ritonavir/ombitasvir plus dasabuvir with or without weight-based RBV for treatment-naive, HCV genotype 1a-infected patients without cirrhosis. ([Ferenci, 2014](#)) SVR12 was lower in the RBV-free arm than in the RBV-containing arm (90% vs 97%, respectively) owing to higher rates of virologic failure (7.8% vs 2%, respectively), confirming the need for weight-based RBV for patients with HCV genotype 1a. TURQUOISE-II enrolled treatment-naive and -experienced patients (261 patients with HCV genotype 1a) with CTP class A cirrhosis to receive either 12 weeks or 24 weeks of treatment with paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV. Overall, SVR12 rates were 89% in the 12-week arm and 95% in the 24-week arm. ([Ombitasvir/Paritaprevir/Ritonavir prescribing information](#)); ([Poordad, 2014](#)) This difference in SVR12 rate between arms was primarily driven by patients with null response to PEG-IFN and RBV; there was less difference in SVR rates in the patients with cirrhosis who were naive to therapy (92% and 95%, respectively). ([Ombitasvir/Paritaprevir/Ritonavir prescribing information](#)); ([Poordad, 2014](#))

COSMOS was a phase II clinical trial of sofosbuvir plus simeprevir that included treatment-naive and -experienced patients, including those with cirrhosis. ([Lawitz, 2014a](#)) The study enrolled 2 cohorts: cohort 1 included 80 patients with a prior null response to PEG-IFN and RBV with Metavir F0 to F2 fibrosis, and cohort 2 included 87 patients who were either treatment naive or had a prior null response to PEG-IFN and RBV with Metavir stage F3 or F4 fibrosis. Each cohort had 4 arms to address length of treatment (12 weeks vs 24 weeks) and the role of weight-based RBV. Across both cohorts, the SVR12 rate ranged from 79.3% to 100%, with no clear benefit of extended treatment or use of RBV; however, the small size per arm limited the analysis. In a pooled analysis of all RBV-free arms, 95% (20/21) with Metavir stage F0 to F3 fibrosis who received 12 weeks of sofosbuvir and simeprevir achieved SVR12 compared with 86% (6/7)

of those with Metavir stage F4 fibrosis. ([Lawitz, 2014b](#)) Among patients with cirrhosis who received 24 weeks of therapy, the SVR rate was 100% (10/10).

Based on this analysis, the FDA recommends 24 weeks for all patients with cirrhosis, regardless of treatment experience. An ongoing phase III trial (NCT02114151) will provide more definitive information on the response rates of patients with cirrhosis receiving 12 weeks of sofosbuvir and simeprevir without RBV. Until that time, the use of 24 weeks of sofosbuvir and simeprevir for patients with cirrhosis is recommended. Overall, only 6 patients in the COSMOS trial experienced virologic failure (relapse), all had HCV genotype 1a infection, and 4 had virus with the Q80K polymorphism at baseline. When simeprevir is combined with PEG-IFN and RBV, patients with HCV genotype 1a infection and a baseline Q80K polymorphism have higher virologic failure rates than patients with genotype 1b infection or genotype 1a infection without this baseline mutation. This led to the recommendation for baseline Q80K testing for all HCV genotype 1a–infected patients prior to the use of simeprevir with PEG-IFN and RBV. In contrast, the presence of the Q80K polymorphism does not preclude treatment with simeprevir and sofosbuvir, because the SVR rate was high in patients with HCV genotype 1a and the Q80K polymorphism (88%; 51/58). ([Lawitz, 2014b](#)) Given the low but finite failure rates, the role of RBV with simeprevir and sofosbuvir in patients with HCV genotype 1a is unclear. Thus, RBV use may be considered until results from the larger phase III trials can more definitively address these questions (NCT02114151, NCT02114177).

The safety profiles of all the recommended regimens above are excellent. Across numerous phase III programs, less than 1% of patients without cirrhosis discontinued treatment early and adverse events were mild. Most adverse events occurred in RBV-containing arms. Discontinuation rates were higher for patients with cirrhosis (approximately 2% for some trials) but still very low.

B. Genotype 1b

Three options with similar efficacy in general are recommended for treatment-naive patients with HCV genotype 1b infection (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks* is recommended for treatment-naive patients with HCV genotype 1b infection.

***See text for further detail on [length of treatment](#)**

Rating: Class I, Level A

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) for 12 weeks is recommended for treatment-naive patients with HCV genotype 1b infection. Rating: Class I, Level A
The addition of weight-based RBV (1000 mg [<75kg] to 1200 mg [≥75 kg]) is recommended in patients with cirrhosis.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) for 12 weeks (no cirrhosis) or

24 weeks (cirrhosis) is recommended for treatment-naive patients with HCV genotype 1b infection.

Rating: Class IIa, Level B

For HCV genotype 1b–infected, treatment-naive patients, there are 3 regimens of comparable efficacy, as outlined above.

There is no measurable difference demonstrated to date in treatment response to ledipasvir/sofosbuvir for HCV genotype 1 subtypes, thus the supporting evidence remains the same as for HCV genotype 1a–infected patients (see [Genotype 1](#)).

Paritaprevir/ritonavir/ombitasvir plus dasabuvir with or without cirrhosis and RBV (cirrhosis) was approved by the FDA for the treatment of HCV genotype 1b infection in treatment-naive patients based on 3 registration trials: SAPPHIRE-I (151 treatment-naive patients with genotype 1b HCV; those with cirrhosis were excluded), PEARL-III (419 treatment-naive patients, all with genotype 1b; those with cirrhosis were excluded), and TURQUOISE-II (119 treatment-naive and -experienced patients with genotype 1b; those with cirrhosis were included). SAPPHIRE-I reported a high SVR12 rate (98%) with 12 weeks of paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV in patients with HCV genotype 1b. ([Feld, 2014](#)) Given the high SVR12 rates seen in SAPPHIRE-I, PEARL-III was specifically designed to determine the role of weight-based RBV with paritaprevir/ritonavir/ombitasvir plus dasabuvir in treatment-naive patients with HCV genotype 1b without cirrhosis. ([Ferenci, 2014](#)) SVR12 rate was 99% in both arms, confirming the lack of added benefit of weight-based RBV for patients without cirrhosis with HCV genotype 1b. TURQUOISE-II enrolled treatment-naive and -experienced patients with CTP class A cirrhosis to receive either 12 weeks or 24 weeks of treatment with paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV. Overall, SVR12 rates were 98.5% in the 12-week arm and 100% in the 24-week arm. ([Poordad, 2014](#)) Owing to the high SVR rates, a phase III study is currently underway to determine if RBV is required as part of this regimen for HCV genotype 1b–infected patients with cirrhosis (TURQUOISE-III, NCT02219503). Until these data are available, treatment-naive patients with HCV genotype 1b infection and cirrhosis should receive RBV with paritaprevir/ritonavir/ombitasvir plus dasabuvir.

As noted, COSMOS was a phase II clinical trial of sofosbuvir plus simeprevir with or without weight-based RBV for 12 weeks or 24 weeks. ([Lawitz, 2014a](#)) The study enrolled 2 cohorts: cohort 1 included 80 patients with a prior null response to PEG-IFN and RBV with Metavir stage F0 to F2 fibrosis and cohort 2 included 87 patients who were treatment naive or who had a prior null response to PEG-IFN and RBV with Metavir stage F3 or F4 fibrosis. No virologic failure occurred in patients with HCV genotype 1b infection, suggesting high efficacy of this regimen for this genotype. Preliminary data from large prospective observational cohort studies demonstrate high SVR rates (>90%) in patients with HCV genotype 1b infection treated with sofosbuvir plus simeprevir in clinical practice. These data provide additional support for use of this DAA combination for the treatment of HCV genotype 1b. ([Jensen, 2014](#)); ([Dieterich, 2014a](#)) The FDA approval of this combination recommends 24 weeks of the combination without RBV in all patients with cirrhosis. Phase III study data of this regimen should be available in mid-2015 and are expected to provide further insight into the treatment of patients with HCV genotype 1b, including the need for RBV and optimal treatment duration for patients with cirrhosis (NCT02114151, NCT02114177).

The safety profiles to date of all recommended regimens above are excellent. Across numerous phase III programs, less than 1% of patients without cirrhosis discontinued treatment early and adverse events were mild. Most adverse events occurred in RBV-containing arms. Discontinuation rates were higher for patients with cirrhosis (approximately 2% for some trials) but still very low.

The following regimens are NOT recommended for treatment-naïve patients with HCV genotype 1.*

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥75 kg]) for 24 weeks.

Rating: Class IIb, Level A

PEG-IFN and RBV with or without sofosbuvir, simeprevir, telaprevir, or boceprevir for 12 weeks to 48 weeks.

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

*See sections on [HIV/HCV coinfection](#), [decompensated cirrhosis](#), [liver transplantation](#), and [renal impairment](#).

Although regimens of sofosbuvir and RBV or PEG-IFN and RBV plus sofosbuvir, simeprevir, telaprevir, or boceprevir for 12 weeks to 48 weeks (some using response-guided therapy) are also FDA approved, they are inferior to the current recommended regimens. Most of the IFN-containing regimens are associated with higher rates of serious adverse events (eg, anemia and rash), longer treatment duration, high pill burden, numerous drug-drug interactions, more frequent dosing, higher intensity of monitoring for continuation and stopping of therapy, and the requirement to be taken with food or with high-fat meals. Although the phase III NEUTRINO trial reported the highest SVR rate (89%) for an IFN-containing regimen (sofosbuvir [400 mg daily]) in combination with PEG-IFN 2a (180 µg by subcutaneous injection weekly) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥75 kg]) in HCV genotype 1 infection and limited exposure to IFN to just 12 weeks, the safety and tolerability profile limits its usefulness in the setting of FDA-approved, highly efficacious oral DAA combinations. ([Lawitz, 2013a](#))

PEG-IFN and RBV for 48 weeks for treatment-naïve patients with HCV genotype 1 has been superseded by treatments incorporating DAAs and should not be used.

II. Genotype 2

Recommended regimen for treatment-naive patients with HCV genotype 2 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for treatment-naive patients with HCV genotype 2 infection.

Rating: Class I, Level A

Extending treatment to 16 weeks is recommended in patients with cirrhosis.

Rating: Class IIb, Level C

Sofosbuvir (400 mg daily) was combined with weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for treatment-naive patients with HCV genotype 2 infection in 3 clinical trials, each of which enrolled patients with genotype 2 or 3 HCV: FISSION, POSITRON, and VALENCE. ([Lawitz, 2013a](#)); ([Jacobson, 2013c](#)); ([Zeuzem, 2013c](#)) The FISSION study randomized patients to receive daily PEG-IFN and RBV (800 mg) for 24 weeks or sofosbuvir plus daily weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks. ([Lawitz, 2013a](#)) The SVR rate was higher (94%) in patients who received sofosbuvir plus RBV than in those who received PEG-IFN and RBV (78%; 52/67). Across all 3 trials, 201 of the 214 (94%) patients with HCV genotype 2 achieved SVR with sofosbuvir plus RBV. Among patients who did not achieve SVR, sofosbuvir resistance-associated variants (RAVs) were not detected. ([US FDA, 2013a](#)) Based on real-world data from Trio Health, lower response rates were seen in treatment-naive patients with cirrhosis than in those without cirrhosis. ([Dieterich, 2014a](#)) Although data to support extension of therapy are not yet available for treatment-naive patients with HCV genotype 2 infection, longer treatment duration improves SVR in treatment-experienced patients with cirrhosis. ([Jacobson, 2013c](#)) Owing to the small numbers of patients with HCV genotype 2 infection and cirrhosis enrolled in the registration trials, several phase IIIb studies are ongoing to specifically determine the appropriate length of treatment for this subgroup of patients (NCT01962441, NCT 02128542). Until these data are available, extending treatment from 12 weeks to 16 weeks in HCV genotype 2-infected patients with cirrhosis is recommended.

Alternative regimens for treatment-naive patients with HCV genotype 2 infection.

None.

Although there are no alternative regimens listed, several available DAAs have activity in vitro and in vivo against HCV genotype 2. Simeprevir has moderate potency against HCV genotype 2 but has not formally been tested in combination with sofosbuvir in HCV genotype 2 infection. ([Moreno, 2012](#)) Daclatasvir (60 mg/day) with sofosbuvir (400 mg/day) for 24 weeks was associated with high rates of SVR in treatment-naive patients with HCV genotype 2 infection. Although the 50% effective concentration (EC_{50}) for daclatasvir increases several logs in the presence of the prevalent M31 polymorphism, the drug maintains

adequate activity against HCV genotype 2. ([Wang, 2014](#)) Although approved by regulatory authorities in some regions, daclatasvir is not approved by the FDA for use in the United States at this time. For patients who require treatment but cannot tolerate RBV, a discussion with a practitioner with expertise in the treatment of HCV infection is recommended prior to the use of approved RBV-free DAA combinations for HCV genotype 2 infection.

The following regimens are NOT recommended for treatment-naive patients with HCV genotype 2.

PEG-IFN and RBV for 24 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir-, boceprevir-, or ledipasvir-containing regimens

Rating: Class III, Level A

PEG-IFN 2a (180 µg weekly) or PEG-IFN 2b (1.5 µg/kg weekly) plus RBV (800 mg daily) for 24 weeks was compared with sofosbuvir (400 mg daily) plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) in the FISSION trial. ([Lawitz, 2013a](#)) The SVR12 rate achieved with PEG-IFN and RBV was lower than that achieved with sofosbuvir and RBV overall (78% and 95%, respectively) and in the subgroups of patients with or without cirrhosis. Safety and tolerability of PEG-IFN and RBV was inferior to that observed with sofosbuvir and RBV, with greater frequency of reported adverse events and laboratory abnormalities and a higher rate of treatment discontinuation owing to adverse events. Further, therapy with PEG-IFN and RBV is 12 weeks longer than that with sofosbuvir plus RBV.

Because of its poor activity in vitro and in vivo, boceprevir should not be used as therapy for patients with HCV genotype 2 infection. Although telaprevir plus PEG-IFN and RBV has antiviral activity against HCV genotype 2, ([Foster, 2011](#)) the additional adverse effects and longer duration of therapy required do not support use of this regimen. Similarly, although ledipasvir has adequate activity against HCV genotype 2, this is lost in the presence of the highly prevalent M31 polymorphism and thus is not recommended for treatment of HCV genotype 2 infection. ([Nakamoto, 2014](#))

III. Genotype 3

Recommended regimen for treatment-naive patients with HCV genotype 3 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for treatment-naive patients with HCV

genotype 3 infection.

Rating: Class I, Level B

The VALENCE study, which enrolled patients with genotype 2 or 3 HCV infection, assessed the efficacy and safety of sofosbuvir (400 mg daily) plus weight-based RBV for 24 weeks. This trial included 250 treatment-naïve (42%) and -experienced (58%) subjects with HCV genotype 3 infection. The overall SVR12 rate was 84% and was higher among treatment-naïve than -experienced patients (93% vs 77%, respectively). ([Zeuzem, 2014](#)) These results suggest that higher response rates can be achieved with a 24-week duration of sofosbuvir plus RBV than those reported for HCV genotype 3–infected participants receiving 12- or 16-week regimens in the FISSION ([Lawitz, 2013a](#)) (12 weeks, SVR12 rate: 63%), POSITRON, ([Jacobson, 2013c](#)) (12 weeks, SVR 12 rate: 61%) and FUSION (12 weeks, SVR12 rate: 30%; 16 weeks, SVR12 rate: 62%) trials. The primary reason for the higher SVR rate with extended therapy among treatment-naïve patients was a reduction in the relapse rate from 40% to 5%. In a subanalysis, response rates were similarly high among those with (n=45) and without (n=100) cirrhosis (92% and 93%, respectively).

Alternative regimens for treatment-naïve patients with HCV genotype 3 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an acceptable regimen for IFN-eligible, treatment-naïve patients with HCV genotype 3 infection.

Rating: Class IIa, Level A

The combination of sofosbuvir plus PEG-IFN and RBV was evaluated in patients with HCV genotype 3 infection. In 2 phase II clinical trials, PROTON and ELECTRON, 38 of 39 (97%) treatment-naïve patients with HCV genotype 3 infection achieved SVR with sofosbuvir plus PEG-IFN (4-12 weeks of therapy) and RBV. ([Gane, 2013b](#)) For many patients with HCV genotype 3 infection, the adverse effects and increased monitoring requirements of PEG-IFN make this less acceptable than the recommended regimen of sofosbuvir plus weight-based RBV. However, the shortened treatment period may be of interest to some.

The following regimens are NOT recommended for treatment-naïve patients with HCV genotype 3 infection.

PEG-IFN and RBV for 24 weeks to 48 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir-, boceprevir-, or simeprevir-based regimens should not be used for patients with genotype 3 HCV infection.

Rating: Class III, Level A

Although the combination of PEG-IFN and RBV is an FDA-approved regimen for HCV genotype 3 infection, its less acceptable adverse effect profile, need for more intensive monitoring, and overall lower efficacy make it less desirable than the recommended regimen.

Because of their limited in vitro and in vivo activity against HCV genotype 3, boceprevir, telaprevir, and simeprevir should not be used as therapy for patients with HCV genotype 3 infection.

Very limited phase II data are available from a single-center study (ELECTRON-II), which examined ledipasvir/sofosbuvir with (n=26) or without (n=25) RBV for 12 weeks in treatment-naive patients with HCV genotype 3 infection, 15% of whom had cirrhosis. All 26 (100%) patients in the RBV-containing arm achieved SVR12 compared with 16 of 25 (64%) of those in the RBV-free arm. Although these data raise the possibility that the addition of ledipasvir to sofosbuvir and ribavirin may shorten the course of therapy for persons with HCV genotype 3 infection, the high EC₅₀ of ledipasvir for HCV genotype 3 ([Wong, 2013](#)); ([Kohler, 2014](#)) and the homogenous patient population studied limit the generalizability of this study. Until further data are available to confirm these findings, a recommendation for use of this regimen cannot be made at this time. ([Gane, 2013b](#))

ALLY 3 is a phase III study of the investigational once-daily nonstructural protein 5A (NS5A) inhibitor daclatasvir (60 mg) plus sofosbuvir (400 mg) for 12 weeks; the study included 101 treatment-naive patients and demonstrated an SVR12 rate of 90%. In treatment-naive patients without cirrhosis (Metavir F0-F3), 97% achieved SVR12, and in treatment-naive patients with cirrhosis (Metavir F4), 58% achieved SVR12. ([Nelson, 2014](#)) Although approved by regulatory authorities in some regions, daclatasvir is not available for use in the United States at this time.

IV. Genotype 4

Three options with similar efficacy in general are recommended for treatment-naive patients with HCV genotype 4 infection (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for treatment-naive patients with HCV genotype 4 infection.

Rating: Class IIb, Level B

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir

(25 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for treatment-naive patients with HCV genotype 4 infection.

Rating: Class I, Level B

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for treatment-naive patients with HCV genotype 4 infection.

Rating: Class IIa, Level B

PEARL-I was an open-label phase IIb study that included a cohort of 86 treatment-naive patients with HCV genotype 4 infection with or without cirrhosis who received 12 weeks of the daily fixed-dose combination of paritaprevir/ritonavir/ombitasvir with or without weight-based RBV. SVR12 rates were 100% (42/42) in the group receiving RBV and 90.9% (40/44) in the group not receiving RBV. Adverse effects were generally mild, with headache, asthenia, fatigue, and nausea most commonly reported. There were no discontinuations owing to adverse events. ([Pol, 2014](#))

Several studies support the use of sofosbuvir and RBV in treatment-naive patients with HCV genotype 4 infection. In a small study of Egyptian patients in the United States treated with sofosbuvir plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]), SVR12 was achieved in 79% (11/14) and 100% (14/14) of treatment-naive patients treated for 12 weeks and 24 weeks, respectively. ([Ruane, 2014](#)) In a phase II study in Egypt, patients with HCV genotype 4 infection received daily sofosbuvir plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks or 24 weeks; among treatment-naive patients, SVR12 rates were 84% (21/25) and 92% (22/24), respectively. ([Esmat, 2014](#)) PHOTON-2, an open-label study of HIV/HCV-coinfected patients, included 31 treatment-naive patients with HCV genotype 4 infection who received daily sofosbuvir plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks. In this study, 84% of patients (26/31) achieved SVR12. ([Molina, 2014](#))

The SYNERGY trial is an open-label study evaluating 12 weeks of ledipasvir/sofosbuvir in 21 HCV genotype 4-infected patients, of whom 60% were treatment-naive and 43% had advanced fibrosis (Metavir F3 or F4). One patient took the first dose and then withdrew consent. In an interim analysis in which 20 patients had completed posttreatment week 12 follow-up, SVR12 rate was 95% by intention-to-treat analysis and 100% by per-protocol analysis. ([Kapoor, 2014](#))

Alternative regimens for treatment-naive patients with HCV genotype 4 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an acceptable regimen for treatment-naive patients with HCV genotype 4 infection.

Rating: Class II, Level B

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is an acceptable regimen for treatment-naive patients with HCV genotype 4 infection.

Rating: Class IIb, Level B

In the phase III NEUTRINO trial, ([Lawitz, 2013a](#)) 28 treatment-naive patients with HCV genotype 4 infection were treated with sofosbuvir (400 mg daily) plus PEG-IFN 2a (180 μ g weekly) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks. Of the 28 patients with HCV genotype 4 infection, 27 (96%) achieved SVR12. The single patient who did not achieve SVR had cirrhosis and had a relapse after therapy. The adverse event profile was similar to that associated with PEG-IFN and RBV therapy.

There are limited clinical data to date to support the use of the combination of 12 weeks of daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 mg]) in HCV genotype 4–infected patients, although studies are planned. Given the demonstrated activity of simeprevir in vitro and in vivo against HCV genotype 4, this combination may be considered as an alternative regimen. The open-label phase III RESTORE trial assessed the efficacy of simeprevir in combination with PEG-IFN and RBV in 107 patients with HCV genotype 4 infection, including 35 treatment-naive patients. In these treatment-naive patients, daily simeprevir (150 mg) for 12 weeks in combination with PEG-IFN and RBV for 24 weeks to 48 weeks (by response-guided therapy) produced an SVR in 83% (29 of 35). ([Moreno, 2013a](#)) These results are comparable to SVR rates observed with similar regimens in patients with HCV genotype 1 infection, suggesting that efficacy of sofosbuvir plus simeprevir for HCV genotype 4 infection may be roughly in line with the SVR rates of patients with HCV genotype 1 infection shown in the COSMOS trial. This combination has been approved in Europe for patients with HCV genotype 4 infection but is not FDA-approved for use in the United States.

The following regimens are NOT recommended for treatment-naive patients with HCV genotype 4 infection.

PEG-IFN and RBV with or without simeprevir for 24 weeks to 48 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens

Rating: Class III, Level A

PEG-IFN and RBV for 48 weeks was the previously recommended regimen for patients with HCV genotype 4 infection. ([Ghany, 2009](#)); ([AASLD/IDSA/IAS-USA, 2014](#)) Adding sofosbuvir (400 mg daily) to PEG-IFN and RBV increases response rates and markedly shortens therapy with no apparent additional adverse effects. The addition of simeprevir to PEG-IFN and RBV increases response rates but has inferior SVR rates to the other available regimens and requires a longer duration of PEG-IFN and RBV, which increases the risk of adverse events and thus is no longer recommended. ([Moreno, 2013b](#))

Because of their limited activity against genotype 4 HCV in vitro and in vivo, boceprevir and telaprevir should not be used as therapy for patients with HCV genotype 4 infection.

V. Genotype 5 and 6

Few data are available to help guide decision making for patients infected with HCV genotype 5 or 6. Nonetheless, for those patients for whom immediate treatment is required, the following recommendations have been drawn from available data.

Recommended regimen for treatment-naive patients with HCV genotype 5 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is recommended for treatment-naive patients with HCV genotype 5 infection.

Rating: Class IIa, Level B

In the phase III NEUTRINO trial, ([Lawitz, 2013a](#)) treatment-naive patients with HCV genotypes 1 (n=291), 4 (n=28), 5 (n=1), and 6 (n=6) were treated with sofosbuvir (400 mg daily) plus PEG-IFN 2a (180 μ g weekly) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks. All 6 patients with HCV genotype 6 and the 1 patient with HCV genotype 5 achieved SVR12. The adverse event profile in these patients and in the larger study population was similar to that seen with PEG-IFN and RBV therapy.

Alternative regimen for treatment-naive patients with HCV genotype 5 infection.

Weekly PEG-IFN plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 48 weeks is an alternative regimen for IFN-eligible, treatment-naive patients with HCV genotype 5 infection.

Rating: Class IIb, Level A

PEG-IFN and RBV for 48 weeks was the previously recommended regimen for patients infected with HCV

genotype 5, but the availability of recommended regimens that substantially reduce exposure to IFN and RBV make this regimen less appealing.

Recommended regimen for treatment-naive patients with HCV genotype 6 infection.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for treatment-naive patients with HCV genotype 6 infection.

Rating: Class IIa, Level B

Ledipasvir has in vitro activity against most HCV genotype 6 subtypes (exception 6e). ([Wong, 2013](#)); ([Kohler, 2014](#)) A small, 2-center, open-label study (NCT01826981) investigated the safety and in vivo efficacy of ledipasvir/sofosbuvir for 12 weeks in treatment-naive and -experienced patients with HCV genotype 6 infection. Twenty-five patients (92% treatment naive) who were primarily Asian (88%) had infection from 7 different subtypes (32%, 6a; 24%, 6e; 12%, 6l; 8%, 6m; 12%, 6p; 8%, 6q; 4%, 6r). Two patients (8%) had cirrhosis. The SVR12 rate was 96% (24/25), and the 1 patient who experienced relapse had discontinued therapy at week 8 because of drug use. No patient discontinued treatment owing to adverse events.

Alternative regimen for treatment-naive patients with HCV genotype 6 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an alternative regimen for IFN-eligible, treatment-naive patients with HCV genotype 6 infection.

Rating: Class IIa, Level B

In the phase III NEUTRINO trial, ([Lawitz, 2013a](#)) treatment-naive patients with HCV genotypes 1 (n=291), 4 (n=28), 5 (n=1), and 6 (n=6) were treated with sofosbuvir (400 mg daily) plus PEG-IFN 2a (180 μ g weekly) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks. All 6 patients with HCV genotype 6 and the 1 patient with HCV genotype 5 achieved SVR12. The adverse event profile in these patients and in the larger study population was similar to that seen with PEG-IFN and RBV therapy.

The following regimens are NOT recommended for treatment-naive patients with HCV genotype 5 or 6 infection.

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens

Rating: Class III, Level A

Because of their limited activity against genotypes 5 and 6 HCV in vitro and in vivo, boceprevir and telaprevir should not be used as therapy for patients with HCV genotype 5 or 6 infection.

Mixed Genotypes

Rarely, genotyping assays may indicate the presence of a mixed infection (eg, genotypes 1a and 2). Treatment data for mixed genotypes with direct-acting antivirals are sparse, and awaiting availability of a pangenotypic regimen may be considered. Until then, when treatment is necessary, the choice of antiviral combination and duration of treatment should maximize efficacy against each genotype represented in the assay. When the correct combination or duration is unclear, expert consultation should be sought.

Initial Treatment Table: Drug Interactions With Direct-Acting Antivirals and Selected Concomitant Medications

Concomitant Medications	Ledipasvir	Paritaprevir / Ritonavir / Ombitasvir + Dasabuvir	Simeprevir	Sofosbuvir / Velpatasvir / Glecaprevir + Sofosbuvir / Sofosbuvir + Velpatasvir / Sofosbuvir + Velpatasvir + Glecaprevir
Acid-reducing agents*	X	X		
Alfuzosin/tamsulosin		X		
Anticonvulsants	X	X	X	X
Antiretrovirals*	Coming Soon	Coming Soon	Coming Soon	tipranavir
Azole antifungals*		X	X	
Buprenorphine/naloxone		X		
Calcineurin inhibitors*		X	X	
Calcium channel blockers*		X	X	
Cisapride		X	X	
Digoxin	X		X	
Ergot derivatives		X		
Ethinyl estradiol-containing products		X		
Furosemide		X		
Gemfibrozil		X		
Glucocorticoids		X (inhaled, intranasal)	X	
Herbals				
St. John's wort		X	X	X
Milk thistle			X	
Macrolide antimicrobials*			X	
Other antiarrhythmics*		X	X	
Phosphodiesterase type 5 inhibitors*		X	X	

Pimozide		X		
Rifamycin antimicrobials*	X	X	X	X
Salmeterol		X		
Sedatives*		X	X	
Simeprevir	X			
Statins*	X	X	X	

***Some drug interactions are not class specific; see product prescribing information for specific drugs within a class.**

Complete revision made on this section on December 19, 2014. Additional changes made on January 26, 2015.

Initial Treatment Box. Summary of Recommendations for Patients Who are Initiating Therapy for HCV Infection by HCV Genotype

*/

Initial Treatment Box. Summary of Recommendations for Patients Who are Initiating Therapy for HCV Infection by HCV Genotype

Three options with similar efficacy in general are recommended for treatment-naive patients with HCV genotype 1a infection (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks* is recommended for treatment-naive patients with HCV genotype 1a infection.

*See text for further detail on [length of treatment](#)

Rating: Class I, Level A

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks (no cirrhosis) or 24 weeks (cirrhosis) is recommended for treatment-naive patients with HCV genotype 1a infection.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks (no cirrhosis) or 24 weeks (cirrhosis) is recommended for treatment-naive patients with HCV genotype 1a infection.

Rating: Class IIa, Level B

Three options with similar efficacy in general are recommended for treatment-naive patients with HCV genotype 1b infection (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks* is recommended for treatment-naive patients with HCV genotype 1b infection.

*See text for further detail on [length of treatment](#)

Rating: Class I, Level A

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) for 12 weeks is recommended for treatment-naive patients with HCV genotype 1b infection. Rating: Class I, Level A
The addition of weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) is recommended in patients with cirrhosis.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) for 12 weeks (no cirrhosis) or 24 weeks (cirrhosis) is recommended for treatment-naive patients with HCV genotype 1b infection.

Rating: Class IIa, Level B

*The following regimens are NOT recommended for treatment-naive patients with HCV genotype 1.**

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks.

Rating: Class IIb, Level A

PEG-IFN and RBV with or without sofosbuvir, simeprevir, telaprevir, or boceprevir for 12 weeks to 48 weeks.

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

*See sections on [HIV/HCV coinfection](#), [decompensated cirrhosis](#), [liver transplantation](#), and [renal impairment](#).

Recommended regimen for treatment-naive patients with HCV genotype 2 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for treatment-naive patients with HCV genotype 2 infection.

Rating: Class I, Level A

Extending treatment to 16 weeks is recommended in patients with cirrhosis.

Rating: Class IIb, Level C

Alternative regimens for treatment-naive patients with HCV genotype 2 infection.

None.

The following regimens are NOT recommended for treatment-naive patients with HCV genotype 2.

PEG-IFN and RBV for 24 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir-, boceprevir-, or ledipasvir-containing regimens

Rating: Class III, Level A

Recommended regimen for treatment-naive patients with HCV genotype 3 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for treatment-naive patients with HCV genotype 3 infection.

Rating: Class I, Level B

Alternative regimens for treatment-naive patients with HCV genotype 3 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an acceptable regimen for IFN-

eligible, treatment-naive patients with HCV genotype 3 infection.

Rating: Class IIa, Level A

The following regimens are NOT recommended for treatment-naive patients with HCV genotype 3 infection.

PEG-IFN and RBV for 24 weeks to 48 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir-, boceprevir-, or simeprevir-based regimens should not be used for patients with genotype 3 HCV infection.

Rating: Class III, Level A

Three options with similar efficacy in general are recommended for treatment-naive patients with HCV genotype 4 infection (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for treatment-naive patients with HCV genotype 4 infection.

Rating: Class IIb, Level B

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for treatment-naive patients with HCV genotype 4 infection.

Rating: Class I, Level B

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for treatment-naive patients with HCV genotype 4 infection.

Rating: Class IIa, Level B

Alternative regimens for treatment-naive patients with HCV genotype 4 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an acceptable regimen for treatment-naive patients with HCV genotype 4 infection.

Rating: Class II, Level B

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is an acceptable regimen for treatment-naive patients with HCV genotype 4 infection.

Rating: Class IIb, Level B

The following regimens are NOT recommended for treatment-naive patients with HCV genotype 4 infection.

PEG-IFN and RBV with or without simeprevir for 24 weeks to 48 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens

Rating: Class III, Level A

Recommended regimen for treatment-naive patients with HCV genotype 5 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is recommended for treatment-naive patients with HCV genotype 5 infection.

Rating: Class IIa, Level B

Alternative regimen for treatment-naive patients with HCV genotype 5 infection.

Weekly PEG-IFN plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 48 weeks is an alternative regimen for IFN-eligible, treatment-naive patients with

HCV genotype 5 infection.

Rating: Class IIb, Level A

Recommended regimen for treatment-naive patients with HCV genotype 6 infection.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for treatment-naive patients with HCV genotype 6 infection.

Rating: Class IIa, Level B

Alternative regimen for treatment-naive patients with HCV genotype 6 infection.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an alternative regimen for IFN-eligible, treatment-naive patients with HCV genotype 6 infection.

Rating: Class IIa, Level B

The following regimens are NOT recommended for treatment-naive patients with HCV genotype 5 or 6 infection.

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens

Rating: Class III, Level A

RETREATMENT OF PERSONS IN WHOM PRIOR THERAPY HAS FAILED

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

A summary of recommendations for retreatment is found in the [BOX](#).

This section provides guidance on the retreatment of a person with chronic HCV infection in whom prior therapy has failed.

The level of the evidence supporting the best treatment for each patient and the corresponding confidence in the recommendation varies as does the strength of the recommendation, and is rated in the same manner as the section on initial treatment of treatment-naïve patients ([Methods Table 2](#)). In addition, when treatment differs for a particular group (eg, those infected with various genotypes) specific recommendations are given. A regimen is classified as "Recommended" when it is favored for most patients or "Alternative" when it might be optimal in a particular subset of patients in that category. When a treatment is clearly inferior or should not be used, it is classified as "Not Recommended." Unless otherwise indicated, such regimens should not be administered to patients with HCV infection. **Specific considerations of persons with [HIV/HCV coinfection](#), [decompensated cirrhosis \(moderate or severe hepatic impairment; Child Turcotte Pugh \[CTP\] class B or C\)](#), [post-liver transplant HCV infection](#), and those with severe [renal impairment](#) or end-stage renal disease are addressed in other sections of the Guidance.**

When several regimens are offered at the same recommendation level, choice of regimen should be based on patient-specific data, including potential drug interactions. As always, patients receiving direct-acting antiviral (DAA) therapy require careful pretreatment assessment for comorbidities that may influence treatment response. All patients should have careful monitoring during treatment, particularly for anemia if RBV is included in the regimen. (See [Monitoring Section](#))

I. Genotype 1

Three highly potent DAA oral combination regimens are recommended for patients with HCV genotype 1 infection, although there are differences in the recommended regimens based on the viral subtype. With certain treatment regimens, patients infected with genotype 1a may have higher rates of virologic failure than those infected with genotype 1b. Genotype 1 HCV infection that cannot be subtyped should be treated as genotype 1a infection.

The introduction of DAAs into HCV treatment regimens increased the risk of drug interactions with other concomitant medications used by patients, and now with combinations of DAAs, attention to drug interactions is all the more important (see [Drug Interactions Table](#)). The product prescribing information and other resources (eg, <http://www.hep-druginteractions.org>) should be referenced regularly to ensure safety when prescribing DAA regimens. In particular, the daily fixed-dose combination of ledipasvir (90 mg) and sofosbuvir (400 mg) (hereafter ledipasvir/sofosbuvir) has a potential interaction with acid-suppressing medications, for example proton pump inhibitors, which may result in decreased absorption of ledipasvir and lower exposures. Because of over-the-counter access to acid-suppressing medications, a comprehensive assessment of all prescribed and over-the-counter medications is recommended prior to

initiating treatment. If possible, acid-suppressing medications should be held prior to and during the HCV treatment period to optimize ledipasvir exposure. For patients in whom interruption of acid suppression is not possible, dosing of acid suppressants is recommended per the prescribing information.

Three options with similar efficacy in general are recommended for patients with HCV genotype 1a infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for patients with HCV genotype 1a infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1a infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1a infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Three options with similar efficacy in general are recommended for patients with HCV genotype 1b infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for patients with HCV genotype 1b infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) for 12 weeks is recommended for patients with HCV genotype 1b infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1b infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Recommended regimens for patients with HCV genotype 1a or 1b infection who have compensated cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 24 weeks is recommended for patients with HCV genotype 1a or 1b infection who have compensated cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1a or 1b infection who have compensated cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level B

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for patients with HCV genotype 1a infection and for 12 weeks for patients with HCV genotype 1b infection who have compensated cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for patients with HCV genotype 1 infection, regardless of subtype, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Ledipasvir/sofosbuvir has been evaluated in patients with and without cirrhosis in whom prior treatment with PEG-IFN and RBV, with or without HCV protease inhibitors (telaprevir or boceprevir), failed. In the ION-2 study, patients who had not responded to prior PEG-IFN and RBV were treated with ledipasvir/sofosbuvir. This regimen was given for 12 weeks or 24 weeks, with or without RBV. In the population without cirrhosis, the overall response rate was 98% (95% confidence interval [CI], 96%-99%). Specifically, in patients without cirrhosis who did not respond to PEG-IFN and RBV, 33 of 35 (94%) achieved a sustained virologic response (SVR) after treatment with ledipasvir/sofosbuvir alone, and 38 of 38 (100%) patients achieved SVR after treatment with ledipasvir/sofosbuvir and RBV. ([Afdhal, 2014b](#)) This regimen was well tolerated in all groups, with no serious adverse events reported in the 12-week regimen with or without RBV. In the population with cirrhosis, patients treated for 24 weeks had higher SVR rates than those treated for 12 weeks, supporting the recommendation that HCV treatment-experienced patients with cirrhosis receive 24 weeks of treatment without RBV.

In SIRIUS, a double-blind placebo-controlled French study, patients with cirrhosis who did not respond to PEG-IFN and RBV plus telaprevir or boceprevir, were randomized to receive placebo for 12 weeks followed by ledipasvir/sofosbuvir plus RBV for 12 weeks or ledipasvir/sofosbuvir plus placebo for 24 weeks. The SVR rate was similar in each group: 74 of 77 (96%) in the group that received ledipasvir/sofosbuvir plus RBV for 12 weeks (3 patients with relapse) and 75 of 77 (97%) in the group that received ledipasvir/sofosbuvir for 24 weeks (2 patients with relapse). This observation was further supported by a meta-analysis of treatment-naïve and -experienced patients with cirrhosis who were treated with ledipasvir/sofosbuvir in phase II and III studies (including the SIRIUS study). In this analysis, ledipasvir/sofosbuvir for 12 weeks was inferior to ledipasvir/sofosbuvir for 24 weeks and ledipasvir/sofosbuvir plus RBV for 12 weeks; no difference in SVR was detected between the latter 2 groups. Safety and tolerability were similar in each group, and with the exception of anemia, reported adverse events did not differ substantially between patients treated with or without RBV. ([Bourliere, 2014a](#)); ([Bourliere, 2014b](#))

The daily fixed-dose combination of paritaprevir (150 mg), ritonavir (100 mg), and ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) (hereafter paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV) has been investigated for treatment of patients with HCV genotype 1 in whom previous PEG-IFN and RBV therapy failed. ([Zeuzem, 2014](#)) In this phase III trial, patients who did not have cirrhosis who were treated for a total of 12 weeks had a high overall rate of response with 286 of 297 (96.3%) patients achieving SVR at 12 weeks (SVR₁₂). Response rates did not differ substantially when stratified by genotype subtype (genotype 1a, 96.0% [166/173]; genotype 1b, 96.7% [119/123]) or kinetics of prior response to PEG-IFN and RBV (relapse, 95.3% [82/86]; partial response, 100% [139/146]; null response, 95.2% [139/146]). In the PEARL-II study, 179 patients without cirrhosis and HCV genotype 1b infection, in whom previous therapy with PEG-IFN and RBV failed, were treated with paritaprevir/ritonavir/ombitasvir plus dasabuvir with or without weight-based RBV for 12

weeks. ([Andreone, 2014](#)) SVR rates were high in both arms: 100% (91/91) in the RBV-free arm and 96.6% (85/88) in the RBV-containing arm, supporting the recommendation that this regimen may be used without RBV for patients with HCV genotype 1b.

In the TURQUOISE-II study, patients with CTP class A cirrhosis were treated with paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV for 12 weeks or 24 weeks. ([Poordad, 2014](#)) Of the 380 patients enrolled in this study, prior PEG-IFN and RBV therapy had failed in 220. Among the treatment-experienced patients, SVR12 was achieved in 90.2% (110/122) of patients in the 12-week arm and 96.9% (95/98) of patients in the 24-week arm. In multivariate logistic regression analysis, both prior null response to PEG-IFN and RBV therapy and genotype 1a subtype were associated with lower likelihood of SVR in patients who received 12 weeks of therapy, therefore patients with HCV genotype 1a infection should be treated for 24 weeks. Hemoglobin decline to less than 10 g/dL occurred in 7.2% of the 12-week arm and 11.0% of the 24-week arm; however, treatment discontinuation for adverse events was rare overall (2.1%).

COSMOS is a phase IIa randomized trial in which participants received sofosbuvir (400 mg once daily) plus simeprevir (150 mg once daily) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks or 24 weeks. ([Lawitz, 2013a](#)); ([Jacobson, 2013b](#)) Of the 80 patients with a null response to PEG-IFN and RBV treatment who had fibrosis (Metavir stage $\geq F2$) included in this trial, 79% to 96% achieved SVR (79%-96% in the RBV-containing arms and 93% in both RBV-free arms). Although benefit from RBV is not apparent from these results, this study was not powered to show noninferiority of RBV-free regimens. Preliminary data from large prospective observational cohort studies confirm high SVR rates ($>80\%$) in patients treated with sofosbuvir plus simeprevir in clinical practice. ([Jensen, 2014](#)); ([Dieterich, 2014a](#)) These studies also demonstrate higher rates of virologic failure in patients with HCV genotype 1a infection than patients with HCV genotype 1b infection, but additional data have not been provided regarding the contribution of treatment duration (12 weeks vs 24 weeks), use of RBV, and presence of the Q80K polymorphism to observed rates of HCV treatment failure in genotype 1a infection. In patients with cirrhosis, high SVR rates (96%) occurred in patients treated for a longer duration, supporting the recommendation that these patients be treated for 24 weeks. ([Janssen Therapeutics, 2013](#)); ([Lawitz, 2014b](#))

Recommended regimen for patients without advanced fibrosis, in whom a previous sofosbuvir-containing regimen has failed.

Based on the limited data available for effective therapy, patients without an urgent need for HCV treatment should defer antiviral therapy pending additional data or consider treatment within clinical trial settings.

Rating: Class IIb, Level C

Recommended regimen for patients who have advanced fibrosis, in whom a previous sofosbuvir-containing regimen has failed.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for patients who have cirrhosis, in whom a previous sofosbuvir-containing regimen has failed.

Rating : Class IIa, Level C

Emerging data suggest that approximately 10% to 15% of patients with HCV genotype 1 infection treated for 12 weeks with the combination of simeprevir plus sofosbuvir will experience treatment failure, typically owing to viral relapse after discontinuing therapy. Treatment failure appears to be more common in persons infected with HCV genotype 1a and those with cirrhosis. Data from the COSMOS study indicate that treatment failure following simeprevir plus sofosbuvir is associated with resistance to simeprevir with cross-resistance to other HCV nonstructural protein 3/4A (NS3/4A) protease inhibitors such as paritaprevir, telaprevir, and boceprevir. On the other hand, sofosbuvir resistance-associated variants were not observed in the COSMOS trial and are likely to be rare in clinical practice.

To date, clinical experience and trial data on the retreatment of such patients is very limited. However, retreatment after a sofosbuvir-containing treatment failure with a second course of treatment using sofosbuvir plus new agents, or retreatment with the same sofosbuvir-based regimen has been reported.

Retreatment with ledipasvir/sofosbuvir in subjects with HCV genotype 1, with or without cirrhosis, in whom a sofosbuvir-containing regimen failed has been evaluated in 2 small pilot studies utilizing ledipasvir/sofosbuvir for 12 weeks.

With prior failures of 24 weeks of sofosbuvir plus RBV, high SVR rates were noted when patients were retreated with ledipasvir/sofosbuvir for 12 weeks. ([Osinusi, 2014a](#)) Ledipasvir/sofosbuvir plus RBV has also been evaluated in subjects in whom prior treatment with sofosbuvir plus PEG-IFN and RBV or sofosbuvir and RBV failed. In this pilot study, viral response at the end of treatment is promising, however, SVR12 data are pending. ([Wyles, 2014a](#))

In the absence of data, for patients in whom prior treatment with simeprevir plus sofosbuvir failed, strong consideration should be given to enrolling them in clinical trials. For patients with minimal liver disease, consideration should be given to deferral of retreatment pending the availability of data. In patients who require retreatment more urgently, based on emerging data and the expected pattern of HCV drug resistance, patients in whom simeprevir plus sofosbuvir does not result in a cure may be treated with ledipasvir/sofosbuvir with or without RBV for 24 weeks.

Recommended regimen for patients without cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen has failed.

Daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for retreatment of patients without cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen has failed.

Rating: Class I, Level A

Two options with similar efficacy in general are recommended for patients with cirrhosis

who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and an HCV protease inhibitor regimen has failed.

Daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) for 24 weeks is recommended for retreatment of patients with cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen has failed.

Rating: Class I, Level A

Daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for retreatment of patients with cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen has failed.

Rating: Class IIa, Level B

The safety and efficacy of ledipasvir/sofosbuvir were evaluated in subjects in whom prior treatment with an HCV protease inhibitor (telaprevir or boceprevir) plus PEG-IFN and RBV failed. ([Afdhal, 2014b](#)) SVR12 rates with 12- and 24-week regimens were high during both treatment durations (94% and 98%, respectively). Relapse rates in the ION-2 retreatment trial were numerically higher in the 12-week arms than in the 24-week arms. The presence of cirrhosis or NS5A resistance-associated variants at baseline were the major reasons for the higher relapse rate in the 12-week arm. Thus, patients with cirrhosis in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen failed should receive 24 weeks of ledipasvir/sofosbuvir, and patients without cirrhosis should receive 12 weeks of ledipasvir/sofosbuvir. Based on data from the SIRIUS study, patients with cirrhosis in whom a prior protease inhibitor-containing regimen failed may also receive ledipasvir/sofosbuvir plus weight-based RBV for 12 weeks.

The following regimens are NOT recommended for patients with HCV genotype 1 infection, in whom prior treatment that included an HCV protease inhibitor has failed.

Any regimen containing PEG-IFN, including

Simeprevir, PEG-IFN, and RBV

Sofosbuvir, PEG-IFN, and RBV

Telaprevir or boceprevir, PEG-IFN, and RBV

PEG-IFN and RBV alone

Rating: Class IIb Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Any interferon-free regimen containing an HCV protease inhibitor

Simeprevir

Paritaprevir

Rating: Class IIb, Level A

Simeprevir was combined with PEG-IFN and RBV in patients in whom previous PEG-IFN and RBV dual therapy failed, in the phase IIb ASPIRE trial. ([Zeuzem, 2013a](#)); ([Janssen Therapeutics, 2013](#)); (www.fda.gov; [package insert](#)) SVR24 after 48 total weeks of therapy in the simeprevir 150 mg per day arm was 65% in patients with a previous partial response (n=23) and 53% in patients with a prior null response (n=17). Based on relatively poor response, need for prolonged therapy, and poor tolerability, this treatment is no longer recommended.

Sofosbuvir combined with PEG-IFN and RBV has high efficacy in treatment-naive patients but has not been studied prospectively in the treatment-experienced population. Based on limited prospective data and poor tolerability of PEG-IFN-based regimens, this treatment is no longer recommended.

Triple therapy with boceprevir plus PEG-IFN and RBV for 48 weeks may result in SVR for up to 52% of patients with a partial response to previous PEG-IFN and RBV treatment (RESPOND 2; [[Bacon, 2011](#)]), and up to 38% of patients with a prior null response (PROVIDE; [[Di Bisceglie, 2013](#)]). Similarly, telaprevir plus PEG-IFN and RBV resulted in an SVR24 rate of 54% to 59% among patients with a partial response to previous treatment and an SVR24 rate of 29% to 33% among those with a prior null response (REALIZE; [[Zeuzem, 2011](#)]). Because of the relatively poor efficacy, need for prolonged therapy (48 weeks), and poor tolerability, these regimens are no longer recommended.

Monotherapy with PEG-IFN, RBV, or any of the available DAAs is ineffective; further, DAA monotherapy leads to rapid selection of resistant variants.

II. Genotype 2

Recommended regimen for patients with HCV genotype 2 infection, in whom prior PEG-IFN and RBV treatment has failed.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks to 16 weeks is recommended for patients with HCV

genotype 2 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

High SVR12 rates were demonstrated in patients with HCV genotype 2 in whom prior treatment with PEG-IFN and RBV failed who were retreated with 12 weeks of sofosbuvir plus RBV. Limited data are available for treatment-experienced patients with HCV genotype 2 infection and cirrhosis; however, in the FUSION study, numerically higher SVR12 rates were seen with extension of therapy from 12 weeks (60%) to 16 weeks (78%). ([Jacobson, 2013b](#)) In contrast, the VALENCE trial found high SVR12 rates among HCV genotype 2–infected persons with cirrhosis after only 12 weeks of sofosbuvir plus RBV (88%). ([Zeuzem, 2013b](#)) Thus, definitive recommendations on the appropriate duration of sofosbuvir and RBV for treatment-experienced, HCV genotype 2–infected persons with cirrhosis cannot be made at this time. The decision to extend therapy to 16 weeks should be made on a case-by-case basis.

Alternative regimen for patients in whom previous PEG-IFN and RBV treatment failed who have HCV genotype 2 infection and are eligible to receive IFN.

Retreatment with daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an alternative for patients in whom previous PEG-IFN and RBV treatment failed who have HCV genotype 2 infection and are eligible to receive IFN.

Rating: Class IIa Level B

In recognition of the potential limitations of sofosbuvir plus RBV in harder-to-treat, HCV genotype 2–infected patients with a prior treatment failure, particularly those with cirrhosis, combination therapy with PEG-IFN has been studied. The LONESTAR-2 trial (an open-label, single-site, single-arm, phase II trial) evaluated PEG-IFN (180 μ g weekly), sofosbuvir (400 mg daily), and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg] daily in 2 divided doses for 12 weeks) in treatment-experienced patients with HCV genotype 2 or 3. ([Lawitz, 2014a](#)) Cirrhosis was present at baseline in 61% of patients. SVR12 was achieved in 22 of 23 (96%) persons with HCV genotype 2 infection. For patients with and without cirrhosis, SVR occurred in 13 of 14 (93%) and 9 of 9 (100%), respectively. Despite the limitations of this small study and accounting for the potential challenges inherent with IFN therapy, sofosbuvir plus PEG-IFN and RBV is an alternative 12-week regimen for HCV genotype 2–infected patients with cirrhosis.

The following regimens are NOT recommended for patients with HCV genotype 2 infection in whom prior HCV therapy with PEG-IFN and RBV has failed.

PEG-IFN and RBV with or without telaprevir or boceprevir

Rating: Class IIb, Level A

Fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg)

Rating: Class III, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

No HCV protease inhibitors have been approved or are indicated for the treatment of HCV genotype 2 infection. However, there is in vitro and in vivo evidence that simeprevir has activity against HCV genotype 2. Although PEG-IFN plus RBV has been the mainstay of treatment for HCV genotype 2, it requires a longer duration of therapy, is less efficacious, and has more adverse effects than the regimen recommended above.

III. Genotype 3

Recommended regimen for patients with HCV genotype 3 infection in whom prior PEG-IFN and RBV treatment has failed.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for treatment of HCV genotype 3 infection in patients in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level B

The phase III FUSION trial compared 12 weeks (n=103) with 16 weeks (n=98) of daily sofosbuvir (400 mg) and weight-based RBV in patients with HCV genotype 2 or 3 in whom previous PEG-IFN and RBV therapy failed. Of all patients, 63% had HCV genotype 3 and 34% had cirrhosis. Because persons who had experienced prior relapses to IFN-based therapy accounted for 75% of patients, the number of patients with a prior nonresponse in the study was limited. The SVR rate for patients with HCV genotype 3 in the 12-week arm was 30% (19% among patients with cirrhosis and 37% among patients without cirrhosis). Extending therapy to 16 weeks increased the SVR rate to 62%.

Based on results from the FUSION study, the phase III multicenter, randomized, placebo-controlled VALENCE trial was amended to evaluate the effects of extending sofosbuvir plus RBV therapy to 24 weeks in all patients with HCV genotype 3. As with the FUSION study, most (65%) treatment-experienced patients had a relapse. The SVR₁₂ rate after 24 weeks of therapy for treatment-experienced patients with HCV genotype 3 was 79% (60% among patients with cirrhosis and 87% in those without cirrhosis). The increased efficacy with 24 weeks of sofosbuvir plus RBV therapy across all fibrosis stages combined with a

favorable safety and tolerability profile supports the recommendation to use 24 weeks of sofosbuvir plus RBV for all HCV genotype 3–infected patients, despite the minimal number of patients studied to date. The response rate for HCV genotype 3–infected patients with cirrhosis treated for 24 weeks in the VALENCE trial (60%) was similar to that observed after 16 weeks of treatment in the FUSION trial (61%).

Alternate regimen for patients with HCV genotype 3 who are eligible to receive IFN, in whom prior PEG-IFN and RBV treatment has failed.

Retreatment with daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an alternative for patients with HCV genotype 3 infection who are eligible to receive IFN, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa Level B

Choice of specific regimen may be influenced by previous or anticipated tolerance to PEG-IFN or by the presence of advanced fibrosis or cirrhosis. For most patients, the ease of administration and tolerability of sofosbuvir plus RBV will outweigh any potential benefit associated with the addition of PEG-IFN. However, for HCV genotype 3–infected patients who have cirrhosis, responses to sofosbuvir and RBV alone for 24 weeks were suboptimal.

In the LONESTAR-2 study, adding 12 weeks of PEG-IFN to the sofosbuvir and RBV regimen resulted in numerically higher response rates among persons with HCV genotype 3 than those obtained with sofosbuvir and RBV alone for 24 weeks. Of HCV genotype 3–infected patients with and without cirrhosis, 10 of 12 (83%) achieved SVR. Given the limited number of patients in this demographic in both the VALENCE and LONESTAR-2 studies, these differences in response rates should be interpreted with caution.

The following regimens are NOT recommended for patients with HCV genotype 3 infection in whom prior PEG-IFN and RBV treatment has failed.

PEG-IFN and RBV for 24 weeks to 48 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir-, boceprevir-, or simeprevir-based regimens should not be used for patients with HCV genotype 3 infection.

Rating: Class III, Level A

No HCV protease inhibitors have been approved or are indicated for the treatment of HCV genotype 3 infection. Although PEG-IFN plus RBV has been the mainstay of treatment of HCV genotype 3 infection, it is less efficacious and has more adverse effects than the recommended regimens.

IV. Genotype 4

Data are limited to help guide decision making for patients infected with HCV genotype 4. Nonetheless, for patients in whom treatment is required, the following recommendations can be made.

Four options with similar efficacy in general are recommended for patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is an acceptable regimen for patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Daily sofosbuvir (400 mg) for 12 weeks and daily weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is recommended for retreatment of IFN-eligible patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for retreatment of patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

PEARL-I was an open-label phase IIb study that included a cohort of 49 treatment-experienced patients without cirrhosis with HCV genotype 4 infection who received 12 weeks of paritaprevir/ritonavir/ombitasvir with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]). In intention-to-treat analysis, SVR12 was achieved in 100% (41/41) of patients. This regimen was well tolerated with no serious adverse events reported. ([Pol, 2014](#))

Sofosbuvir-based regimens have also shown efficacy in patients infected with HCV genotype 4. Sofosbuvir administered with PEG-IFN and RBV for 12 weeks was investigated in the phase III NEUTRINO trial. ([Lawitz, 2013a](#)) Of the 28 treatment-naive patients with HCV genotype 4 infection, 27 (96%) achieved SVR12. In a pilot study of treatment-experienced patients of Egyptian ancestry with HCV genotype 4 infection, patients were randomized to receive sofosbuvir and RBV for 12 weeks or 24 weeks. SVR12 rate was numerically higher in the 24-week arm (89% [24/27] in the 24-week arm vs 70% [19/27] in the 12-week arm), supporting the recommendation for longer treatment duration with a sofosbuvir and RBV regimen. ([Esmat, 2014](#)) In the SYNERGY trial, 20 patients with HCV genotype 4 were treated with ledipasvir/sofosbuvir for 12 weeks. Of these patients, 40% were treatment experienced and 40% had advanced fibrosis. Preliminary data support efficacy, with 95% achieving SVR12 in an intention-to-treat analysis. (Kapoor, 2014)

The following regimens are NOT recommended for patients with HCV genotype 4 infection in whom prior PEG-IFN and RBV treatment has failed.

PEG-IFN and RBV with or without telaprevir or boceprevir

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

PEG-IFN and RBV for 48 weeks was previously recommended for patients with HCV genotype 4 infection. ([AASLD/IDSA/IAS-USA, 2014](#)) Adding sofosbuvir (400 mg daily) to PEG-IFN and RBV increases response rates and markedly shortens therapy with no apparent additional adverse effects. The addition of simeprevir to PEG-IFN and RBV increases response rates but has inferior SVR rates to the other available regimens and requires a longer duration of PEG-IFN and RBV, which increases the risk of adverse events and thus is no longer recommended. ([Moreno, 2013b](#))

Because of their limited activity against HCV genotype 4 in vitro and in vivo, boceprevir and telaprevir should not be used as therapy for patients with HCV genotype 4 infection.

V. Genotype 5 and 6

Few data are available to help guide decision making for patients infected with HCV genotype 5 or 6. Nonetheless, for those patients for whom immediate treatment is required, the following recommendations have been drawn from available data.

Recommended regimen for patients with HCV genotype 5 infection in whom prior treatment has failed.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is recommended for patients with HCV genotype 5 infection in whom prior treatment has failed.

Rating: Class IIa, Level B

In the phase III NEUTRINO trial, ([Lawitz, 2013a](#)) treatment-naïve patients with HCV genotypes 1 (n=291), 4 (n=28), 5 (n=1), and 6 (n=6) were treated with sofosbuvir (400 mg daily) plus PEG-IFN 2a (180 µg weekly) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks. All 6 patients with HCV genotype 6 and the 1 patient with HCV genotype 5 achieved SVR12. The adverse event profile in these patients and in the larger study population was similar to that seen with PEG-IFN and RBV therapy.

Alternative regimen for patients with HCV genotype 5 infection in whom prior treatment has failed.

Weekly PEG-IFN plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 48 weeks is an alternative regimen for IFN-eligible patients with HCV genotype 5 infection in whom prior treatment has failed.

Rating: Class IIb, Level A

PEG-IFN and RBV for 48 weeks was the previously recommended regimen for patients infected with HCV genotype 5, but the availability of recommended regimens that substantially reduce exposure to IFN and RBV make this regimen less appealing.

Recommended regimen for patients with HCV genotype 6 infection in whom prior treatment has failed.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for patients with HCV genotype 6 infection in whom prior treatment has failed.

Rating: Class IIa, Level B

Ledipasvir has in vitro activity against most HCV genotype 6 subtypes (exception 6e). ([Wong, 2013](#)); ([Kohler, 2014](#)) A small, 2-center, open-label study (NCT01826981) investigated the safety and in vivo efficacy of ledipasvir/sofosbuvir for 12 weeks in treatment-naive and -experienced patients with HCV genotype 6 infection. Twenty-five patients (92% treatment naive) who were primarily Asian (88%) had infection from 7 different subtypes (32%, 6a; 24%, 6e; 12%, 6l; 8%, 6m; 12%, 6p; 8%, 6q; 4%, 6r). Two patients (8%) had cirrhosis. The SVR12 rate was 96% (24/25), and the 1 patient who experienced relapse had discontinued therapy at week 8 because of drug use. No patient discontinued treatment owing to adverse events.

Alternative regimen for patients with HCV genotype 6 infection in whom prior treatment has failed.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an alternative regimen for patients with HCV genotype 6 infection in whom prior treatment has failed who are IFN-eligible.

Rating: Class IIa, Level B

In the phase III NEUTRINO trial, this regimen has been shown to have efficacy in a small number of patients with HCV genotype 6 infection (6 of 6 patients with SVR12). ([Lawitz, 2013a](#))

The following regimens are NOT recommended for patients with HCV genotype 5 or 6 infection in whom prior treatment has failed.

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens

Rating: Class III, Level A

Because of their limited activity against genotypes 5 and 6 HCV in vitro and in vivo, boceprevir and telaprevir should not be used as therapy for patients with HCV genotype 5 or 6 infection.

Mixed Genotypes

Rarely, genotyping assays may indicate the presence of a mixed infection (eg, genotypes 1a and 2). Treatment data for mixed genotypes with direct-acting antivirals are sparse, and awaiting availability of a pangenotypic regimen may be considered. Until then, when treatment is necessary, the choice of antiviral combination and duration of treatment should maximize efficacy against each genotype represented in the assay. When the correct combination or duration is unclear, expert consultation should be sought.

Complete revision made to this section on December 19, 2014. Additional changes made on January 26, 2015.

Retreatment Box. Summary of Recommendations for Patients in Whom Previous Treatment Has Failed

Retreatment Box. Recommendations for Patients in Whom Previous Treatment Has Failed

Three options with similar efficacy in general are recommended for patients with HCV genotype 1a infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for patients with HCV genotype 1a infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1a infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1a infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Three options with similar efficacy in general are recommended for patients with HCV genotype 1b infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12

weeks is recommended for patients with HCV genotype 1b infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) for 12 weeks is recommended for patients with HCV genotype 1b infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1b infection who do not have cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Recommended regimens for patients with HCV genotype 1a or 1b infection who have compensated cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 24 weeks is recommended for patients with HCV genotype 1a or 1b infection who have compensated cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1a or 1b infection who have compensated cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level B

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for patients with HCV genotype 1a infection and for 12 weeks for patients with HCV genotype 1b infection who have compensated cirrhosis, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for patients with HCV genotype 1 infection, regardless of subtype, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Recommended regimen for patients without advanced fibrosis, in whom a previous sofosbuvir-containing regimen has failed.

Based on the limited data available for effective therapy, patients without an urgent need for HCV treatment should defer antiviral therapy pending additional data or consider treatment within clinical trial settings.

Rating: Class IIb, Level C

Recommended regimen for patients who have advanced fibrosis, in whom a previous sofosbuvir-containing regimen has failed.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for patients who have cirrhosis, in whom a previous sofosbuvir-containing regimen has failed.

Rating : Class IIa, Level C

Recommended regimen for patients without cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen has failed.

Daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for retreatment of patients without cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen has failed.

Rating: Class I, Level A

Two options with similar efficacy in general are recommended for patients with cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and an HCV protease inhibitor regimen has failed.

Daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) for 24 weeks is recommended for retreatment of patients with cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen has failed.

Rating: Class I, Level A

Daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for retreatment of patients with cirrhosis who have HCV genotype 1 infection, regardless of subtype, in whom a prior PEG-IFN, RBV, and HCV protease inhibitor regimen has failed.

Rating: Class IIa, Level B

The following regimens are NOT recommended for patients with HCV genotype 1 infection, in whom prior treatment that included an HCV protease inhibitor has failed.

Any regimen containing PEG-IFN, including

Simeprevir, PEG-IFN, and RBV

Sofosbuvir, PEG-IFN, and RBV

Telaprevir or boceprevir, PEG-IFN, and RBV

PEG-IFN and RBV alone

Rating: Class IIb Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Any interferon-free regimen containing an HCV protease inhibitor

Simeprevir

Paritaprevir

Rating: Class IIb, Level A

Recommended regimen for patients with HCV genotype 2 infection, in whom prior PEG-IFN and RBV treatment has failed.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks to 16 weeks is recommended for patients with HCV genotype 2 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level A

Alternative regimen for patients in whom previous PEG-IFN and RBV treatment failed who have HCV genotype 2 infection and are eligible to receive IFN.

Retreatment with daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an alternative for patients in whom previous PEG-IFN and RBV treatment failed who have HCV genotype 2 infection and are eligible to receive IFN.

Rating: Class IIa Level B

The following regimens are NOT recommended for patients with HCV genotype 2 infection in whom prior HCV therapy with PEG-IFN and RBV has failed.

PEG-IFN and RBV with or without telaprevir or boceprevir

Rating: Class IIb, Level A

Fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg)

Rating: Class III, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Recommended regimen for patients with HCV genotype 3 infection in whom prior PEG-IFN and RBV treatment has failed.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for treatment of HCV genotype 3 infection in patients in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class I, Level B

Alternate regimen for patients with HCV genotype 3 who are eligible to receive IFN, in whom prior PEG-IFN and RBV treatment has failed.

Retreatment with daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an alternative for patients with HCV genotype 3 infection who are eligible to receive IFN, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa Level B

The following regimens are NOT recommended for patients with HCV genotype 3 infection in whom prior PEG-IFN and RBV treatment has failed.

PEG-IFN and RBV for 24 weeks to 48 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir-, boceprevir-, or simeprevir-based regimens should not be used for patients with HCV genotype 3 infection.

Rating: Class III, Level A

Four options with similar efficacy in general are recommended for patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed (listed in alphabetic order; see text).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is an acceptable regimen for patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Daily sofosbuvir (400 mg) for 12 weeks and daily weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is recommended for retreatment of IFN-eligible patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for retreatment of patients with HCV genotype 4 infection, in whom prior PEG-IFN and RBV treatment has failed.

Rating: Class IIa, Level B

The following regimens are NOT recommended for patients with HCV genotype 4 infection in whom prior PEG-IFN and RBV treatment has failed.

PEG-IFN and RBV with or without telaprevir or boceprevir

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Recommended regimen for patients with HCV genotype 5 infection in whom prior treatment has failed.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is recommended for patients with HCV genotype 5 infection in whom prior treatment has failed.

Rating: Class IIa, Level B

Alternative regimen for patients with HCV genotype 5 infection in whom prior treatment has failed.

Weekly PEG-IFN plus weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 48 weeks is an alternative regimen for IFN-eligible patients with HCV genotype 5 infection in whom prior treatment has failed.

Rating: Class IIb, Level A

Recommended regimen for patients with HCV genotype 6 infection in whom prior treatment has failed.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 12 weeks is recommended for patients with HCV genotype 6 infection in whom prior treatment has failed.

Rating: Class IIa, Level B

Alternative regimen for patients with HCV genotype 6 infection in whom prior treatment has failed.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) plus weekly PEG-IFN for 12 weeks is an alternative regimen for patients with HCV genotype 6 infection in whom prior treatment has failed who are IFN-eligible.

Rating: Class IIa, Level B

The following regimens are NOT recommended for patients with HCV genotype 5 or 6 infection in whom prior treatment has failed.

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens

Rating: Class III, Level A

MONITORING PATIENTS WHO ARE STARTING HEPATITIS C TREATMENT, ARE ON TREATMENT, OR HAVE COMPLETED THERAPY

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

A summary of recommendations for monitoring is found in the [BOX](#).

This section provides guidance on monitoring patients with chronic hepatitis C who are starting treatment, are on treatment, or have completed treatment. The section is divided into 3 parts: pretreatment and on-treatment monitoring, posttreatment follow-up for persons in whom treatment has failed to clear virus, and posttreatment follow-up for those who achieved a sustained virologic response (SVR; virologic cure).

Recommended assessments prior to starting antiviral therapy.

Assessment of potential drug-drug interactions with concomitant medications is recommended prior to starting antiviral therapy.

The following laboratory tests are recommended within 12 weeks prior to starting antiviral therapy:

Complete blood count (CBC); international normalized ratio (INR)

Hepatic function panel (albumin, total and direct bilirubin, alanine aminotransferase, aspartate aminotransferase, and alkaline phosphatase levels)

Thyroid-stimulating hormone (TSH) if IFN is used

Calculated glomerular filtration rate (GFR)

The following laboratory testing is recommended at any time prior to starting antiviral therapy:

HCV genotype and subtype

Quantitative HCV viral load, except in the circumstance that a quantitative viral load will influence duration of therapy

Rating for all statements above: Class I, Level C

Recommended monitoring during antiviral therapy.

Clinic visits or telephone contact are recommended as clinically indicated during treatment to ensure medication adherence and to monitor for adverse events and potential drug-drug interactions with newly prescribed medications.

Complete blood count (CBC), creatinine level, calculated glomerular filtration rate (GFR), and hepatic function panel are recommended after 4 weeks of treatment and as clinically indicated. Thyroid-stimulating hormone (TSH) is recommended every 12 weeks for patients receiving IFN. More frequent assessment for drug-related toxic effects (eg, CBC for patients receiving RBV) is recommended as clinically indicated.

Prompt discontinuation of therapy is recommended for any a) 10-fold increase in alanine aminotransferase (ALT) activity at week 4; or b) any increase in ALT of less than 10-fold at week 4 that is accompanied by any weakness, nausea, vomiting, or jaundice, or accompanied by increased bilirubin, alkaline phosphatase, or international normalized ratio. Asymptomatic increases in ALT of less than 10-fold elevated at week 4 should be closely monitored and repeated at week 6 and week 8.

Rating: Class I, Level B

Quantitative HCV viral load testing is recommended after 4 weeks of therapy and at 12 weeks following completion of therapy. Antiviral drug therapy should NOT be interrupted or discontinued if HCV RNA levels are not performed or available during treatment.

Quantitative HCV viral load testing can be considered at the end of treatment and 24 weeks or longer following the completion of therapy.

Rating: Class I, Level B

Recommendations for discontinuation of treatment because of lack of efficacy.

If quantitative HCV viral load is detectable at week 4 of treatment, repeat quantitative HCV RNA viral load testing is recommended after 2 additional weeks of treatment (treatment week 6). If quantitative HCV viral load has increased by greater than 10-fold ($>1 \log_{10}$ IU/mL) on repeat testing at week 6 (or thereafter), then discontinuation of HCV treatment is recommended.

The significance of a positive HCV RNA test result at week 4 that remains positive, but lower, at week 6 or week 8 is unknown. No recommendation to stop therapy or extend therapy can be provided at this time.

Rating: Class III, Level C

Recommended monitoring for pregnancy-related issues prior to and during antiviral therapy that includes RBV.

Women of childbearing age should be cautioned not to become pregnant while receiving RBV-containing antiviral regimens, and for up to 6 months after stopping.

Rating: Class I, Level C

Serum pregnancy testing is recommended for women of childbearing age prior to beginning treatment with a regimen that includes RBV.

Rating: Class I, Level C

Assessment of contraceptive use and of possible pregnancy is recommended at appropriate intervals during (and for 6 months after) RBV treatment for women of childbearing potential, and for female partners of men who receive RBV treatment.

Rating: Class I, Level C

The following regimens are NOT recommended with regard to pregnancy-related issues.

Treatment with RBV is NOT recommended for pregnant women or for women who are unwilling to adhere to use of adequate contraception, including those who are receiving RBV themselves or are sexual partners of male patients who are receiving RBV.

Rating: Class III, Level C

Female patients who have received RBV and sexual partners of male patients who have received RBV should NOT become pregnant for at least 6 months after stopping RBV.

Rating: Class III, Level C

Pretreatment and On-Treatment Monitoring

The pretreatment testing described here assumes that a decision to treat with antiviral medications has already been made and that the testing involved in deciding to treat, including testing for HCV genotype and assessment of hepatic fibrosis, has already been completed [see [Whom and When to Treat](#)].

Prior to starting treatment, patients should be evaluated for potential drug-drug interactions with selected antiviral medications (eg, <http://www.hep-druginteractions.org/>). Patients should also be educated on the proper administration of medications (eg, dose of medications, frequency of taking medicines, with or without food, missed doses, expected duration, adverse effects, etc), the crucial importance of adherence, and the necessity for close supervision and blood tests during and after treatment.

During treatment, individuals should be followed up at clinically appropriate intervals to ensure medication adherence, assess adverse events and potential drug-drug interactions, and monitor blood test results necessary for patient safety. Frequency and type of contact (eg, clinic visit, phone call, etc) are variable, but need to be sufficient to assess patient safety and response to treatment, as outlined above.

The assessment of HCV viral load at week 4 of therapy is useful to determine initial response to therapy and adherence. In phase III clinical trials, almost all patients who did not have cirrhosis had undetectable HCV RNA level at week 4; those with cirrhosis may require more than 4 weeks of treatment before HCV RNA level is undetectable. There are minimal data on how to use HCV RNA level during treatment to determine when to stop treatment for futility. The current recommendation to repeat quantitative HCV RNA testing at week 4 of treatment and to discontinue treatment if the quantitative HCV RNA level increases by more than 10-fold ($>1 \log_{10}$ IU/mL) is based on expert opinion. There are no data to support stopping treatment based on detectable HCV RNA results at weeks 2, 3, or 4 of treatment, or that detectable HCV RNA level at these time points signifies medication nonadherence. Although HCV RNA testing is recommended at week 4 of treatment, the absence of an HCV RNA level at week 4 is not a reason to discontinue treatment. Quantitative HCV RNA level testing at the end of treatment will help to differentiate viral breakthrough from relapse, if necessary. Some may choose to forego end-of-treatment viral load testing, given the high rates of viral response with the newer regimens, and to focus on the week 12 posttreatment viral load. Virologic relapse is rare at 12 or more weeks after completing treatment.

Nevertheless, repeat quantitative HCV RNA testing can be considered at 24 or more weeks after discontinuing treatment for selected patients.

The availability of IFN-free treatment regimens has simplified hepatitis C therapy by allowing shorter-duration, all-oral therapy for most patients. However, PEG-IFN and RBV-based regimens are beneficial for selected patients, and these require specific monitoring for the toxic effects (eg, anemia or neutropenia) associated with PEG-IFN or RBV use. ([RBV prescribing information, 2014a](#)); ([PEG-IFN prescribing information, 2014b](#)) In patients with a history of cardiovascular disease, RBV dose reduction to 600 mg per day is recommended for those with hemoglobin (Hgb) level below 10 g/dL and discontinuation is recommended for those with Hgb below 8.5 g/dL. In addition, although the newer all-oral regimens are generally well tolerated, adverse effects do occur. Expansion of therapy into a large population of patients may reveal toxic effects that are not apparent in registration trials. Furthermore, drug-drug interactions are possible.

RBV causes fetal death and fetal abnormalities in animals and thus it is imperative for persons of childbearing potential who receive the drug to use at least 2 reliable forms of effective contraception during treatment and for a period of 6 months thereafter. It is recommended that the health care practitioner document the discussion of potential teratogenic effects of RBV in the patient's medical record. Sofosbuvir, ledipasvir, paritaprevir, ombitasvir, and dasabuvir are pregnancy category B, although there are limited data on the use of these drugs in pregnancy. It is recommended that female patients have a thorough discussion of potential pregnancy-related drug effects prior to starting antiviral treatment. Given the relatively short duration of treatment and the potential to use RBV-free regimens in many patients, the potential risks and benefits of delaying pregnancy until HCV antiviral treatment is completed should be considered. The education of patients and caregivers about potential adverse effects and their management is an integral component of treatment and is important for a successful outcome in all patient populations.

Monitoring Patients Who Have Completed Treatment

Patients who do not achieve an SVR, because of failure of the treatment to clear or to maintain clearance of HCV infection with relapse after treatment completion, have ongoing HCV infection and the possibility of continued liver injury and transmission. Such patients should be monitored for progressive liver disease and considered for retreatment when alternative treatments are available. Patients who have undetectable HCV RNA in the serum, when assessed by a sensitive polymerase chain reaction (PCR) assay, 12 or more weeks after completing treatment, are deemed to have achieved an SVR. In these patients, HCV-related liver injury stops, although the patients remain at risk for non-HCV-related liver disease, such as fatty liver disease or alcoholic liver disease. Patients with cirrhosis remain at risk for developing hepatocellular carcinoma.

Recommended monitoring for patients in whom treatment failed to achieve a sustained virologic response.

Disease progression assessment every 6 months to 12 months with a hepatic function panel, complete blood count (CBC), and international normalized ration (INR) is recommended.

Rating: Class I, Level C

Surveillance for hepatocellular carcinoma with ultrasound testing every 6 months is recommended for patients with advanced fibrosis (ie, Metavir stage F3 or F4).

Rating: Class I, Level C

Endoscopic surveillance for esophageal varices is recommended if cirrhosis is present.

Rating: Class I, Level A

Evaluation for retreatment is recommended as effective alternative treatments become available.

Rating: Class I, Level C

The following monitoring is NOT recommended during therapy.

Routine monitoring for HCV drug resistance–associated variants during therapy is NOT recommended.

Rating: Class III, Level C

Patients in whom treatment failed to achieve an SVR remain at risk for ongoing liver injury and progression of liver fibrosis. ([Dienstag, 2011](#)) Thus, patients in whom treatment fails should be monitored for signs and symptoms of cirrhosis. There is currently no conclusive evidence to suggest that failure of antiviral treatment results in more severe liver injury or more rapidly progressive liver disease than would have occurred if the patient had not received treatment.

A small number of patients in whom an initial antiviral treatment failed have achieved SVR when treated with the same drugs for a longer duration, or when treated with alternative antiviral regimens. ([Lawitz, 2014a](#)) Thus, patients in whom treatment has failed to achieve an SVR should be considered for treatment when alternative antiviral regimens are available. Advice from a physician experienced in HCV treatment may be beneficial when considering retreatment after antiviral therapy failure.

Patients in whom antiviral therapy failed to achieve an SVR may harbor viruses that are resistant to 1 or more of the antivirals at the time of virologic “breakthrough.” ([Lawitz, 2014a](#)); ([Schneider, 2014](#)) However, there is no evidence to date that the presence of resistance-associated variants (RAVs) results in more progressive liver injury than would have occurred if the patient did not have resistant viruses. The presence

of baseline RAVs does not preclude achieving an SVR with a combination direct-acting antiviral regimen. Furthermore, RAVs are often not detectable with routine (population sequencing) detection methods, nor with more sensitive tests of HCV variants, after patients are followed up for several months. ([Schneider, 2014](#)) Subsequent retreatment with combination antivirals, particularly regimens containing antiviral drugs that have a high barrier to resistance, such as nonstructural protein 5B nucleotide polymerase inhibitors (eg, sofosbuvir), may overcome the presence of resistance to 1 or more antivirals. Certain telaprevir and boceprevir RAVs may impact simeprevir activity in vitro. However, with the exception of testing for Q80K polymorphism at baseline in patients with HCV genotype 1a infection before treatment with simeprevir plus PEG-IFN and RBV, testing for RAVs before repeat antiviral treatment is not routinely recommended. If in doubt, consultation with an expert in the treatment of HCV infection may be useful.

Recommended follow-up for patients who achieve a sustained virologic response (SVR).

For patients who do not have advanced fibrosis (ie, those with Metavir stage F0-F2), recommended follow-up is the same as if they were never infected with HCV.

Rating: Class I, Level B

Assessment for HCV recurrence or reinfection is recommended only if the patient has ongoing risk for HCV infection or otherwise unexplained hepatic dysfunction develops. In such cases, a quantitative HCV RNA assay rather than an anti-HCV serology test is recommended to test for HCV recurrence or reinfection.

Rating: Class I, Level A

Surveillance for hepatocellular carcinoma with twice-yearly ultrasound testing is recommended for patients with advanced fibrosis (ie, Metavir stage F3 or F4) who achieve an SVR.

Rating: Class I, Level C

A baseline endoscopy is recommended to screen for varices if cirrhosis is present. Patients in whom varices are found should be treated and followed up as indicated.

Rating: Class I, Level C

Assessment of other causes of liver disease is recommended for patients who develop persistently abnormal liver tests after achieving an SVR.

Rating: Class I, Level C

With the advent of highly effective HCV antiviral regimens, the likelihood of achieving an SVR among adherent, immunologically competent, treatment-naïve patients with compensated liver disease generally exceeds 90%. Of patients who achieved an SVR with PEG-IFN and RBV treatment, more than 99% have remained free of HCV infection when followed up for 5 years after completing treatment. ([Manns, 2013](#)) Thus, achieving an SVR is considered a virologic cure of HCV infection.

SVR typically aborts progression of liver injury with regression of liver fibrosis in most but not all treated patients. ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([George, 2009](#)); ([Morgan, 2013](#)); ([Singal, 2010](#)) Because of lack of progression, patients without advanced liver fibrosis (ie, Metavir stage F0-F2) who achieve an SVR should receive standard medical care that is recommended for patients who were never infected with HCV.

Among patients with advanced liver fibrosis (ie, Metavir stage F3 or F4) who achieve an SVR, decompensated liver disease (with the exception of hepatocellular carcinoma) rarely develops during follow-up, and overall survival is prolonged. ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([George, 2009](#)); ([Morgan, 2013](#)); ([Singal, 2010](#)) Patients who have advanced fibrosis or cirrhosis continue to be at risk for development of hepatocellular carcinoma after achieving an SVR, although the risk in these patients is lower than the risk in persistently viremic patients. ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([George, 2009](#)); ([Morgan, 2013](#)); ([Singal, 2010](#)) Patients with cirrhosis who achieve SVR experience increased survival (compared with patients with cirrhosis who are untreated or in whom treatment fails), but still may be at some risk for hepatocellular carcinoma; thus, they should continue to undergo regular surveillance for hepatocellular carcinoma despite the lowered risk that results after viral eradication. ([Bruix, 2011](#)) The risk of hepatocellular carcinoma among patients with advanced fibrosis prior to treatment but who have regression to minimal fibrosis after treatment is not known. In the absence of data to the contrary, such patients remain at some risk for hepatocellular carcinoma and should be monitored at regular intervals for hepatocellular carcinoma.

Liver fibrosis and liver function test results improve in most patients who achieve an SVR. ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([George, 2009](#)); ([Morgan, 2013](#)); ([Singal, 2010](#)) Bleeding from esophageal varices is rare after an SVR. ([Morisco, 2013](#)); ([Morgan, 2010](#)); ([George, 2009](#)); ([Morgan, 2013](#)); ([Singal, 2010](#)) Patients with cirrhosis should receive routine surveillance endoscopy for detection of esophageal varices if not previously done and these should be treated or followed up as indicated. ([Garcia-Tsao, 2007](#))

Patients in whom an SVR is achieved but who have another potential cause of liver disease (eg, excessive alcohol use, metabolic syndrome with or without proven fatty liver disease, or iron overload) remain at risk for progression of fibrosis. It is recommended that such patients be educated about the risk of liver disease and monitored for liver disease progression with periodic physical examinations, blood tests, and potentially, tests of liver fibrosis by a liver disease specialist.

Periodically testing patients with ongoing risk for HCV infection (eg, illicit drug use, high-risk sexual exposure) for HCV reinfection is recommended. Flares in liver enzyme test results should prompt evaluation of possible de novo reinfection with HCV through a new exposure (see [Management of Acute HCV Infection](#)). Antibody to HCV (anti-HCV) remains positive in most patients following an SVR. Thus, testing for reinfection with HCV is recommended and should be performed with an assay that detects HCV RNA (eg, a quantitative HCV RNA test).

Monitoring for HCV during chemotherapy and immunosuppression.

Prospective monitoring for HCV recurrence among patients who achieved a sustained virologic response and who are receiving immunosuppressive treatment (eg, systemic corticosteroids, antimetabolites, chemotherapy, etc) is NOT routinely recommended.

Rating: Class III, Level C

Acute liver injury is common among patients receiving chemotherapy or immunosuppressive agents; thus, testing for hepatitis viruses should be included in the laboratory assessment of the cause of liver injury. However, while individuals with inactive (no detectable virus) or past hepatitis B virus infection may experience reactivation and clinically apparent hepatitis during immunosuppressive treatment or chemotherapy, this does not occur with hepatitis C infection. Although some patients with active HCV infection, primarily those with hematologic malignancy, may have a flare in their liver enzymes during chemotherapy, this is unusual. ([Mahale, 2012](#)) Furthermore, reactivation of past HCV infection, such as after SVR or spontaneous clearance, is not anticipated since there is no residual reservoir for the virus. Thus, in this latter group, routine testing of HCV RNA during immunosuppressive treatment or prophylactic administration of antivirals during immunosuppressive treatment is not recommended.

Changes made on this section on December 19, 2014.

Monitoring Box. Summary of the Recommendations for Monitoring Patients Who Are Starting HCV Treatment, Are On Treatment, Or Have Completed Therapy

Monitoring Box. Summary of the Recommendations for Monitoring Patients Who Are Starting HCV Treatment, Are On Treatment, Or Have Completed Therapy

Recommended assessments prior to starting antiviral therapy.

Assessment of potential drug-drug interactions with concomitant medications is recommended prior to starting antiviral therapy.

The following laboratory tests are recommended within 12 weeks prior to starting antiviral therapy:

Complete blood count (CBC); international normalized ratio (INR)

Hepatic function panel (albumin, total and direct bilirubin, alanine aminotransferase, aspartate aminotransferase, and alkaline phosphatase levels)

Thyroid-stimulating hormone (TSH) if IFN is used

Calculated glomerular filtration rate (GFR)

The following laboratory testing is recommended at any time prior to starting antiviral therapy:

HCV genotype and subtype

Quantitative HCV viral load, except in the circumstance that a quantitative viral load will influence duration of therapy

Rating for all statements above: Class I, Level C

Recommended monitoring during antiviral therapy.

Clinic visits or telephone contact are recommended as clinically indicated during treatment to ensure medication adherence and to monitor for adverse events and potential drug-drug interactions with newly prescribed medications.

Complete blood count (CBC), creatinine level, calculated glomerular filtration rate (GFR), and hepatic function panel are recommended after 4 weeks of treatment and as clinically indicated. Thyroid-stimulating hormone (TSH) is recommended every 12 weeks for patients receiving IFN. More frequent assessment for drug-related toxic effects (eg, CBC for patients receiving RBV) is recommended as clinically indicated.

Prompt discontinuation of therapy is recommended for any a) 10-fold increase in alanine aminotransferase (ALT) activity at week 4; or b) any increase in ALT of less than 10-fold at week 4 that is accompanied by any weakness, nausea, vomiting, or jaundice, or accompanied by increased bilirubin, alkaline phosphatase, or international normalized ratio. Asymptomatic increases in ALT of less than 10-fold elevated at week 4 should be closely monitored and repeated at week 6 and week 8.

Rating: Class I, Level B

Quantitative HCV viral load testing is recommended after 4 weeks of therapy and at 12 weeks following completion of therapy. Antiviral drug therapy should NOT be interrupted or discontinued if HCV RNA levels are not performed or available during treatment.

Quantitative HCV viral load testing can be considered at the end of treatment and 24 weeks or longer following the completion of therapy.

Rating: Class I, Level B

Recommendations for discontinuation of treatment because of lack of efficacy.

If quantitative HCV viral load is detectable at week 4 of treatment, repeat quantitative HCV RNA viral load testing is recommended after 2 additional weeks of treatment (treatment week 6). If quantitative HCV viral load has increased by greater than 10-fold ($>1 \log_{10}$ IU/mL) on repeat testing at week 6 (or thereafter), then

discontinuation of HCV treatment is recommended.

The significance of a positive HCV RNA test result at week 4 that remains positive, but lower, at week 6 or week 8 is unknown. No recommendation to stop therapy or extend therapy can be provided at this time.

Rating: Class III, Level C

Recommended monitoring for pregnancy-related issues prior to and during antiviral therapy that includes RBV.

Women of childbearing age should be cautioned not to become pregnant while receiving RBV-containing antiviral regimens, and for up to 6 months after stopping.

Rating: Class I, Level C

Serum pregnancy testing is recommended for women of childbearing age prior to beginning treatment with a regimen that includes RBV.

Rating: Class I, Level C

Assessment of contraceptive use and of possible pregnancy is recommended at appropriate intervals during (and for 6 months after) RBV treatment for women of childbearing potential, and for female partners of men who receive RBV treatment.

Rating: Class I, Level C

The following regimens are NOT recommended with regard to pregnancy-related issues.

Treatment with RBV is NOT recommended for pregnant women or for women who are unwilling to adhere to use of adequate contraception, including those who are receiving RBV themselves or are sexual partners of male patients who are receiving RBV.

Rating: Class III, Level C

Female patients who have received RBV and sexual partners of male patients who have received RBV should NOT become pregnant for at least 6 months after stopping RBV.

Rating: Class III, Level C

Recommended monitoring for patients in whom treatment failed to achieve a sustained virologic response.

Disease progression assessment every 6 months to 12 months with a hepatic function panel, complete blood count (CBC), and international normalized ration (INR) is recommended.

Rating: Class I, Level C

Surveillance for hepatocellular carcinoma with ultrasound testing every 6 months is recommended for patients with advanced fibrosis (ie, Metavir stage F3 or F4).

Rating: Class I, Level C

Endoscopic surveillance for esophageal varices is recommended if cirrhosis is present.

Rating: Class I, Level A

Evaluation for retreatment is recommended as effective alternative treatments become available.

Rating: Class I, Level C

The following monitoring is NOT recommended during therapy.

Routine monitoring for HCV drug resistance–associated variants during therapy is NOT recommended.

Rating: Class III, Level C

Recommended follow-up for patients who achieve a sustained virologic response (SVR).

For patients who do not have advanced fibrosis (ie, those with Metavir stage F0-F2), recommended follow-up is the same as if they were never infected with HCV.

Rating: Class I, Level B

Assessment for HCV recurrence or reinfection is recommended only if the patient has ongoing risk for HCV infection or otherwise unexplained hepatic dysfunction develops. In such cases, a quantitative HCV RNA assay rather than an anti-HCV

serology test is recommended to test for HCV recurrence or reinfection.

Rating: Class I, Level A

Surveillance for hepatocellular carcinoma with twice-yearly ultrasound testing is recommended for patients with advanced fibrosis (ie, Metavir stage F3 or F4) who achieve an SVR.

Rating: Class I, Level C

A baseline endoscopy is recommended to screen for varices if cirrhosis is present. Patients in whom varices are found should be treated and followed up as indicated.

Rating: Class I, Level C

Assessment of other causes of liver disease is recommended for patients who develop persistently abnormal liver tests after achieving an SVR.

Rating: Class I, Level C

Monitoring for HCV during chemotherapy and immunosuppression.

Prospective monitoring for HCV recurrence among patients who achieved a sustained virologic response and who are receiving immunosuppressive treatment (eg, systemic corticosteroids, antimetabolites, chemotherapy, etc) is NOT routinely recommended.

Rating: Class III, Level C

UNIQUE PATIENT POPULATIONS: PATIENTS WITH HIV/HCV COINFECTION

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

The summary of recommendations for HIV-coinfected patients is in the [BOX](#).

This section provides guidance on the treatment of chronic HCV infection in HIV/HCV-coinfected patients. For individuals with acute HCV infection, please refer to the [Acute HCV](#) section. HIV/HCV-coinfected patients suffer from more liver-related morbidity and mortality, nonhepatic organ dysfunction, and overall mortality than HCV-monoinfected patients. ([Lo Re, 2014](#)); ([Chen, 2009](#)) Even in the potent HIV antiretroviral therapy era, HIV infection remains independently associated with advanced liver fibrosis and cirrhosis in patients with HCV coinfection. ([Thein, 2008b](#)); ([de Ledinghen, 2008](#)); ([Fierer, 2013](#)); ([Kirk, 2013](#))

Similar to HCV-monoinfected patients, HIV/HCV-coinfected patients cured with PEG-IFN and RBV have lower rates of hepatic decompensation, hepatocellular carcinoma, and liver-related mortality. ([Berenguer, 2009](#)); ([Limketkai, 2012](#)); ([Mira, 2013](#)) Uptake of HCV therapy was lower in the HIV/HCV-coinfected population, owing to historically lower response rates, patient comorbidities, patient and practitioner perceptions, and adverse events associated with IFN-based therapy. ([Mehta, 2006b](#)); ([Thomas, 2008](#)) With the availability of HCV direct-acting antivirals (DAAs), these barriers should diminish; however, treatment of HIV/HCV-coinfected patients requires continued awareness and attention to the complex drug interactions that can occur between DAAs and antiretroviral medications. A chart of known interactions with most US Food and Drug Administration (FDA)-approved agents is found [here](#).

Recommendations related to HCV medication interactions with HIV antiretroviral medications.

Antiretroviral drug switches, when needed, should be done in collaboration with the HIV practitioner. For HIV antiretroviral and HCV direct-acting antiviral combinations not addressed below, expert consultation is recommended.

Rating: Class I, Level A

Fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg):

Because ledipasvir increases tenofovir levels, concomitant use mandates consideration of creatinine clearance (CrCl) rate and should be avoided in those with CrCl below 60 mL/min. Because potentiation of this effect is expected when tenofovir is used with ritonavir-boosted HIV protease inhibitors, ledipasvir should be avoided with this combination (pending further data) unless antiretroviral regimen cannot be changed and the urgency of treatment is high.

Rating: Class IIa, Level C

For combinations expected to increase tenofovir levels, baseline and ongoing assessment for tenofovir nephrotoxicity is recommended.

Rating: Class IIa, Level C

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) (hereafter paritaprevir/ritonavir/ombitasvir plus dasabuvir):

Paritaprevir/ritonavir/ombitasvir plus dasabuvir should be used with antiretroviral drugs with which it does not have substantial interactions: raltegravir (and probably dolutegravir), enfuvirtide, tenofovir, emtricitabine, lamivudine, and atazanavir.

The dose of ritonavir used for boosting of HIV protease inhibitors may need to be adjusted (or held) when administered with paritaprevir/ritonavir/ombitasvir plus dasabuvir and then restored when HCV treatment is completed. The HIV protease inhibitor should be administered at the same time as the fixed-dose HCV combination.

Rating: Class IIa, Level C

Simeprevir:

Simeprevir should only be used with antiretroviral drugs with which it does not have clinically significant interactions: raltegravir (and probably dolutegravir), rilpivirine, maraviroc, enfuvirtide, tenofovir, emtricitabine, lamivudine, and abacavir.

Rating: Class IIa, Level B

The following are NOT recommended or should not be used.

Antiretroviral treatment interruption to allow HCV therapy is NOT recommended.

Rating: Class III, Level A

Fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) (hereafter ledipasvir/sofosbuvir) should NOT be used with cobicistat and elvitegravir, pending further data.

Rating: Class III, Level C

Sofosbuvir or ledipasvir/sofosbuvir should NOT be used with tipranavir.

Rating: Class III, Level B

Fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) (hereafter paritaprevir/ritonavir/ombitasvir plus dasabuvir) should NOT be used with efavirenz, rilpivirine, darunavir, or ritonavir-boosted lopinavir.

Rating: Class III, Level B

Paritaprevir/ritonavir/ombitasvir with or without dasabuvir should NOT be used in HIV/HCV-coinfected individuals who are not taking antiretroviral therapy.

Rating: Class III, Level B

Simeprevir should NOT be used with efavirenz, etravirine, nevirapine, cobicistat, or any HIV protease inhibitors.

Rating: Class III, Level B

RBV should NOT be used with didanosine, stavudine, or zidovudine.

Rating: Class III, Level B

Pharmacokinetics and Drug Interactions

Extensive recommendations for antiretroviral therapy use, including in persons anticipating HCV treatment, are found at jama.jamanetwork.com and aidsinfo.nih.gov.

Antiretroviral regimen switches may be performed to allow compatibility of DAAs, with the goal of maintaining HIV suppression without compromising future options. Considerations include prior treatment

history, responses to antiretroviral therapy, resistance profiles, and drug tolerance. ([Gunthard, 2014](#)); ([Panel on Antiretroviral Guidelines for Adults and Adolescents, 2014](#); DHHS Guidelines; [aidsinfo.nih.gov](#)) Treatment interruption in HIV/HCV-coinfected individuals is not recommended, as it is associated with increased cardiovascular events ([Strategies for Management of Antiretroviral Therapy \(SMART\) Study Group, 2006](#)) and increased rates of fibrosis progression and liver-related events. ([Tedaldi, 2008](#)); ([Thorpe, 2011](#)) If HCV treatment is nonurgent and antiretroviral compatibility and safety with DAAs is unclear, expert consultation should be sought or postponing HCV treatment should be considered until additional data are available.

Sofosbuvir

Sofosbuvir is not metabolized by the hepatic P450 enzyme complex and is a substrate (but not an inhibitor) of drug transporters, p-glycoprotein (P-gp), and breast cancer resistance protein (BCRP). It is not a substrate of organic anion-transporting polypeptide (OATP). Drug interaction studies with antiretroviral drugs (ie, efavirenz, tenofovir, emtricitabine, rilpivirine, ritonavir-boosted darunavir, and raltegravir) in noninfected persons identified no clinically significant interactions ([Kirby, 2013](#)). Sofosbuvir is not recommended for use with tipranavir because of the potential of this antiretroviral drug to induce P-gp (see [sofosbuvir prescribing](#) information).

Ledipasvir

Ledipasvir is minimally metabolized and is a substrate and inhibitor of drug transporters P-gp and BCRP. At high doses (exceeding clinical concentrations), it is an inhibitor of OATPs. Drug interaction studies of ledipasvir and sofosbuvir with antiretroviral drugs in noninfected persons did not identify clinically significant interactions with raltegravir, rilpivirine, abacavir, lamivudine, or emtricitabine. ([German, 2014](#)) Interactions with dolutegravir, maraviroc, and enfuvirtide are not expected based on their pharmacologic profiles. Ledipasvir area under the curve (AUC) serum levels are decreased by 34% when coadministered with efavirenz-containing regimens and increased by 96% when coadministered with ritonavir-boosted atazanavir. ([German, 2014](#)) No dose adjustments of ledipasvir are recommended to account for these interactions, as their clinical significance is unknown and ledipasvir is available only as a part of a fixed-dose combination with sofosbuvir.

Ledipasvir plus Sofosbuvir

Fixed-dose combination of ledipasvir (90 mg) and sofosbuvir (400 mg) (hereafter ledipasvir/sofosbuvir) increases tenofovir levels when coadministered with rilpivirine. The effect of ledipasvir/sofosbuvir on tenofovir levels was greater in magnitude when coadministered with efavirenz. Although not yet studied in HIV/HCV-coinfected individuals, there is concern of an even greater increase in tenofovir levels when ledipasvir/sofosbuvir is administered with tenofovir and HIV protease inhibitors. (see ledipasvir/sofosbuvir [prescribing information](#)) In the [ERADICATE study](#), ledipasvir/sofosbuvir was administered to 37 HIV/HCV-coinfected patients taking combination antiretroviral therapy, including 16 taking regimens containing tenofovir, emtricitabine, and efavirenz, and all with baseline creatinine clearance (CrCl) rate of 60 mL/min or higher. ([Osinusi, 2014a](#)) Changes in creatinine level or glomerular filtration rate (GFR) in these 37 patients were similar to patients not taking antiretroviral therapy. Further safety data from the phase III ION-4 study are described [below](#) regarding interactions between ledipasvir/sofosbuvir and raltegravir, rilpivirine, or efavirenz, each in combination with fixed-dose tenofovir and emtricitabine (hereafter tenofovir/emtricitabine).

Renal parameters should therefore be checked at baseline and regularly while on therapy when

ledipasvir/sofosbuvir is administered with tenofovir-containing regimens. Baseline parameters should include measuring creatinine level, electrolytes (including phosphorus), and urinary protein and glucose levels, according to recent guidelines for management of chronic kidney disease in those with HIV that include indications for nephrology consultation. ([Lucas, 2014](#)) Changing antiretroviral therapy or delaying HCV treatment if nonurgent may be considered for those at high risk for renal toxicity (especially those with CrCl between 30 mL/min and 60 mL/min or who have preexisting evidence of Fanconi syndrome) and particularly those taking tenofovir and a ritonavir-boosted HIV protease inhibitor, as there are currently few efficacy or safety data for these combinations. (see ledipasvir/sofosbuvir prescribing information) If the urgency of HCV treatment and the risk of switching antiretroviral regimens are both high and there is no safer alternative to ledipasvir/sofosbuvir, then frequent monitoring (every 2-4 weeks) of urine parameters is recommended for concomitant use with tenofovir and a ritonavir-boosted HIV protease inhibitor. Tenofovir should also be properly dosed and adjusted for CrCl rate at baseline and while on therapy. ([Lucas, 2014](#))

Fixed-dose Paritaprevir, Ritonavir, and Ombitasvir plus Dasabuvir

Paritaprevir is an inhibitor of the OATP1B1 bilirubin transporter and may increase indirect bilirubin, especially in those taking atazanavir. ([Eron, 2014](#)) Paritaprevir is metabolized by cytochrome P450 3A4 (CYP3A4) and has therefore been studied with a variety of antiretroviral drugs that are also metabolized by this enzyme; it requires ritonavir boosting to achieve long-lasting serum levels. As ritonavir has anti-HIV activity, HIV/HCV-coinfected patients should have achieved HIV RNA suppression prior to initiation of this regimen; those not taking antiretroviral therapy should avoid use of this fixed-dose combination due to the potential for low-dose ritonavir to select for HIV protease inhibitor resistance.

Paritaprevir has been studied as part of the entire regimen with the boosted HIV protease inhibitors atazanavir, darunavir, and lopinavir in noninfected individuals. There were minimal changes to maximum concentration (C_{max}) and AUC for these antiretroviral drugs, although trough concentrations were increased for atazanavir and lopinavir, and decreased for darunavir. Although paritaprevir C_{max} and AUC values were minimally to moderately increased, no dose adjustment has been recommended. ([Khatri, 2014a](#)) Because 100 mg of ritonavir is coformulated with paritaprevir and ombitasvir, the total dose of ritonavir must be carefully considered when using ritonavir-containing antiretroviral regimens. Coadministration with ritonavir-boosted lopinavir would result in 300 mg daily of ritonavir, a dose associated with substantial gastrointestinal adverse effects, this combination is not recommended. The daily fixed-dose combination of paritaprevir (150 mg), ritonavir (100 mg), and ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 mg]) (hereafter paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV) was not added to unboosted HIV protease inhibitors. Twenty-eight HIV/HCV-coinfected subjects already taking ritonavir-boosted atazanavir (with the ritonavir coming from the HCV regimen during the time of coadministration) were treated with a paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV regimen as part of the TURQUOISE-1 study. ([Sulkowski, 2015](#))

In noninfected volunteers, when paritaprevir/ritonavir/dasabuvir was combined with efavirenz, emtricitabine, and tenofovir, clinically significant gastrointestinal and neurologic adverse events occurred, coincident with elevations of alanine aminotransferase levels. When the entire regimen was combined with rilpivirine, exposures to rilpivirine were substantially increased. Therefore, rilpivirine and efavirenz should not be used with paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV. In contrast, noninfected volunteer studies did not reveal notable pharmacologic interactions with tenofovir/emtricitabine (when tested separately from other fixed-dose combinations) or raltegravir. ([Khatri, 2014b](#))

Simeprevir

Simeprevir is metabolized primarily by CYP3A4 and is therefore susceptible to drug interactions with inhibitors and inducers of the enzyme. Simeprevir is also an inhibitor of the OATP and P-gp transporters, leading to additional drug interaction concerns. Drug interaction studies with antiretroviral drugs in noninfected volunteers suggested no substantial interactions with tenofovir, rilpivirine, or raltegravir; however, simeprevir concentrations were substantially decreased when dosed with efavirenz and substantially increased when dosed with ritonavir-boosted darunavir. Use with efavirenz, etravirine, cobicistat, or boosted HIV protease inhibitors is not recommended. ([Kiser, 2013](#))

RBV

RBV has the potential for dangerous drug interactions with didanosine resulting in mitochondrial toxicity with hepatomegaly and steatosis, pancreatitis, and lactic acidosis; thus, the concomitant administration of these 2 drugs is contraindicated. ([Fleischer, 2004](#)) The combined use of RBV and zidovudine has been reported to increase the rates of anemia and the need for RBV dose reduction; thus, zidovudine is not recommended for use with RBV. ([Alvarez, 2006](#))

Recommended regimens for HIV/HCV-coinfected individuals.

HIV/HCV-coinfected persons should be treated and retreated the same as persons without HIV infection, after recognizing and managing interactions with antiretroviral medications (see [Initial Treatment of HCV Infection](#) and [Retreatment of Persons in Whom Prior Therapy Has Failed](#) sections).

Rating: Class I, Level B

Although fewer HIV/HCV-coinfected patients have been treated than HCV-monoinfected patients in trials of DAAs, efficacy rates thus far have been remarkably similar between the groups. ([Sulkowski, 2013a](#)); ([Sulkowski, 2013d](#)); ([Sulkowski, 2014](#)); ([Dieterich, 2014b](#)); ([Rodriguez-Torres, 2013](#)); ([Osinusi, 2015](#)); ([Sulkowski, 2015](#)); ([Dieterich, 2015](#)) Thus, results from HCV monoinfection studies largely justify the recommendations for HIV/HCV coinfection (discussed in the [Initial Treatment](#) and [Retreatment](#) sections). Discussion specific to HIV/HCV coinfection studies is included here.

The safety and efficacy of 12 weeks of ledipasvir/sofosbuvir was evaluated in the phase II ERADICATE study, which treated 50 HIV/HCV-coinfected, HCV genotype 1, treatment-naive patients without cirrhosis from an urban population in a single-center, open-label clinical trial. ([Osinusi, 2015](#)) Thirteen patients were not receiving antiretroviral therapy and 37 patients were on protocol-allowed medications (tenofovir, emtricitabine, rilpivirine, raltegravir, and efavirenz). Although the inclusion criteria for patients receiving antiretroviral therapy allowed CD4+ T cell counts of greater than 100/ μ L, the median CD4+ T cell count was 576/ μ L. Overall, 98% achieved sustained virologic response at 12 weeks (SVR12; 13/13 in treatment-naive arm and 36/37 in treatment-experienced arm). There were no deaths, discontinuations, or clinically significant serious adverse events. Renal function was monitored frequently during this trial and after administration of study drugs using a battery of tests (serum creatinine, estimated GFR, CrCl, urinary beta-2 microglobulin, proteinuria, and glycosuria). No clinically significant changes in these parameters or renal toxicity were observed. **A larger (ION-4) study reported similar outcomes with ledipasvir/sofosbuvir (**

[Naggie, 2015](#)). A total of 335 HCV treatment-naive and -experienced HIV-coinfected patients were enrolled and received ledipasvir/sofosbuvir once daily for 12 weeks. Patients received tenofovir/emtricitabine with raltegravir (44%), efavirenz (48%), or rilpivirine (9%). HCV genotypes included were 1a (75%), 1b (23%), and 4 (2%); 20% of patients had cirrhosis, 34% were black, and 55% had not responded to prior HCV treatment. Overall, the SVR12 rate was 96% (321/335); 2 patients had on-treatment virologic failure judged to be a result of nonadherence, 10 had virologic relapse after discontinuing treatment, 1 died from endocarditis associated with injection drug use, and 1 was lost to follow-up. SVR12 rate was 94% (63/67) among patients with cirrhosis and 97% (179/185) among treatment-experienced patients. No patients discontinued the study drug because of an adverse event. Although all patients had GFRs above 60 mL/min at study entry, drug interaction studies suggested that some patients would have elevated tenofovir levels. There were 4 patients in whom serum creatinine level rose to 0.4 mg/dL or higher: 2 remained on tenofovir, 1 had the tenofovir dose reduced, and the other stopped taking tenofovir. Neither study reported clinically significant changes in CD4 cell counts or HIV RNA levels in the study subjects. Thus, these data suggest that 12 weeks of ledipasvir/sofosbuvir is a safe and effective regimen for HIV/HCV-coinfected patients with HCV genotype 1 taking select antiretroviral therapy. ([Osinusi, 2015](#)); ([Naggie, 2015](#)) There are no data regarding an 8-week duration of ledipasvir/sofosbuvir in HIV/HCV-coinfected patients. Therefore, a shortened treatment course for persons who do not have HIV infection or cirrhosis and who have HCV RNA levels less than 6 million IU/mL cannot yet be recommended.

Paritaprevir/ritonavir/ombitasvir plus dasabuvir was FDA-approved for use in HCV genotypes 1a and 1b because of its efficacy and safety in [treatment-naive patients](#) and [PEG-IFN and RBV treatment-experienced](#) patients with and without cirrhosis. Available information about response rates with this regimen in HIV/HCV-coinfected patients comes from the first part of the phase II TURQUOISE-I study. In this study, treatment-naive (n=42) and -experienced (n=21) patients were randomized to receive either 12 weeks or 24 weeks of paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV. Of the 63 study subjects, 12 had cirrhosis, 56 had HCV genotype 1a, and 7 had HCV genotype 1b. Two study-permitted antiretroviral regimens were chosen based on healthy volunteer pharmacokinetic data: 35 patients entered taking tenofovir/emtricitabine with raltegravir and 28 patients entered taking tenofovir/emtricitabine with ritonavir-boosted atazanavir (with the ritonavir coming from the HCV regimen during the time of coadministration). Of the 31 patients to receive 12 weeks of paritaprevir/ritonavir/ombitasvir plus dasabuvir and RBV, 29 (93.5%) achieved SVR12, 1 relapsed, and 1 withdrew consent from study participation. Similarly, of the 32 subjects in the 24-week arm, 29 (90.6%) achieved SVR12, 1 experienced viral breakthrough, and 2 had apparent HCV reinfection. No treatment-related serious adverse events occurred and no subjects discontinued treatment because of medication intolerance. ([Sulkowski, 2015](#))

The combination of simeprevir plus sofosbuvir with or without RBV has been studied in the phase II COSMOS trial in patients with HCV mono-infection. ([Jacobson, 2013b](#)) This study is the main basis for the recommendation supporting the use of this all-oral combination for HCV genotype 1a or 1b mono-infection. Sofosbuvir plus simeprevir has been used anecdotally in patients with HIV/HCV coinfection, with a recent report of achieving SVR in 11 (92%) of 12 patients. ([Del Bello, 2014](#)) Despite the dearth of study data, this regimen may be considered for the treatment of HCV genotype 1 infection in patients with HIV infection who are receiving antiretroviral therapy that may be coadministered with [simeprevir](#) and [sofosbuvir](#).

Similarly, few data exist for the combination of sofosbuvir plus simeprevir for the retreatment of HCV infection in HIV/HCV-coinfected patients. However, preliminary results obtained for HCV-mono-infected patients, including those with prior treatment failure and advanced fibrosis, support the expectation that this regimen will be highly effective in HIV/HCV-coinfected patients receiving compatible antiretroviral therapy as described above (see [Retreatment](#) of HCV-mono-infected patients). ([Jacobson, 2013b](#))

The combined analysis of the phase III PHOTON 1 and PHOTON 2 studies was recently presented. These trials treated HIV/HCV-coinfected patients with HCV genotype 1 (treatment-naïve), 2, 3, or 4 (treatment-naïve and -experienced) with 400 mg of sofosbuvir and weight-based RBV. Treatment-naïve patients with HCV genotype 1 (n=226) or 4 (n=31) received 24 weeks of treatment. Treatment-naïve patients with HCV genotype 2 (n=45) received 12 weeks of treatment. Treatment-naïve patients with HCV genotype 3 received 12 weeks (n=42) or 24 weeks (n=57) of treatment. All treatment-experienced patients with HCV genotype 2, 3, or 4 (n=30, n=66, and n=31, respectively) were treated for 24 weeks. Patients with compensated cirrhosis (15%) were included. Antiretroviral regimens allowed included combinations of tenofovir and emtricitabine with efavirenz, raltegravir, ritonavir-boosted atazanavir, ritonavir-boosted darunavir, or rilpivirine. High SVR12 rates were observed for all HCV genotypes (81% for genotype 1, 89% for genotype 2, 84% each for genotypes 3 and 4). SVR12 rates were lower for patients with cirrhosis with HCV genotype 1a and treatment-experienced patients with HCV genotype 3 who were treated for 24 weeks (65% vs 85%, respectively, and 95% vs 79%, respectively) but not for others. Sofosbuvir and RBV were well tolerated. Of note, 6 patients had transient breakthrough of HIV viral load, although none required change of antiretroviral drugs. Sofosbuvir and RBV can be an effective therapy for HIV/HCV-coinfected patients, particularly for those with HCV genotypes 2, 3, or 4. ([Rockstroh, 2014](#)); ([Sulkowski, 2014](#)); ([Molina, 2015](#))

Because of response rates observed in the PHOTON 1 and PHOTON 2 trials that are similar to those seen in studies of HCV-monoinfected patients, sofosbuvir plus RBV for 12 weeks to 16 weeks is recommended for treatment-naïve, HIV/HCV-coinfected patients with HCV genotype 2, and sofosbuvir plus RBV for 24 weeks is recommended for HCV genotypes 3 and 4. Current recommendations for the retreatment of HIV-infected patients coinfecting with HCV genotype 2 or 3 are the same as those for HCV-monoinfected patients. For HCV genotype 1 infection, sofosbuvir plus RBV for 24 weeks is not recommended because of the availability of more effective regimens, assuming antiretroviral regimen compatibility can be achieved.

ALLY-2 is a phase III clinical trial that evaluated the 12-week regimen of sofosbuvir with daclatasvir (a pangenotypic NS5A inhibitor not yet approved in the United States but approved in the European Union) in patients with HIV/HCV coinfection and HCV genotypes 1 to 4. ([Wyles, 2015](#)) This open-label clinical trial enrolled both treatment-naïve (n=151) and -experienced (n=52) HIV/HCV-coinfected patients. Treatment-naïve patients were randomly assigned (2:1), with stratification by cirrhosis status and HCV genotype, to receive 12 weeks or 8 weeks of once-daily sofosbuvir 400 mg along with daclatasvir 60 mg (dose adjusted based on antiretroviral regimen); treatment-experienced patients received sofosbuvir and daclatasvir for 12 weeks. Genotype distribution was 83%, 9%, 6%, and 2% of patients, respectively, for genotypes 1, 2, 3, and 4 HCV infection, and 14% of all participants had cirrhosis. Antiretroviral drugs allowed were ritonavir-boosted (/r) darunavir, atazanavir/r, lopinavir/r, efavirenz, nevirapine, rilpivirine, raltegravir, and dolutegravir. The combination of daclatasvir and sofosbuvir once daily for 12 weeks achieved SVR12 in 97% of HIV/HCV-coinfected patients with HCV genotype 1, 2, 3, or 4, and was safe and well tolerated. Ninety-seven percent of treatment-naïve patients and 98% of -experienced patients achieved SVR. However, among patients who received 8 weeks of combination therapy, only 76% of patients achieved SVR. Factors associated with relapse in this patient group included high baseline HCV RNA level (>2 million IU/mL) (69%), concomitant use of a boosted darunavir-based antiretroviral regimen with 30 mg of daclatasvir (67%), and the presence of cirrhosis (60%). More data are needed in certain subgroups (eg, patients with HCV genotype 3 and cirrhosis who had lower response rates to this regimen and patients without HIV infection) ([Nelson, 2014](#)) and for patients receiving darunavir with daclatasvir 60 mg (as now recommended).

Data are lacking regarding the use of sofosbuvir among HIV/HCV-coinfected patients with HCV genotype 5 or 6. Given the evidence of the safety and efficacy of sofosbuvir-based regimens for HIV/HCV-coinfected

individuals infected with other HCV genotypes and the efficacy data from HCV-monoinfected individuals with these genotypes, the recommended regimens for treatment-naive and -experienced patients with HIV/HCV coinfection and these genotypes are the same as those for HCV-monoinfected patients.

In general, few HIV/HCV-coinfected patients with [cirrhosis](#) have been included in clinical trials of DAAs, and no data are available regarding HIV/HCV-coinfected patients with [renal insufficiency](#) or solid organ [transplantation](#). Despite a lack of data, it is highly likely that response rates are similar to those of HCV-monoinfected patients, as no study thus far in the DAA era has showed a lower efficacy for HIV/HCV-coinfected patients. Therefore, the respective guidance from these sections should be followed if treatment is otherwise warranted, with consideration of drug interactions.

Some HIV/HCV-coinfected patients must remain on antiretroviral regimens with drug interactions that absolutely preclude otherwise recommended DAAs that are currently available. For these situations, expert consultation is recommended.

No data currently exist to guide recommendations for the retreatment of HIV/HCV-coinfected patients or for the retreatment of simeprevir- or sofosbuvir-experienced individuals. When treatment is necessary, guidelines for HCV-monoinfected individuals are recommended.

The following regimens are NOT recommended for treatment-naive or -experienced HIV/HCV-coinfected patients.

PEG-IFN and RBV with or without simeprevir, telaprevir, or boceprevir for 24 weeks to 48 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Owing to its prolonged treatment course, adverse effects, and poor response rates, PEG-IFN with RBV is no longer recommended for the treatment of patients with any HCV genotype who are coinfecting with HIV. Telaprevir, boceprevir, or simeprevir combined with PEG-IFN and RBV have similar reported efficacy and safety in patients with HIV/HCV coinfection and HCV genotype 1 to that for patients with HCV monoinfection and genotype 1, but require 24 weeks to 48 weeks of HCV treatment. ([Sulkowski, 2013a](#)); ([Sulkowski, 2013d](#)); ([Dieterich, 2014b](#)) Telaprevir, boceprevir, and simeprevir are each substrates and to varying degrees inhibitors of CYP3A4 and thus have substantial drug interactions with antiretroviral drugs. ([van Heeswijk, 2011a](#)); ([van Heeswijk, 2011b](#)); ([Kakuda, 2012](#)); ([Johnson, 2013](#)); ([Kasserra, 2011](#)); ([Hulskotte, 2013](#)); ([Garraffo, 2013](#)); ([de Kanter, 2012](#)); ([Hammond, 2013](#)); ([Vourvahis, 2013](#)) Owing to the adverse effect profile, a prolonged required course of PEG-IFN and RBV, lower efficacy than other recommended regimens, and substantial drug interactions, these regimens are no longer recommended for HIV/HCV-coinfected patients.

Because of their limited activity in vitro and in vivo against HCV genotypes 2 and 3, boceprevir, telaprevir,

and simeprevir should not be used as therapy for HIV/HCV-coinfected patients with HCV genotype 2 or 3 infection. Boceprevir and telaprevir also have limited activity against HCV genotype 4 and should not be used as therapy for HIV/HCV-coinfected patients with HCV genotype 4 infection. There are currently insufficient data to support a recommendation for the use of simeprevir for HCV genotype 4 infection in HIV/HCV-coinfected patients.

Mixed Genotypes

Rarely, genotyping assays may indicate the presence of a mixed infection (eg, genotypes 1a and 2). Treatment data for mixed genotypes with DAAs are sparse, and awaiting availability of a pangenotypic regimen may be considered. Until then, when treatment is necessary, the choice of antiviral combination and duration of treatment should maximize efficacy against each genotype represented in the assay. When the correct combination or duration is unclear, expert consultation should be sought.

Complete revision made to this section on December 19, 2014. Additional changes made on March 13, 2015.

Unique Patient Populations: HIV/HCV Coinfection Box. Summary of Recommendations for HIV/HCV-Coinfected Patients Who are Being Treated for HCV, by Genotype

Unique Patient Populations: HIV/HCV Coinfection Box. Recommendations for HIV/HCV-Coinfected Patients Who are Being Treated for HCV, by Genotype

Recommendations related to HCV medication interactions with HIV antiretroviral medications.

Antiretroviral drug switches, when needed, should be done in collaboration with the HIV practitioner. For HIV antiretroviral and HCV direct-acting antiviral combinations not addressed below, expert consultation is recommended.

Rating: Class I, Level A

Fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg):

Because ledipasvir increases tenofovir levels, concomitant use mandates consideration of creatinine clearance (CrCl) rate and should be avoided in those with CrCl below 60 mL/min. Because potentiation of this effect is expected when tenofovir is used with ritonavir-boosted HIV protease inhibitors, ledipasvir should be avoided with this combination (pending further data) unless antiretroviral regimen cannot be changed and the urgency of treatment is high.

Rating: Class IIa, Level C

For combinations expected to increase tenofovir levels, baseline and ongoing assessment for tenofovir nephrotoxicity is recommended.

Rating: Class IIa, Level C

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25

mg) plus twice-daily dosed dasabuvir (250 mg) (hereafter paritaprevir/ritonavir/ombitasvir plus dasabuvir):

Paritaprevir/ritonavir/ombitasvir plus dasabuvir should be used with antiretroviral drugs with which it does not have substantial interactions: raltegravir (and probably dolutegravir), enfuvirtide, tenofovir, emtricitabine, lamivudine, and atazanavir.

The dose of ritonavir used for boosting of HIV protease inhibitors may need to be adjusted (or held) when administered with paritaprevir/ritonavir/ombitasvir plus dasabuvir and then restored when HCV treatment is completed. The HIV protease inhibitor should be administered at the same time as the fixed-dose HCV combination.

Rating: Class IIa, Level C

Simeprevir:

Simeprevir should only be used with antiretroviral drugs with which it does not have clinically significant interactions: raltegravir (and probably dolutegravir), rilpivirine, maraviroc, enfuvirtide, tenofovir, emtricitabine, lamivudine, and abacavir.

Rating: Class IIa, Level B

The following are NOT recommended or should not be used.

Antiretroviral treatment interruption to allow HCV therapy is NOT recommended.

Rating: Class III, Level A

Fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) (hereafter ledipasvir/sofosbuvir) should NOT be used with cobicistat and elvitegravir, pending further data.

Rating: Class III, Level C

Sofosbuvir or ledipasvir/sofosbuvir should NOT be used with tipranavir.

Rating: Class III, Level B

Fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) (hereafter paritaprevir/ritonavir/ombitasvir plus dasabuvir) should NOT be used with efavirenz, rilpivirine, darunavir, or ritonavir-boosted lopinavir.

Rating: Class III, Level B

Paritaprevir/ritonavir/ombitasvir with or without dasabuvir should NOT be used in HIV/HCV-coinfected individuals who are not taking antiretroviral therapy.

Rating: Class III, Level B

Simeprevir should NOT be used with efavirenz, etravirine, nevirapine, cobicistat, or any HIV protease inhibitors.

Rating: Class III, Level B

RBV should NOT be used with didanosine, stavudine, or zidovudine.

Rating: Class III, Level B

Recommended regimens for HIV/HCV-coinfected individuals.

HIV/HCV-coinfected persons should be treated and retreated the same as persons without HIV infection, after recognizing and managing interactions with antiretroviral medications (see [Initial Treatment of HCV Infection](#) and [Retreatment of Persons in Whom Prior Therapy Has Failed](#) sections).

Rating: Class I, Level B

The following regimens are NOT recommended for treatment-naïve or -experienced HIV/HCV-coinfected patients.

PEG-IFN and RBV with or without simeprevir, telaprevir, or boceprevir for 24 weeks to 48 weeks

Rating: Class IIb, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A



UNIQUE PATIENT POPULATIONS: PATIENTS WITH DECOMPENSATED CIRRHOSIS

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

The summary of recommendations for patients with decompensated cirrhosis is in the [BOX](#).

Decompensated Cirrhosis: Genotype 1 and 4

Patients with HCV genotype 1 or 4 with [decompensated cirrhosis](#) (moderate or severe hepatic impairment; [Child Turcotte Pugh \[CTP\] class B or C](#)) should be referred to a medical practitioner with expertise in that condition (ideally in a liver transplant center).

Rating: Class I, Level C

Recommended regimen for patients with [decompensated cirrhosis](#) (moderate or severe hepatic impairment; [CTP class B or C](#)) who may or may not be candidates for liver transplantation, including those with hepatocellular carcinoma. This regimen should be used only by highly experienced HCV practitioners.

Daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) and RBV (initial dose of 600 mg, increased as tolerated) for 12 weeks is recommended for patients with [decompensated cirrhosis](#).

Rating: Class IIb, Level C

For patients with [decompensated cirrhosis](#) and anemia or RBV intolerance, daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) for 24 weeks is recommended.

Rating: Class IIb, Level C

For patients with [decompensated cirrhosis](#) in whom prior sofosbuvir-based treatment has failed, daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) and RBV (initial dose of 600 mg, increased as tolerated) for 24 weeks is an alternative regimen.

Rating: Class IIb, Level C

The SOLAR-2 study was a multicenter randomized controlled trial of 108 patients with HCV genotypes 1 and 4 who had decompensated cirrhosis. Study participants who were treatment-naïve or -experienced, with Child Turcotte Pugh (CTP) class B cirrhosis (score 7 to 9) or CTP class C cirrhosis (score 10 to 12), were randomly assigned to receive daily fixed-dose combination ledipasvir (90 mg) and sofosbuvir (400 mg) (hereafter ledipasvir/sofosbuvir) and RBV (initial dose of 600 mg, increased as tolerated) for 12 weeks or 24 weeks. All participants had a hemoglobin level greater than 10 g/dL and a creatinine clearance rate greater than 40 mL/min. ([Flamm, 2014](#))

Excluding 6 patients who had received a transplant, sustained virologic response (SVR) was achieved in 87% of those given the 12-week treatment course and 89% of those given the 24-week treatment course. Posttherapy virologic relapse occurred in 8% and 4% of the 12- and 24-week groups, respectively. Total bilirubin and serum albumin levels improved substantially at week 4 posttherapy compared with baseline in both treatment groups. Baseline CTP and Model for End-Stage Liver Disease (MELD) scores improved in more than 50% of the treated patients, but some patients did have worsening hepatic function. During the course of the study, 5 (5%) patients died from various causes but none of the deaths were attributed to antiviral therapy. Grade 3 or 4 adverse events were more common in the 24-week arm (34%) than in the 12-week arm (15%). These results indicate that a 12-week course of ledipasvir/sofosbuvir and RBV (initial dose of 600 mg, increased as tolerated) is an appropriate regimen for patients with decompensated cirrhosis who are infected with HCV genotype 1 or 4. Such therapy may lead to objective improvements in hepatic function and reduce the likelihood of recurrent HCV infection after subsequent transplantation.

Most patients who started RBV at 600 mg per day did not receive higher doses. As of December 2014, there are no data from studies of ledipasvir/sofosbuvir without RBV in patients with decompensated cirrhosis. However, a pilot study of 14 patients with compensated cirrhosis and HCV genotype 1 infection in whom prior sofosbuvir-based therapy had failed demonstrated that ledipasvir/sofosbuvir for 12 weeks was associated with a 100% SVR rate. ([Osinusi, 2014b](#)) In addition, preliminary results of a study of 51 HCV genotype 1–infected patients in whom prior sofosbuvir-based therapy had failed demonstrated that a 12-week course of ledipasvir/sofosbuvir and RBV (initial dose of 600 mg, increased as tolerated) led to a 98% rate of SVR at 4 weeks (SVR4). ([Wyles, 2014b](#))

Another multicenter double-blind study from France reported on the use of daily ledipasvir/sofosbuvir for 24 weeks compared with daily ledipasvir/sofosbuvir and RBV for 12 weeks, with a 12-week placebo phase, in 154 patients with compensated cirrhosis and HCV genotype 1 infection in whom prior PEG-IFN and RBV treatment had failed (for most, treatment with PEG-IFN, RBV, and a protease inhibitor had also failed). ([Bourliere, 2014a](#)) The mean MELD score was 7 (range, 6-16), 26% had varices, and 13% had low serum albumin levels. The SVR12 rates were 96% with the 12-week regimen and 97% with the 24-week regimen. The most common adverse events were asthenia, headache, and pruritus, but the frequency of severe adverse events and the need for early drug discontinuation were low in both treatment groups. In light of these results, it is reasonable to consider daily ledipasvir/sofosbuvir and RBV for 12 weeks in patients with decompensated cirrhosis in whom prior sofosbuvir-based treatment has failed.

Decompensated Cirrhosis: Genotype 2 and 3

Patients with HCV genotype 2 or 3 [decompensated cirrhosis](#) (moderate or severe hepatic impairment; Child Turcotte Pugh [CTP] class B or C) should be referred to a medical practitioner with expertise in that condition (ideally in a liver transplant center).

Rating: Class I, Level C

Recommended regimen for patients with HCV genotype 2 or 3 who have [decompensated cirrhosis](#) (moderate or severe hepatic impairment; [CTP class B or C](#)) who may or may not be candidates for liver transplantation, including those with hepatocellular carcinoma. This regimen should be used only by highly experienced HCV practitioners.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) (with consideration of the patient's creatinine clearance rate and hemoglobin level) for up to 48 weeks is recommended for patients with HCV genotype 2 or 3 who have [decompensated cirrhosis](#). This regimen should be used only by highly experienced HCV practitioners.

Rating: Class IIb, Level B

In one study, 61 patients with HCV infection and hepatocellular carcinoma meeting Milan criteria for liver transplantation were treated with sofosbuvir plus RBV for up to 48 weeks. ([Curry, 2014](#)) At the time of treatment initiation, the median MELD score was 8 (range, 6-14), 17 patients had CTP scores of 7 or 8 (CTP class B cirrhosis), 45 (73%) patients had HCV genotype 1, 8 (13%) had HCV genotype 2, and 7 (11%) had HCV genotype 3. Forty-six patients underwent liver transplantation. At 12 weeks posttransplant, 30 of the 43 patients (70%) had undetectable HCV RNA levels, consistent with prevention of recurrent HCV infection. Ten patients experienced recurrent HCV infection, 9 of whom had undetectable HCV RNA levels for a duration less than 30 days pretransplant.

The most common adverse effects were fatigue (38%), anemia (21%), and headache (23%); adverse effects led to treatment discontinuation for 2 patients (3%), and 12 patients (20%) required a dose reduction of RBV. The only independent predictor of posttransplant SVR12 was the number of days of undetectable HCV RNA level pretransplant. In addition, 10 of the 11 (91%) subjects with HCV genotype 2 or 3 achieved SVR12, and only 19 of the 29 (65%) patients with HCV genotype 1 achieved SVR12. These data suggest that sofosbuvir and RBV can be given to liver transplant candidates with hepatocellular carcinoma and mildly decompensated cirrhosis but that more than 30 days of undetectable HCV RNA level are required to achieve SVR12 posttransplant.

In a sofosbuvir compassionate-use program for patients with severe recurrent HCV infection following liver transplantation who were predicted to have a less than 6-month survival rate, ([Forns, 2013b](#)) 78 subjects were treated: 44 patients were treated with sofosbuvir plus RBV, and 32 patients also received PEG-IFN. At treatment initiation, the median MELD score was 16 (range, 6-43), and fibrosing cholestatic hepatitis was documented in 20 patients. After week 12 of treatment, 91% of patients treated with sofosbuvir plus RBV and 75% of those treated with the addition of PEG-IFN achieved HCV RNA levels below the lower limit of quantification. Of 27 patients evaluated at 12 weeks posttreatment, 15 patients (56%) achieved SVR. Overall, 75% had improved or stable clinical liver disease, including improvement in hyperbilirubinemia and coagulopathy and decrease in MELD score. In this very sick population, 8 patients died and most deaths were caused by liver disease progression.

Finally, preliminary data from the ELECTRON-2 study of daily ledipasvir/sofosbuvir with or without weight-based RBV (n=26 and n=25, respectively) in previously untreated patients with HCV genotype 3 without

cirrhosis have been presented. ([Gane, 2014](#)) In this study, the SVR12 rate was 100% in the group that received ledipasvir/sofosbuvir with RBV compared with only 64% in the group that received ledipasvir/sofosbuvir alone. Although there are currently no data regarding the use of ledipasvir/sofosbuvir with RBV in patients with decompensated cirrhosis and HCV genotype 3, this regimen may be of value if proven safe and effective.

The following regimens are NOT recommended for patients with [decompensated cirrhosis](#) (moderate or severe hepatic impairment; Child Turcotte Pugh [class B or C](#)).

Any IFN-based therapy

Rating: Class III, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir-, boceprevir-, or simeprevir-based regimens

Pariteprevir-, ombitasvir-, or dasabuvir-based regimens

Rating: Class III, Level A

IFN should not be given to patients with decompensated cirrhosis (moderate or severe hepatic impairment; [CTP class B or C](#)) because of the potential for worsening hepatic decompensation. Neither telaprevir nor boceprevir should be used for this population because they must be coadministered with PEG-IFN and RBV. Very minimal data exist for the use of simeprevir in patients with decompensated cirrhosis. Until additional data become available, simeprevir should not be used in patients with decompensated cirrhosis.

Mixed Genotypes

Rarely, genotyping assays may indicate the presence of a mixed infection (eg, genotypes 1a and 2). Treatment data for mixed genotypes with direct-acting antivirals are sparse, and awaiting availability of a pangenotypic regimen may be considered. Until then, when treatment is necessary, the choice of antiviral combination and duration of treatment should maximize efficacy against each genotype represented in the assay. When the correct combination or duration is unclear, expert consultation should be sought.

Complete revision made to this section on December 19, 2014. Additional changes made on December 29, 2014 and January 26, 2015.

Unique Patient Populations: Cirrhosis Box. Summary of Recommendations for Patients with Decompensated Cirrhosis

Unique Patient Populations: Cirrhosis Box. Summary of Recommendations for Patients with Decompensated Cirrhosis

Patients with HCV genotype 1 or 4 with [decompensated cirrhosis](#) (moderate or severe hepatic impairment; [Child Turcotte Pugh \[CTP\] class B or C](#)) should be referred to a medical practitioner with expertise in that condition (ideally in a liver transplant center).

Rating: Class I, Level C

Recommended regimen for patients with [decompensated cirrhosis](#) (moderate or severe hepatic impairment; [CTP class B or C](#)) who may or may not be candidates for liver transplantation, including those with hepatocellular carcinoma. This regimen should be used only by highly experienced HCV practitioners.

Daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) and RBV (initial dose of 600 mg, increased as tolerated) for 12 weeks is recommended for patients with [decompensated cirrhosis](#).

Rating: Class IIb, Level C

For patients with [decompensated cirrhosis](#) and anemia or RBV intolerance, daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) for 24 weeks is recommended.

Rating: Class IIb, Level C

For patients with [decompensated cirrhosis](#) in whom prior sofosbuvir-based treatment has failed, daily fixed-dose combination ledipasvir (90 mg)/sofosbuvir (400 mg) and RBV (initial dose of 600 mg, increased as tolerated) for 24 weeks is an alternative regimen.

Rating: Class IIb, Level C

Patients with HCV genotype 2 or 3 [decompensated cirrhosis](#) (moderate or severe hepatic impairment; Child Turcotte Pugh [[CTP](#)] class B or C) should be referred to a medical practitioner with expertise in that condition (ideally in a liver transplant center).

Rating: Class I, Level C

Recommended regimen for patients with HCV genotype 2 or 3 who have [decompensated cirrhosis](#) (moderate or severe hepatic impairment; [CTP class B or C](#)) who may or may not be candidates for liver transplantation, including those with hepatocellular carcinoma. This regimen should be used only by highly experienced HCV practitioners.

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) (with consideration of the patient's creatinine clearance rate and hemoglobin level) for up to 48 weeks is recommended for patients with HCV genotype 2 or 3 who have [decompensated cirrhosis](#). This regimen should be used only by highly experienced HCV practitioners.

Rating: Class IIb, Level B

The following regimens are NOT recommended for patients with [decompensated cirrhosis](#) (moderate or severe hepatic impairment; Child Turcotte Pugh [class B or C](#)).

Any IFN-based therapy

Rating: Class III, Level A

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir-, boceprevir-, or simeprevir-based regimens

Pariteprevir-, ombitasvir-, or dasabuvir-based regimens

Rating: Class III, Level A

UNIQUE PATIENT POPULATIONS: PATIENTS WHO DEVELOP RECURRENT HCV INFECTION POST-LIVER TRANSPLANTATION

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

The summary of recommendations for patients who develop recurrent HCV infection post-liver transplantation is in the [BOX](#).

Recommended regimen for treatment-naive and -experienced patients with HCV genotype 1 or 4 infection in the allograft, including [compensated cirrhosis](#).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) with weight-based RBV (1000 mg [<75 kg] to 1200 mg [\geq 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1 or 4 infection in the allograft, including [compensated cirrhosis](#).

Rating: Class I, Level B

Alternative regimen for treatment-naive patients with HCV genotype 1 or 4 infection in the allograft with compensated liver disease who are RBV intolerant or ineligible.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 24 weeks is an alternative regimen for treatment-naive patients with HCV genotype 1 or 4 infection in the allograft with compensated liver disease who are RBV intolerant or ineligible.

Rating: Class I, Level B

Alternative regimen for patients with HCV genotype 1 in the allograft, including [compensated cirrhosis](#).

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [\geq 75 kg]) for 12 weeks for an alternative regimen for patients with HCV genotype 1 in the allograft, including [compensated cirrhosis](#).

Rating: Class I, Level B

Recommended regimen for treatment-naive and -experienced patients with HCV genotype 2 in the allograft, including [compensated cirrhosis](#).

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for patients with HCV genotype 2 in the allograft, including [compensated cirrhosis](#).

Rating: Class IIb, Level C

Recommended regimen for treatment-naive and -experienced patients with HCV genotype 3 in the allograft, including [compensated cirrhosis](#).

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for treatment-naive patients with HCV genotype 3 infection in the allograft, including [compensated cirrhosis](#).

Rating: Class I, Level B

Recommended regimen for treatment-naive and -experienced liver transplant recipients with [decompensated cirrhosis](#) (Child Turcotte Pugh class B or C) due to HCV genotype 3 infection in the allograft.

Daily sofosbuvir (400 mg) and low initial dose of RBV (600 mg, increasing as tolerated) for 24 weeks is recommended for liver transplant recipients with [decompensated cirrhosis](#) (Child Turcotte Pugh class B or C) due to HCV genotype 3 infection in the allograft.

Rating: Class I, Level B

Alternative regimen for patients with HCV genotype 1 in the allograft, including early (Metavir fibrosis stage F0-F2) recurrence.

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 mg]) for 24 weeks is an alternative regimen for patients with HCV genotype 1 in the allograft, without cirrhosis (Metavir fibrosis stage F0-F2).

Rating: Class I, Level B

The SOLAR-1 study was a large, multicenter, randomized controlled trial that included liver transplant recipients ($n=223$) across a broad spectrum of histologic and clinical severity of recurrence ($n=111$ with Metavir fibrosis stage F0-F3; $n=51$ with HCV genotype 1 or 4 and compensated Child Turcotte Pugh [CTP] class A cirrhosis; $n=61$ with decompensated CTP class B or C cirrhosis). Study participants were randomly assigned to receive fixed-dose combination ledipasvir (90 mg) and sofosbuvir (400 mg) (hereafter ledipasvir/sofosbuvir) and RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for either 12 weeks or 24 weeks. On an intention-to-treat basis, sustained virologic response (SVR) was achieved in 96% of patients with Metavir stage F0 to F3 fibrosis and in 96% of those with compensated cirrhosis, in both the 12- and 24-week arms; all patients received RBV. RBV dose was weight based for patients with Metavir fibrosis stage of F0 to F3 and CTP class A cirrhosis. For patients with CTP class B or C cirrhosis, RBV was initiated at 600 mg daily followed by dose escalation as tolerated. (Reddy, 2014) Only 2% of patients discontinued treatment owing to adverse events. Efficacy was lower in patients with CTP class B cirrhosis (85% SVR at 12 weeks [SVR12]) or CTP class C cirrhosis (60% SVR12), with no increase in SVR observed in patients who received 24 weeks of treatment. Mortality rate was 10% during the study among patients with CTP class B or C cirrhosis.

As the importance of RBV cannot be ascertained from the SOLAR study, in which all patients received RBV, the safest presumption is that RBV may contribute to the high SVR12 rates observed. In a previous study of a similar patient population to that of the SOLAR study, 40 patients with recurrent HCV infection following liver transplantation were treated for 24 weeks with sofosbuvir plus RBV, with SVR12 achieved in 70%. (Charlton, 2014) Although the basis for attenuated SVR rate observed in patients with more advanced HCV infection post-liver transplant, these results together with those of the sofosbuvir compassionate-use program (Forns, 2013a) suggest that the optimal period to initiate therapy may be the first 6 months to 12 months post-transplant to minimize the likelihood of having to treat patients with more advanced liver disease.

No data on ledipasvir/sofosbuvir are available for patients with HCV genotype 3 infection in the posttransplant setting. Very limited phase II data are available from a single-center study (ELECTRON-II) that examined ledipasvir/sofosbuvir used with ($n=26$) or without ($n=25$) RBV for 12 weeks in treatment-naive patients with HCV genotype 3 infection; 15% of patients had cirrhosis. All 26 (100%) patients in the

RBV-containing arm achieved SVR12 compared with 16 of 25 (64%) of those in the RBV-free arm. Although these data raise the possibility that the addition of ledipasvir to sofosbuvir and RBV may shorten the course of therapy for persons with HCV genotype 3 infection, the high effective concentration (EC₅₀) of ledipasvir for HCV genotype 3 ([Wong, 2013](#)); ([Kohler, 2014](#)) and the homogenous patient population studied limit the generalizability of this study. Until further data are available to confirm these findings, a recommendation for use of this regimen cannot be made at this time. ([Gane, 2013b](#))

In a multicenter study of 34 liver transplant recipients with mild recurrence (Metavir fibrosis stage of F0-F2) of HCV genotype 1 infection, fixed-dose combination paritaprevir (150 mg), ritonavir (100 mg), and ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) (hereafter paritaprevir/ritonavir/ombitasvir plus dasabuvir) and RBV (1000 mg [75 kg] to 1200 mg [\geq 75 kg]) was given for 24 weeks and achieved an SVR24 rate of 96%. ([Mantry, 2014](#)) Because of the drug-drug interactions between ritonavir and calcineurin inhibitors, prospective dose adjustments were needed for cyclosporine and tacrolimus. Interactions between ritonavir and other medications commonly taken by liver transplant recipients are also possible and will require detailed consideration when using this regimen. The efficacy and tolerability of this regimen in patients with more advanced HCV infection post-liver transplant are unknown.

Prospective studies of simeprevir with sofosbuvir in the posttransplant setting are ongoing. A retrospective multicenter analysis of sofosbuvir (400 mg daily) plus simeprevir (150 mg daily) with or without RBV in 77 recipients reported an SVR4 rate of 92%. ([Pungpapong, 2014](#)) The coadministration of single-dose cyclosporine with simeprevir resulted in a 19% increase in cyclosporine concentrations and no change in simeprevir concentrations (see [simeprevir prescribing information](#)). However, in an interim analysis of an ongoing study (TMC435HPC3016), concomitant use of simeprevir (plus daclatasvir and RBV) with cyclosporine at steady state resulted in an approximately 6-fold increase in plasma concentrations of simeprevir compared with historical data of simeprevir in the absence of cyclosporine. This interaction may be caused by inhibition of organic ion-transporting polypeptide 1B1 (OATP1B1), p-glycoprotein (P-gp), and cytochrome P450 3A (CYP3A) by cyclosporine. Although the current US Food and Drug Administration labeling does not recommend against coadministration of simeprevir and cyclosporine, given these new findings, simeprevir should not be coadministered with cyclosporine.

The coadministration of single-dose tacrolimus with simeprevir did not result in a notable change of tacrolimus concentrations (see [simeprevir prescribing information](#)). In an ongoing study, concomitant use of simeprevir with tacrolimus also resulted in a 2-fold increase in plasma concentrations of simeprevir compared with historical data (see [simeprevir prescribing information](#)). Based on phase I studies, a 2-fold increase in simeprevir concentrations is unlikely to be clinically significant.

Clinicians may consider the use of sofosbuvir plus simeprevir in patients receiving tacrolimus with therapeutic drug monitoring, particularly in those expected to have difficulty tolerating RBV (eg, patients with impaired renal function or anemia) or who are unable to forego proton pump inhibitor therapy (proton pump inhibitors attenuate ledipasvir absorption by >90%). A further option in patients who are RBV intolerant is 24 weeks of ledipasvir/sofosbuvir.

The following regimens are NOT recommended for treatment-naive patients with compensated allograft HCV infection.

Regimens containing PEG-IFN

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens should not be used for patients with compensated allograft HCV infection.

Rating: Class III, Level A

Telaprevir or boceprevir should not be used in the post–liver transplant population because of associated toxicities and drug interactions with calcineurin inhibitors.

Decompensated Cirrhosis

Recommended regimen for treatment-naive and -experienced liver transplant recipients with [decompensated cirrhosis](#) (Child Turcotte Pugh class B or C) due to HCV genotype 1 or 4 infection in the allograft.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) with a low initial dose of RBV (600 mg, increasing as tolerated) for 12 weeks is recommended for liver transplant recipients with [decompensated cirrhosis](#) (Child Turcotte Pugh class B or C) due to HCV genotype 1 or 4 infection in the allograft.

Rating: Class I, Level B

Recommended regimen for treatment-naive and -experienced liver transplant recipients with [decompensated cirrhosis](#) (Child Turcotte Pugh class B or C) due to HCV genotype 2 infection in the allograft.

Daily sofosbuvir (400 mg) and RBV (initial dose 600 mg/day, increased monthly by 200 mg/day as tolerated to weight-based dose of 1000 mg [<75 kg] to 1200 mg [≥ 75 kg] mg) for 24 weeks is recommended for liver transplant recipients with [decompensated cirrhosis](#) (Child Turcotte Pugh class B or C) due to HCV genotype 2 infection in the allograft.

Rating: Class IIb, Level C

Recommended regimen for treatment-naive and -experienced patients with decompensated cirrhosis (Child Turcotte Pugh class B or C) due to HCV genotype 3 infection in the allograft.

Daily sofosbuvir (400 mg) and low initial dose of RBV (600 mg, increasing as tolerated) for 24 weeks is recommended for persons with decompensated cirrhosis (Child Turcotte Pugh class B or C) due to HCV genotype 3 infection in the allograft.

Rating: Class I, Level B

The following regimens are NOT recommended for patients with decompensated allograft HCV infection.

Regimens containing PEG-IFN

Regimens containing simeprevir

Fixed-dose combination of paritaprevir (150 mg), ritonavir (100 mg), and ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg])

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens

Rating: Class III, Level A

Mixed Genotypes

Rarely, genotyping assays may indicate the presence of a mixed infection (eg, genotypes 1a and 2). Treatment data for mixed genotypes with direct-acting antivirals are sparse, and awaiting availability of a pangentypic regimen may be considered. Until then, when treatment is necessary, the choice of antiviral combination and duration of treatment should maximize efficacy against each genotype represented in the assay. When the correct combination or duration is unclear, expert consultation should be sought.

Complete revision made to this section on December 19, 2014. Additional changes made on January 19 and 26, 2015.

Unique Patient Populations: Post-Liver Transplantation Box. Summary of Recommendations for Patients Who Develop Recurrent HCV Infection Post-Liver Transplantation

Recommended regimen for treatment-naive and -experienced patients with HCV genotype 1 or 4 infection in the allograft, including [compensated cirrhosis](#).

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) with weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks is recommended for patients with HCV genotype 1 or 4 infection in the allograft, including [compensated cirrhosis](#).

Rating: Class I, Level B

Alternative regimen for treatment-naive patients with HCV genotype 1 or 4 infection in the allograft with compensated liver disease who are RBV intolerant or ineligible.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) for 24 weeks is an alternative regimen for treatment-naive patients with HCV genotype 1 or 4 infection in the allograft with compensated liver disease who are RBV intolerant or ineligible.

Rating: Class I, Level B

Alternative regimen for patients with HCV genotype 1 in the allograft, including [compensated cirrhosis](#).

Daily sofosbuvir (400 mg) plus simeprevir (150 mg) with or without weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 12 weeks for an alternative regimen for patients with HCV genotype 1 in the allograft, including [compensated cirrhosis](#).

Rating: Class I, Level B

Recommended regimen for treatment-naive and -experienced patients with HCV genotype 2 in the allograft, including [compensated cirrhosis](#).

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for patients with HCV genotype 2 in the

allograft, including [compensated cirrhosis](#).

Rating: Class IIb, Level C

Recommended regimen for treatment-naïve and -experienced patients with HCV genotype 3 in the allograft, including [compensated cirrhosis](#).

Daily sofosbuvir (400 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg]) for 24 weeks is recommended for treatment-naïve patients with HCV genotype 3 infection in the allograft, including [compensated cirrhosis](#).

Rating: Class I, Level B

Recommended regimen for treatment-naïve and -experienced liver transplant recipients with [decompensated cirrhosis](#) (Child Turcotte Pugh class B or C) due to HCV genotype 3 infection in the allograft.

Daily sofosbuvir (400 mg) and low initial dose of RBV (600 mg, increasing as tolerated) for 24 weeks is recommended for liver transplant recipients with [decompensated cirrhosis](#) (Child Turcotte Pugh class B or C) due to HCV genotype 3 infection in the allograft.

Rating: Class I, Level B

Alternative regimen for patients with HCV genotype 1 in the allograft, including early (Metavir fibrosis stage F0-F2) recurrence.

Daily fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and weight-based RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 mg]) for 24 weeks is an alternative regimen for patients with HCV genotype 1 in the allograft, without cirrhosis (Metavir fibrosis stage F0-F2).

Rating: Class I, Level B

The following regimens are NOT recommended for treatment-naïve patients with compensated allograft HCV infection.

Regimens containing PEG-IFN

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens should not be used for patients with compensated allograft HCV infection.

Rating: Class III, Level A

Recommended regimen for treatment-naive and -experienced liver transplant recipients with decompensated cirrhosis (Child Turcotte Pugh class B or C) due to HCV genotype 1 or 4 infection in the allograft.

Daily fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg) with a low initial dose of RBV (600 mg, increasing as tolerated) for 12 weeks is recommended for liver transplant recipients with decompensated cirrhosis (Child Turcotte Pugh class B or C) due to HCV genotype 1 or 4 infection in the allograft.

Rating: Class I, Level B

Recommended regimen for treatment-naive and -experienced liver transplant recipients with decompensated cirrhosis (Child Turcotte Pugh class B or C) due to HCV genotype 2 infection in the allograft.

Daily sofosbuvir (400 mg) and RBV (initial dose 600 mg/day, increased monthly by 200 mg/day as tolerated to weight-based dose of 1000 mg [<75 kg] to 1200 mg [\geq 75 kg] mg) for 24 weeks is recommended for liver transplant recipients with decompensated cirrhosis (Child Turcotte Pugh class B or C) due to HCV genotype 2 infection in the allograft.

Rating: Class IIb, Level C

Recommended regimen for treatment-naive and -experienced patients with decompensated cirrhosis (Child Turcotte Pugh class B or C) due to HCV genotype 3 infection in the allograft.

Daily sofosbuvir (400 mg) and low initial dose of RBV (600 mg, increasing as tolerated) for 24 weeks is recommended for persons with decompensated cirrhosis (Child Turcotte Pugh class B or C) due to HCV genotype 3 infection in the allograft.

Rating: Class I, Level B

The following regimens are NOT recommended for patients with decompensated allograft HCV infection.

Regimens containing PEG-IFN

Regimens containing simeprevir

Fixed-dose combination of paritaprevir (150 mg), ritonavir (100 mg), and ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) and RBV (1000 mg [<75 kg] to 1200 mg [≥ 75 kg])

Monotherapy with PEG-IFN, RBV, or a direct-acting antiviral

Rating: Class III, Level A

Telaprevir- or boceprevir-based regimens

Rating: Class III, Level A

Minor changes were made on January 19, 2015.

UNIQUE PATIENT POPULATIONS: PATIENTS WITH RENAL IMPAIRMENT

/*-->*/

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

The summary of recommendations for patients with renal impairment, including severe renal impairment (creatinine clearance <30 mL/min) or end-stage renal disease requiring hemodialysis or peritoneal dialysis is found in the [BOX](#).

Recommended dosage adjustments for patients with renal impairment, including severe renal impairment (creatinine clearance [CrCl] >30 mL/min) or end-stage renal disease (ESRD).

For patients with mild to moderate renal impairment (CrCl >30 mL/min), no dosage adjustment is required when using sofosbuvir, simeprevir, fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg), or fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) to treat or retreat HCV infection in patients with appropriate genotypes.

Rating: Class I, Level A

For patients with CrCl below 30 mL/min, treatment can be contemplated after consultation with an expert, because safety and efficacy data are not available for these patients.

Rating: Class IIb, Level C

Sofosbuvir enters the hepatocyte where it is metabolized to its active form, GS-461203. The downstream inactive nucleoside metabolite GS-331007 is almost exclusively eliminated from the body renally, mediated through a combination of glomerular filtration and active tubular secretion. Results of phase II and III clinical trials of sofosbuvir have excluded patients with serum creatinine levels greater than 2.5 mg/dL or creatinine clearance (CrCl) level less than 60 mL/min. The pharmacokinetics of a single 400 mg dose of sofosbuvir were assessed in persons not infected with HCV (study P7977-0915) with mild (estimated glomerular filtration rate [eGFR] >50 mL/min/1.73 m² and <80 mL/min/1.73 m²), moderate (eGFR >30 mL/min/1.73 m² and <50 mL/min/1.73 m²), or severe renal impairment (eGFR <30 mL/min/1.73 m²) and persons with end-stage renal disease (ESRD) requiring hemodialysis.

Relative to persons with normal renal function (eGFR >80 mL/min/1.73 m²), the sofosbuvir area under the

curve (AUC; 0-inf) was 61% higher, 107% higher, and 171% higher in subjects with mild, moderate, and severe renal impairment, respectively; GS-331007 AUC (0-inf) was 55% higher, 88% higher, and 451% higher, respectively. In subjects with ESRD (relative to subjects with normal renal function), sofosbuvir and GS-331007 AUC (0-inf) were 28% higher and 1280% higher, respectively, when sofosbuvir was dosed 1 hour before hemodialysis compared with 60% higher and 2070% higher, respectively, when sofosbuvir was dosed 1 hour after hemodialysis. No dosage adjustment is required for patients with mild or moderate renal impairment. The safety of sofosbuvir has not been established in patients with severe renal impairment or ESRD. Therefore, a dose recommendation cannot be provided for these populations at this time, although a dedicated study to evaluate optimal dosing of sofosbuvir in HCV-infected patients with severe renal impairment or ESRD on hemodialysis is currently underway.

No clinically relevant changes in ledipasvir pharmacokinetics were found between volunteers with normal renal function and those with severe renal impairment (eGFR <30 mL/min by Cockcroft-Gault) after a single dose of 90 mg of ledipasvir was administered.

Unique Patient Populations Table: Dose Adjustments Needed for Patients With Renal Impairment

Renal Impairment	eGFR / CrCl level (mL/min)	IFN	RBV	Sofosbuvir r	Ledipasvir r	Ombitasvir r	Dasabuvir r	Paritaprevir r	Simeprevir
Mild	50-80	180 ug PEG-IFN (2a); PEG-IFN (2b) 1.5 ug/Kg	Standard	Standard	Standard	Standard	Standard	Standard	Standard
Moderate	30-50	180ug PEG-IFN (2a); PEG-IFN (2b) 1 ug/kg (25% reduction)	Alternating doses 200 mg and 400 mg every other day	Standard	Standard	Standard	Standard	Standard	Standard
Severe	<30	135 ug PEG-IFN (2a); PEG-IFN (2b) 1 ug/kg (50% reduction)	200 mg/d	Data not available	Data not available	Data not available	Data not available	Data not available	Standard
ESRD/HD		PEG-IFN (2a) 135 µg/wk or PEG-IFN (2b) 1 µg/kg/wk or standard IFN 3 mU 3x/wk	200 mg/d	Data not available	Data not available	Data not available	Data not available	Data not available	Data not available

Abbreviations: CrCl, creatinine clearance; eGFR, estimated glomerular filtration rate; ESRD, end-stage renal disease; HD, hemodialysis.

Complete revision made to this section on December 19, 2014.

Unique Patient Populations: Renal Impairment Box. Summary of Recommendations for Patients with Renal Impairment Including Severe Renal Impairment (CrCl <30 ML/min) or ESRD Requiring Hemodialysis or Peritoneal Dialysis

Renal Impairment Box. Summary of Recommendations for Patients with Renal Impairment, Including Severe Renal Impairment (CrCl <30 ML/min) or ESRD Requiring Hemodialysis or Peritoneal Dialysis

Recommended dosage adjustments for patients with renal impairment, including severe renal impairment (creatinine clearance [CrCl] >30 mL/min) or end-stage renal disease (ESRD).

For patients with mild to moderate renal impairment (CrCl >30 mL/min), no dosage adjustment is required when using sofosbuvir, simeprevir, fixed-dose combination of ledipasvir (90 mg)/sofosbuvir (400 mg), or fixed-dose combination of paritaprevir (150 mg)/ritonavir (100 mg)/ombitasvir (25 mg) plus twice-daily dosed dasabuvir (250 mg) to treat or retreat HCV infection in patients with appropriate genotypes.

Rating: Class I, Level A

For patients with CrCl below 30 mL/min, treatment can be contemplated after consultation with an expert, because safety and efficacy data are not available for these patients.

Rating: Class IIb, Level C

MANAGEMENT OF ACUTE HCV INFECTION

Expansions and notes for abbreviations used in this section can be found in [Methods Table 3](#).

A summary of recommendations for Managing Acute HCV Infection is found in the [BOX](#).

This section provides guidance on the diagnosis and medical management of acute HCV infection, which is defined as presenting within 6 months of the exposure. During this time, there is a 20% to 50% chance of spontaneous resolution of infection. ([Kamal, 2008](#)) In the past, cure rates of acute infection with IFN-based treatment were very high. ([Grebel, 2014](#)) The present guidance reflects current trends transitioning toward safer, IFN-sparing treatments for chronic infection and the implications for the approach to acute hepatitis C treatment.

Acute HCV infection may result from exposure to the virus through a variety of routes. The highest risk is associated with repeated parenteral exposures from contaminated equipment in the setting of injection drug use (or user; IDU). Lower rates of HCV transmission occur in the setting of needlestick injuries in which health care workers are exposed to the blood of an HCV-infected patient. Heterosexual exposure risk is very low, but transmission rates are much higher among HIV-infected men who have sex with men (MSM) who have unprotected sex, particularly among those who engage in high-risk sexual practices that increase trauma to the mucosal membranes and exposure to blood, and possibly the presence of other ulcerative sexually transmitted infections. ([Boesecke, 2012](#))

Recommended testing for diagnosing acute HCV infection.

HCV antibody and HCV RNA testing are recommended when acute HCV infection is suspected due to exposure, clinical presentation, or elevated aminotransferase levels (see [Figure](#)).

Rating: Class I, Level C

Recommendations for HCV testing are also found in the HCV [Testing and Linkage to Care](#) section.

Diagnosis of acute infection permits estimation of annual incidence rates and transmission patterns, thereby facilitating implementation and assessment of prevention programs. At the individual level, a diagnosis of acute infection expedites linkage to care, counseling regarding high-risk behavior, and timely interventions to reduce transmission of infection and progression of liver disease. ([Bruneau, 2014](#)) Indeed, persons involved in high-risk behavior are known to practice serosorting, defined as using anti-HCV antibody serostatus, to determine whether to engage in high-risk behaviors with certain individuals. ([Smith, 2013](#)) Thus, undiagnosed acutely infected persons may be at greater risk of transmitting HCV to their presumably seronegative contacts than would be expected by chance.

The best laboratory evidence to support a diagnosis of acute HCV infection is (1) a positive HCV RNA test in the setting of a negative HCV antibody (identification during the seronegative “window” period), ([Cox,](#)

2005) or (2) a positive HCV antibody test after prior negative HCV antibody test (termed seroconversion). There are rare instances in which these approaches may be misleading, such as in immunosuppressed individuals with impaired antibody production. ([Chamot, 1990](#))

Discrete Exposure

The above types of clear laboratory documentation of acute infection are easiest to achieve when there has been a discrete exposure (eg, after new onset or a change in drug injection practice, a percutaneous needlestick exposure to an HCV-infected individual, a potentially nonsterile tattoo, or sexual assault). In those instances, baseline HCV antibody and RNA testing should be done within 48 hours of the exposure to document whether there was antecedent HCV infection (see [Figure](#)). If baseline testing is negative, repeat testing is recommended. Frequency of testing can be tailored based on management objectives (eg, monthly testing to identify and treat acute infection). If baseline anti-HCV antibody testing is positive but RNA testing is negative, repeat HCV RNA and alanine aminotransferase (ALT) testing is recommended to identify an acute reinfection. When baseline HCV antibody and RNA testing are both positive, the person most likely already has chronic HCV infection from prior exposures. The frequency of repeat testing should reflect management goals. At a minimum, repeat testing should be done 4 months to 6 months later. When earlier identification of infection or reinfection is desired, HCV RNA and ALT testing every 4 weeks to 6 weeks is recommended for 6 months.

No Discrete Exposure

Often, however, individuals suspected of having acute HCV infection do not have a discrete exposure or have no prior baseline testing, making a diagnosis of acute infection more difficult (see [Table](#) below). Acute infection should be suspected if there is a new rise in the ALT level without an alternate cause. ([Blackard, 2008](#)); ([Kim, 2013](#)) Acute infection should also be suspected when there are low (especially $<10^4$ IU/mL) or fluctuating ($>1 \log_{10}$ IU/mL) HCV RNA values, or spontaneous clearance, which do not commonly occur outside of the first 6 months after acute HCV infection. ([McGovern, 2009](#)) A low signal-to-cutoff ratio of HCV antibody along with detectable HCV RNA may also be suggestive of the early weeks of acute primary infection, although this information may need to be specifically requested from the testing laboratory. ([Araujo, 2011](#)) Patients suspected of having acute HCV infection should also have laboratory evaluation to exclude other or coexisting causes of acute hepatitis (eg, hepatitis A virus, hepatitis B virus, or autoimmune hepatitis) and should be tested for HIV.

Preexposure or postexposure prophylaxis with antiviral therapy is NOT recommended.

Rating: Class III, Level C

Although new antiviral treatment regimens are highly efficacious and more tolerable than IFN-based therapy, there are no data on the efficacy or cost-effectiveness of antiviral therapy for preexposure or postexposure prophylaxis of HCV infection. Some studies have shown that postexposure treatment with IFN-based regimens does not prevent infection. ([Nakano, 1995](#)); ([Arai, 1996](#))

Table. Interpretation of Blood Testing During Diagnosis of Acute HCV Infection

Test	Interpretation for Diagnosis of Acute HCV Infection
HCV antibody	<ul style="list-style-type: none">▪ May be negative in the first 6 weeks after exposure▪ May be delayed or absent when the individual is immunosuppressed▪ Presence alone does not distinguish between acute and chronic infection▪ Low signal-to-cutoff ratio may be present during acute HCV infection or represent a false-positive result
HCV RNA	<ul style="list-style-type: none">▪ Viral fluctuations greater than 1 log₁₀ IU/mL may indicate acute HCV infection▪ May be transiently negative during acute HCV infection▪ Alone does not distinguish between acute and chronic infection
Alanine aminotransferase (ALT)	<ul style="list-style-type: none">▪ Fluctuating peaks during acute HCV infection suggest acute infection▪ May be normal during acute HCV infection▪ May be elevated due to other liver insults such as alcohol consumption

Recommendations for medical management and monitoring in acute HCV infection.

Regular laboratory monitoring is recommended in the setting of acute HCV infection until the alanine aminotransferase (ALT) level normalizes and HCV RNA becomes repeatedly undetectable, suggesting spontaneous resolution.

Rating: Class I, Level B

Monitoring HCV RNA (eg, every 4 weeks to 8 weeks) for 6 months to 12 months is recommended to determine spontaneous clearance of HCV infection versus persistence of infection.

Rating: Class I, Level B

Counseling is recommended for patients with acute HCV infection to avoid hepatotoxic insults, including hepatotoxic drugs (eg, acetaminophen) and alcohol consumption, and to reduce the risk of HCV transmission to others.

Rating: Class I, Level C

Referral to an addiction medicine specialist is recommended for patients with acute HCV infection related to injection drug use.

Rating: Class I, Level B

The patient with acute HCV infection should be counseled to reduce behaviors that could result in transmission, such as sharing of injection equipment or high-risk sexual practices. Because the risk of transmission of other infections is higher in the acute infection phase, some experts counsel patients with acute infection to consider using barrier precautions even in stable monogamous relationships. (see [Testing and Linkage to Care](#)) For individuals with acute HCV infection who have a history of recent IDU, referral to an addiction medicine specialist is recommended when appropriate. ([Litwin, 2009](#)); ([Strathdee, 2005](#))

Patients with acute HCV infection are often asymptomatic or have nonspecific symptoms (fatigue, anorexia, mild or moderate abdominal pain, low-grade fever, nausea, vomiting) that frequently are not recognized as being associated with acute HCV infection. A small proportion (<25%) of patients with acute HCV infection will develop jaundice. Patients diagnosed with acute HCV infection should be initially monitored with hepatic panels (ALT, aspartate aminotransferase [AST], bilirubin, and international normalized ratio [INR] in the setting of increasing bilirubin) at 2- to 4-week intervals. ([Blackard, 2008](#)) Laboratory monitoring should continue until the ALT level normalizes and HCV RNA becomes undetectable, suggesting spontaneous resolution. If this does not occur, frequency of laboratory monitoring for patients with persistently detectable HCV RNA and elevated ALT level should follow recommendations for monitoring patients with chronic HCV infection. (see [Monitoring](#))

HCV infection will spontaneously clear in 20% to 50% of patients. ([Kamal, 2008](#)) In at least two-thirds of patients, this will occur within 6 months of the estimated time of infection (median 16.5 weeks); only 11% of those who remain viremic at 6 months will spontaneously clear infection at some later time. ([Grebely, 2014](#)) Thus, detectable HCV RNA at 6 months after the time of infection will identify most persons who need HCV therapy. (see [When and in Whom to Treat](#)) Those with spontaneous clearance should not be treated with antiviral therapy, but they should be counseled about the possibility of reinfection and tested routinely for reinfection if they engage in ongoing risk behaviors (see [Testing and Linkage to Care](#)). Of note, transient suppression of viremia can occur in those with acute HCV infection, even in those who progress to chronic infection. Thus, a single undetectable HCV RNA value is insufficient to declare spontaneous clearance. ([Villano, 1999](#)); ([Mosley, 2008](#)) (see [Testing and Linkage to Care](#))

Predictors of spontaneous clearance include jaundice, elevated ALT level, hepatitis B virus surface antigen (HBsAg) positivity, female sex, younger age, HCV genotype 1, and host genetic polymorphisms, most notably those near the IL28B gene. ([Kamal, 2008](#)); ([Mosley, 2008](#))

There is no need to alter concomitant medications that are metabolized by hepatic enzymes unless there is concern for developing acute liver failure (eg, increasing bilirubin level and INR). Acetaminophen and alcohol consumption should be avoided during acute HCV infection. ([Proeschold-Bell, 2012](#)); ([Dieperink, 2010](#)); ([Whitlock, 2004](#)) Hospitalization is rarely indicated unless nausea and vomiting are severe. Although acute liver failure is very rare (<1%), it represents a serious and life-threatening complication of acute HCV infection. Patients with an INR above 1.5 or those who exhibit any signs of acute liver failure (eg, hepatic encephalopathy) should be referred to a liver transplant center immediately. The use of HCV antiviral regimens in acute liver failure should be managed by a clinician experienced in HCV treatment, ideally in consultation with a liver transplant specialist.

Recommended treatment for patients with acute HCV infection.

If the practitioner and patient have decided that a delay in treatment initiation is acceptable, monitoring for spontaneous clearance is recommended for a minimum of 6 months. When the decision is made to initiate treatment after 6 months, treating as described for chronic hepatitis C is recommended (see [Initial Treatment of HCV Infection](#) and [When and in Whom to Treat](#))

Rating: Class IIa, Level C

If a decision has been made to initiate treatment during the acute infection period, monitoring HCV RNA for at least 12 weeks to 16 weeks is recommended to allow for spontaneous clearance before starting treatment.

Rating: Class IIa, Level C

Recommended regimens for patients with acute HCV infection.

Owing to high efficacy and safety, the same regimens recommended for chronic HCV infection (see [Initial Treatment of HCV Infection](#) and [When and in Whom to Treat](#)) are also recommended for acute infection.

Rating: Class IIa, Level C

Alternative regimen for patients with acute HCV infection who are eligible to receive IFN.

PEG with or without RBV for 16 weeks (for those with HCV genotype 2 or 3 who have a rapid virologic response) to 24 weeks (for those with HCV genotype 1).

Rating: Class II, Level A

For patients in whom HCV infection spontaneously clears, treatment is NOT recommended.

Rating: Class III, Level B

When the efficacy of the treatment of acute HCV infection (particularly for genotype 1) was superior to the treatment of chronic infection, there was a strong impetus to identify and treat acute HCV infection. (See 2009 AASLD guidelines, [\[Ghany, 2009\]](#)) The current availability of IFN-sparing HCV treatments that have high safety and efficacy for chronic HCV infection reduces (and possibly eliminates) the “efficacy advantage” of early treatment. Indeed, a randomized controlled study showed that delaying treatment was not inferior to early treatment, and many who received early IFN-based therapy were unable to complete treatment because of adverse effects. ([Deterding, 2013](#)) Until data documenting the efficacy and safety of treatment of acute hepatitis C with IFN-sparing therapy are available, monitoring for spontaneous clearance for minimum of 6 months before initiating treatment is recommended. When the decision is made to initiate treatment after 6 months, treating as described for chronic hepatitis C is recommended.

Some argue that the benefits of waiting until 6 months to document chronic hepatitis C and of using well-studied treatments for chronic hepatitis C outweigh the disadvantages of delaying treatment of acute infection for many patients. On the other hand, for some persons, there may be additional benefits of early treatment that include prevention of transmission to others (eg, IDUs or surgeons), prevention of severe complications (eg, someone with underlying compensated cirrhosis superinfected with acute HCV infection), and decreasing the chance of being lost to follow-up. If for these reasons, a decision has been made to initiate treatment during the acute infection period, the same regimens recommended for chronic HCV infection (see [Initial Treatment of HCV Infection](#) and [When and in Whom to Treat](#) sections) are also recommended for acute infection given their high efficacy and safety in chronic HCV infection.

Although data on direct-acting antiviral drugs are limited for acute HCV infection, there are data documenting the efficacy of IFN-based HCV treatment when given within 12 weeks to 16 weeks of infection, as described in the 2009 AASLD guidelines. ([Ghany, 2009](#)) This prior AASLD recommendation was later supported by a meta-analysis that reported sustained virologic response (SVR) in 82.5% of acutely infected patients who started IFN-based treatment within 12 weeks of diagnosis compared with SVR rates of 66.9% in those starting treatment between 12 weeks and 24 weeks and 62.5% in those starting treatment after 24 weeks. ([Corey, 2010](#)) Thus, clinicians should recognize that IFN-based treatment requires a balance between being prompt (treatment within 12 weeks to 16 weeks) and allowing enough time for spontaneous resolution. By analogy with studies of treatment of chronic infection, the duration of treatment of HCV acute genotype 2 or 3 infection with PEG-IFN and RBV may be shortened to 16 weeks for patients in whom a rapid virologic response is achieved. ([Dalgard, 2004](#)); ([Mangia, 2005](#)); ([von Wagner, 2005](#)); ([Yu, 2007](#)); ([Shiffman, 2007](#))

Mixed Genotypes

Rarely, genotyping assays may indicate the presence of a mixed infection (eg, genotypes 1a and 2). Treatment data for mixed genotypes with direct-acting antivirals are sparse, and awaiting availability of a pangentotypic regimen may be considered. Until then, when treatment is necessary, the choice of antiviral combination and duration of treatment should maximize efficacy against each genotype represented in the assay. When the correct combination or duration is unclear, expert consultation should be sought.

Changes made on February 2, 2015.

Acute Box. Recommendations for Management of Acute HCV Infection

Recommended testing for diagnosing acute HCV infection.

HCV antibody and HCV RNA testing are recommended when acute HCV infection is suspected due to exposure, clinical presentation, or elevated aminotransferase levels (see [Figure](#)).

Rating: Class I, Level C

Preexposure or postexposure prophylaxis with antiviral therapy is NOT recommended.

Rating: Class III, Level C

Recommendations for medical management and monitoring in acute HCV infection.

Regular laboratory monitoring is recommended in the setting of acute HCV infection until the alanine aminotransferase (ALT) level normalizes and HCV RNA becomes repeatedly undetectable, suggesting spontaneous resolution.

Rating: Class I, Level B

Monitoring HCV RNA (eg, every 4 weeks to 8 weeks) for 6 months to 12 months is recommended to determine spontaneous clearance of HCV infection versus persistence of infection.

Rating: Class I, Level B

Counseling is recommended for patients with acute HCV infection to avoid hepatotoxic insults, including hepatotoxic drugs (eg, acetaminophen) and alcohol consumption, and to reduce the risk of HCV transmission to others.

Rating: Class I, Level C

Referral to an addiction medicine specialist is recommended for patients with acute HCV infection related to injection drug use.

Rating: Class I, Level B

Recommended treatment for patients with acute HCV infection.

If the practitioner and patient have decided that a delay in treatment initiation is acceptable, monitoring for spontaneous clearance is recommended for a minimum of 6 months. When the decision is made to initiate treatment after 6 months, treating as described for chronic hepatitis C is recommended (see [Initial Treatment of HCV Infection](#) and [When and in Whom to Treat](#))

Rating: Class IIa, Level C

If a decision has been made to initiate treatment during the acute infection period, monitoring HCV RNA for at least 12 weeks to 16 weeks is recommended to allow for spontaneous clearance before starting treatment.

Rating: Class IIa, Level C

Recommended regimens for patients with acute HCV infection.

Owing to high efficacy and safety, the same regimens recommended for chronic HCV infection (see [Initial Treatment of HCV Infection](#) and [When and in Whom to Treat](#)) are also recommended for acute infection.

Rating: Class IIa, Level C

Alternative regimen for patients with acute HCV infection who are eligible to receive IFN.

PEG with or without RBV for 16 weeks (for those with HCV genotype 2 or 3 who have a rapid virologic response) to 24 weeks (for those with HCV genotype 1).


Rating: Class II, Level A

For patients in whom HCV infection spontaneously clears, treatment is NOT recommended.

Rating: Class III, Level B

Acute Figure. Testing Algorithm for Discrete Recognized Hepatitis C Virus (HCV) Exposure

Figure. Testing Algorithm for Discrete Recognized Hepatitis C Virus (HCV) Exposure^a



Baseline testing within 48 hours of exposure^e

^a Often there is no discrete exposure or the entry to health care occurs with jaundice or elevated liver enzymes. In those instances, baseline testing cannot be done and the diagnosis of acute infection is more challenging (see text).

^b Repeat HCV Ab is not needed if it is positive at baseline. Frequency of testing can be tailored based on management objectives (eg, monthly testing to identify and treat acute infection).

^c Some would treat after waiting 8 weeks to 12 weeks for spontaneous clearance (see text). Benefits of HCV antiviral therapy or IFN-based (alternative) within 12 weeks of acute infection are that this may decrease transmission risk to others (eg, among injection drug users or surgeons), prevent severe complications (eg, underlying cirrhosis superinfected with acute HCV infection), and minimize chance of being lost to follow-up.

^d If there were additional exposures in the preceding 6 months, a patient with a new diagnosis who is HCV RNA and HCV Ab positive may still be in the acute infection phase. Symptoms, high ALT level, or viral fluctuations may help distinguish acute from chronic HCV.

^e Baseline testing should be done within 48 hours of exposure to determine existing infection status: HCV RNA, HCV Ab, and ALT.

REFERENCES

AASLD/IDSA/IAS-USA. Recommendations for testing, managing, and treating hepatitis C. www.hcvguidelines.org. Accessed on January 29, 2014.

Aberg JA, Gallant JE, Ghanem KG, Emmanuel P, Zingman BS, Horberg MA. Primary Care Guidelines for the Management of Persons Infected With HIV: 2013 Update by the HIV Medicine Association of the Infectious Diseases Society of America. *Clin Infect Dis*. 2013.

Afdhal N, Zeuzem S, Kwo P, et al. Ledipasvir and sofosbuvir for untreated HCV genotype 1 infection. *N Engl J Med*. 2014a;370(20):1889-1898.

Afdhal N, Reddy KR, Nelson DR, et al. Ledipasvir and sofosbuvir for previously treated HCV genotype 1 infection. *N Engl J Med*. 2014b;370(16):1483-1493.

Agnello V, Chung RT, Kaplan LM. A role for hepatitis C virus infection in type II cryoglobulinemia. *N Engl J Med*. 1992;327(21):1490-1495.

Alter MJ, Kuhnert WL, Finelli L. Guidelines for laboratory testing and result reporting of antibody to hepatitis C virus. Centers for Disease Control and Prevention. *MMWR Recomm Rep*. 2003;52(RR-3):1-13, 15.

Alvarez D, Dieterich DT, Brau N, Moores L, Ball L, Sulkowski MS. Zidovudine use but not weight-based ribavirin dosing impacts anaemia during HCV treatment in HIV-infected persons. *J Viral Hepat*. 2006;13:683-689.

American Heart Association. http://my.americanheart.org/idc/groups/ahamah-public/@wcm/@sop/documents/downloadable/ucm_319826.pdf. Accessed on January 27, 2014.

Amon JJ, Garfein RS, Ahdieh-Grant L, et al. Prevalence of hepatitis C virus infection among injection drug users in the United States, 1994-2004. *Clin Infect Dis*. 2008;46(12):1852-1858.

Andreone P, Colombo MG, Enejosa JV, et al. ABT-450, ritonavir, ombitasvir, and dasabuvir achieves 97% and 100% sustained virologic response with or without ribavirin in treatment-experienced patients with HCV genotype 1b infection. *Gastroenterology*. 2014;147(2):359-365.

Arai Y, Noda K, Enomoto N, et al. A prospective study of hepatitis C virus infection after needlestick accidents. *Liver*. 1996;16(5):331-334.

Arase Y, Suzuki F, Suzuki Y, et al. Sustained virological response reduces incidence of onset of type 2 diabetes in chronic hepatitis C. *Hepatology*. 2009;49(3):739-744.

Araujo AC, Astrakhantseva IV, Fields HA, Kamili S. Distinguishing acute from chronic hepatitis C virus (HCV) infection based on antibody reactivities to specific HCV structural and nonstructural proteins. *J Clin Microbiol*. 2011;49(1):54-57.

Armstrong GL, Wasley A, Simard EP, McQuillan GM, Kuhnert WL, Alter MJ. The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann Intern Med.* 2006;144(10):705-714.

Arora S, Thornton K, Murata G, et al. Outcomes of treatment for hepatitis C virus infection by primary care providers. *N Engl J Med.* 2011;364(23):2199-2207.

Aspinall EJ, Corson S, Doyle JS, et al. Treatment of hepatitis C virus infection among people who are actively injecting drugs: a systematic review and meta-analysis. *Clin Infect Dis.* 2013;57(Suppl 2):S80-S89.

Backus LI, Boothroyd DB, Phillips BR, Belperio P, Halloran J, Mole LA. A sustained virologic response reduces risk of all-cause mortality in patients with hepatitis C. *Clin Gastroenterol Hepatol.* 2011;9(6):509-516.

Bacon BR, Gordon SC, Lawitz E, et al. Boceprevir for previously treated chronic HCV genotype 1 infection. *N Engl J Med.* 2011;364(13):1207-1217.

Bedossa P, Dargère D, Paradis V. Sampling variability of liver fibrosis in chronic hepatitis C. *Hepatology.* 2003;38(6):1449-1457.

Bellentani S, Pozzato G, Saccoccio G, et al. Clinical course and risk factors of hepatitis C virus related liver disease in the general population: report from the Dionysos study. *Gut.* 1999;44(6):874-880.

Benhamou Y, Bochet M, Di Martino V, et al. Liver fibrosis progression in human immunodeficiency virus and hepatitis C virus coinfecting patients. The Multivirc Group. *Hepatology.* 1999;30(4):1054-1058.

Benhamou Y, Di Martino V, Bochet M, et al. Factors affecting liver fibrosis in human immunodeficiency virus- and hepatitis C virus-coinfecting patients: impact of protease inhibitor therapy. *Hepatology.* 2001;34(2):283-287.

Berenguer J, Álvarez-Pellicer J, Martín PM, et al. Sustained virological response to interferon plus ribavirin reduces liver-related complications and mortality in patients coinfecting with human immunodeficiency virus and hepatitis C virus. *Hepatology.* 2009;50(2):407-413.

Berenguer M, Palau A, Aguilera V, Rayon JM, Juan FS, Prieto M. Clinical benefits of antiviral therapy in patients with recurrent hepatitis C following liver transplantation. *Am J Transplant.* 2008;8(3):679-687.

Berenguer M, Schuppan D. Progression of liver fibrosis in post-transplant hepatitis C: mechanisms, assessment and treatment. *J Hepatol.* 2013;58(5):1028-1041.

Blackard JT, Shata MT, Shire NJ, Sherman KE. Acute hepatitis C virus infection: a chronic problem. *Hepatology.* 2008;47(1):321-331.

Bochud PY, Cai T, Overbeck K, et al. Genotype 3 is associated with accelerated fibrosis progression in chronic hepatitis C. *J Hepatol.* 2009;51(4):655-666.

Boesecke C, Rockstroh JK. Acute hepatitis C in patients with HIV. *Semin Liver Dis.* 2012;32(2):130-137.

Bonder A, Afdhal N. Utilization of FibroScan in clinical practice. *Curr Gastroenterol Rep.* 2014;16(2):372.

Bonkovsky HL, Snow KK, Malet PF, et al. Health-related quality of life in patients with chronic hepatitis C and advanced fibrosis. *J Hepatol.* 2007;46(3):420-431.

Bourliere M, Bronowicki J, de Ledinghen V et al. Ledipasvir/sofosbuvir fixed dose combination is safe and efficacious in cirrhotic patients who have previously failed protease-inhibitor based triple therapy. [Abstract LB-6.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014a; Boston, MA.

Bourliere M, Sulkowski MS, Omata M et al. An integrated safety and efficacy analysis of >500 patients with compensated cirrhosis treated with ledipasvir/sofosbuvir with or without ribavirin. [Abstract 82.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014b; Boston, MA

Boursier J, de L, V, Zarski JP, et al. Comparison of eight diagnostic algorithms for liver fibrosis in hepatitis C: new algorithms are more precise and entirely noninvasive. *Hepatology.* 2012;55(1):58-67.

Bräu N, Salvatore M, Ríos-Bedoya CF, et al. Slower fibrosis progression in HIV/HCV-coinfected patients with successful HIV suppression using antiretroviral therapy. *J Hepatol.* 2006;44:47-55.

Bravo MJ, Vallejo F, Barrio G, et al. HCV seroconversion among never-injecting heroin users at baseline: no predictors identified other than starting injection. *Int J Drug Policy.* 2012;23(5):415-419.

Bruggmann P, Litwin AH. Models of care for the management of hepatitis C virus among people who inject drugs: one size does not fit all. *Clin Infect Dis.* 2013;57 Suppl 2:S56-S61.

Bruix J, Sherman M. Management of hepatocellular carcinoma: an update. *Hepatology.* 2011;53(3):1020-1022.

Bruneau J, Zang G, Abrahamowicz M, Jutras-Aswad D, Daniel M, Roy E. Sustained drug use changes after hepatitis C screening and counseling among recently infected persons who inject drugs: a longitudinal study. *Clin Infect Dis.* 2014;58(6):755-761.

Butt AA, McGinnis K, Skanderson M, Justice AC. A comparison of treatment eligibility for hepatitis C virus in HCV-monoinfected versus HCV/HIV-coinfected persons in electronically retrieved cohort of HCV-infected veterans. *AIDS Res Hum Retroviruses.* 2011;27(9):973-979.

Castera L. Noninvasive methods to assess liver disease in patients with hepatitis B or C. *Gastroenterology.* 2012;142(6):1293-1302.

Castera L, Sebastiani G, Le BB, de L, V, Couzigou P, Alberti A. Prospective comparison of two algorithms combining non-invasive methods for staging liver fibrosis in chronic hepatitis C. *J*

Hepatol. 2010;52(2):191-198.

Centers for Disease Control and Prevention. Recommendations for preventing transmission of infections among chronic hemodialysis patients. *MMWR Recomm Rep.* 2001;50(RR-5):1-43.

Centers for Disease Control and Prevention. Recommendations for prevention and control of hepatitis C virus (HCV) infection and HCV-related chronic disease. *MMWR Morb Mortal Wkly Rep.* 1998;47(RR-19):1-39.

Centers for Disease Control and Prevention. Recommendations for identification and public health management of persons with chronic hepatitis B virus infection. *MMWR.* 2008;57(RR-8)

Centers for Disease Control and Prevention (CDC). Testing for HCV infection: an update of guidance for clinicians and laboratorians. *MMWR Morb Mortal Wkly Rep.* 2013;62(18):362-365.

Chamot E, Hirschel B, Wintch J, et al. Loss of antibodies against hepatitis C virus in HIV-seropositive intravenous drug users. *AIDS.* 1990;4(12):1275-1277.

Charlton M, Seaberg E, Wiesner R, et al. Predictors of patient and graft survival following liver transplantation for hepatitis C. *Hepatology.* 1998;28(3):823-830.

Charlton MR, Gane E, Manns M, et al. Sofosbuvir and Ribavirin for the Treatment of Established Recurrent Hepatitis C Infection After Liver Transplantation: Preliminary Results of a Prospective, Multicenter Study. *Hepatology: Special Issue: The 64th Annual Meeting of the American Association for the Study of Liver Diseases: The Liver Meeting 2013.* 2013;Vol 58(4):1378A.

Charlton MR. E-mail communication. August 14, 2014.

Charlton M, Gane E, Manns MP, et al. Sofosbuvir and ribavirin for treatment of compensated recurrent hepatitis C virus infection after liver transplantation. *Gastroenterology.* 2014b.

Chen TY, Ding EL, Seage Iii GR, Kim AY. Meta-analysis: increased mortality associated with hepatitis C in HIV-infected persons is unrelated to HIV disease progression. *Clin Infect Dis.* 2009;49(10):1605-1615.

Chew KW, Allen SA, Taylor LE, Rich JD, Feller E. Treatment outcomes with pegylated interferon and ribavirin for male prisoners with chronic hepatitis C. *J Clin Gastroenterol.* 2009;43(7):686-691.

Chou R, Wasson N. Blood tests to diagnose fibrosis or cirrhosis in patients with chronic hepatitis C virus infection. *Ann Intern Med.* 2013b;159(5):372.

Chou R, Wasson N. Blood tests to diagnose fibrosis or cirrhosis in patients with chronic hepatitis C virus infection: a systematic review. *Ann Intern Med.* 2013;158(11):807-820.

Chu CJ, Lee SD. Hepatitis B virus/hepatitis C virus coinfection: epidemiology, clinical features, viral interactions and treatment. *J Gastroenterol Hepatol.* 2008;23(4):512-520.

Clark BT, Garcia-Tsao G, Fraenkel L. Patterns and predictors of treatment initiation and completion in patients with chronic hepatitis C virus infection. *Patient Prefer Adherence.* 2012;6:285-295.

Coilly A, Roche B, Dumortier J, et al. Safety and efficacy of protease inhibitors to treat hepatitis C after liver transplantation: A multicenter experience. *J.Hepatol.* 2014;60(1):78-86.

Coppola N, De PS, Pisaturo M, et al. Sustained virological response to antiviral treatment in chronic hepatitis C patients may be predictable by HCV-RNA clearance in peripheral blood mononuclear cells. *J Clin Virol.* 2013;58(4):748-750.

Corey KE, Mendez-Navarro J, Gorospe EC, Zheng H, Chung RT. Early treatment improves outcomes in acute hepatitis C virus infection: a meta-analysis. *J Viral Hepat.* 2010;17(3):201-207.

Corrao G, Arico S. Independent and combined action of hepatitis C virus infection and alcohol consumption on the risk of symptomatic liver cirrhosis. *Hepatology.* 1998;27(4):914-919.

Cox AL, Netski DM, Mosbrugger T, et al. Prospective evaluation of community-acquired acute-phase hepatitis C virus infection. *Clin Infect Dis.* 2005;40(7):951-958.

Crippin JS, McCashland T, Terrault N, Sheiner P, Charlton MR. A pilot study of the tolerability and efficacy of antiviral therapy in hepatitis C virus-infected patients awaiting liver transplantation. *Liver Transpl.* 2002;8(4):350-355.

Curry MP, et al. Pretransplant sofosbuvir and ribavirin to prevent recurrence of HCV infection after liver transplantation. 64th Annual Meeting of the American Association for the Study of Liver Disease. Nov 1-5, 2013, 2013; Washington, DC.

Curry M, Forns X, Chung RT, et al. Pretransplant sofosbuvir and ribavirin to prevent recurrence of HCV infection after liver transplantation. *Hepatology.* 2013b;58(S1):313A-317A.

Curry MP, Forns X, Chung RT, et al. Sofosbuvir and Ribavirin Prevent Recurrence of HCV Infection After Liver Transplantation: An Open-Label Study. *Gastroenterology.* 2014;148(1):100-107.

Dalgard O, Bjoro K, Hellum KB, et al. Treatment with pegylated interferon and ribavirin in HCV infection with genotype 2 or 3 for 14 weeks: a pilot study. *Hepatology.* 2004;40(6):1260-1265.

de Kanter C, Blonk M, Colbers A et al. The influence of the HCV protease inhibitor boceprevir on the pharmacokinetics of the HIV integrase inhibitor raltegravir [Abstract 772LB]. 19th Conference on Retroviruses and Opportunistic Infections (CROI). March 5-8, 2012; Seattle, WA.

de Ledingham V, Barreiro P, Foucher J, et al. Liver fibrosis on account of chronic hepatitis C is more severe in HIV-positive than HIV-negative patients despite antiretroviral therapy. *J Viral Hepat.* 2008;15(6):427-433.

Del Bello DP, Bichoupan K, Yalamanchili R et al. Real-world data on HIV positive patients with HCV genotype 1,2 and 3 on sofosbuvir- and simeprevir containing regimens. [Abstract 994.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-1, 2014; Boston, MA.

Denniston MM, Klevens RM, McQuillan GM, Jiles RB. Awareness of infection, knowledge of hepatitis C, and medical follow-up among individuals testing positive for hepatitis C: National Health and Nutrition Examination Survey 2001-2008. *Hepatology.* 2012;55(6):1652-1661.

Denniston MM, Jiles RB, Drobeniuc J, et al. Chronic hepatitis C virus infection in the United States, National Health and Nutrition Examination Survey 2003 to 2010. *Ann Intern Med*. 2014;160(5):293-300.

Deterding K, Gruner N, Buggisch P, et al. Delayed versus immediate treatment for patients with acute hepatitis C: a randomised controlled non-inferiority trial. *Lancet Infect Dis*. 2013;13(6):497-506.

Di Bisceglie A, Kuo A, Rustgi V, Sulkowski M. Virological Outcomes and Adherence to Treatment Algorithms in a Longitudinal Study of Patients with Chronic Hepatitis C Treated with Boceprevir or Telaprevir in the U.S. (HCV-TARGET). 64th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 1-5, 2013; Washington, DC.

Di Bisceglie AM, Shiffman ML, Everson GT, et al. Prolonged therapy of advanced chronic hepatitis C with low-dose peginterferon. *N Engl J Med*. 2008;359(23):2429-2441.

Dienstag JL, Ghany MG, Morgan TR, et al. A prospective study of the rate of progression in compensated, histologically advanced chronic hepatitis C. *Hepatology*. 2011;54(2):396-405.

Dieperink E, Ho SB, Heit S, Durfee JM, Thurs P, Willenbring ML. Significant reductions in drinking following brief alcohol treatment provided in a hepatitis C clinic. *Psychosomatics*. 2010;51(2):149-156.

Dieterich D, Rockstroh J, Orkin C, et al. Simeprevir (TMC435) plus peginterferon/ribavirin in patients co-infected with HCV genotype-1 and HIV-1: primary analysis of the C212 study. 14th European AIDs Conference (EACS 2013). Oct 16-19, 2013, 2013; Brussels.

Dieterich D, Bacon B, Flamm SL et al. Evaluation of sofosbuvir and simeprevir-based regimens in the TRIO network: academic and community treatment of a real-world, heterogeneous population. [Abstract 46.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014a;220A; Boston, MA.

Dieterich D, Rockstroh JK, Orkin C, et al. Simeprevir (TMC435) With Pegylated Interferon/Ribavirin in Patients Coinfected With HCV Genotype 1 and HIV-1: A Phase 3 Study. *Clin Infect Dis*. 2014b;59(11):1579-1587.

Dieterich D, Nelson M, Soriano V, et al. Faldaprevir and pegylated interferon alpha-2a/ribavirin in individuals co-infected with hepatitis C virus genotype-1 and HIV. *AIDS*. 2015;29(5):571-581.

Durier N, Nguyen C, White LJ. Treatment of hepatitis C as prevention: a modeling case study in Vietnam. *PLoS One*. 2012;7(4):e34548.

Eron JJ, Lalezari J, Slim J, et al. Safety and efficacy of ombitasvir - 450/r and dasabuvir and ribavirin in HCV/HIV-1 co-infected patients receiving atazanavir or raltegravir ART regimens. *J Int AIDS Soc*. 2014;17(4 Suppl 3):19500.

Esmat GE, Shiha G, Omar RF et al. Sofosbuvir plus ribavirin in the treatment of egyptian patients with chronic genotype 4 HCV infection. [Abstract 959.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.

Everhart JE, Lok AS, Kim HY, et al. Weight-related effects on disease progression in the hepatitis C antiviral long-term treatment against cirrhosis trial. *Gastroenterology*. 2009;137(2):549-557.

Everhart JE, Wright EC, Goodman ZD, et al. Prognostic value of Ishak fibrosis stage: findings from the hepatitis C antiviral long-term treatment against cirrhosis trial. *Hepatology*. 2010;51(2):585-594.

Everson GT. Treatment of patients with hepatitis C virus on the waiting list. *Liver Transpl*. 2003;9(11):S90-S94.

Everson GT, Hoefs JC, Seeff LB, et al. Impact of disease severity on outcome of antiviral therapy for chronic hepatitis C: Lessons from the HALT-C trial. *Hepatology*. 2006;44(6):1675-1684.

Everson GT, Trotter J, Forman L, et al. Treatment of advanced hepatitis C with a low accelerating dosage regimen of antiviral therapy. *Hepatology*. 2005;42(2):255-262.

Fabrizi F, Dixit V, Messa P. Antiviral therapy of symptomatic HCV-associated mixed cryoglobulinemia: meta-analysis of clinical studies. *J Med Virol*. 2013;85(6):1019-1027.

Fabrizi F, Martin P, Dixit V, Messa P. Meta-analysis of observational studies: hepatitis C and survival after renal transplant. *J Viral Hepat*. 2014;21(5):314-324.

Fabrizi F, Messa P, Martin P. Health-related quality of life in dialysis patients with HCV infection. *Int J Artif Organs*. 2009;32(8):473-481.

Fabrizi F, Poordad FF, Martin P. Hepatitis C infection and the patient with end-stage renal disease. *Hepatology*. 2002;36(1):3-10.

Fabrizi F, Takkouche B, Lunghi G, Dixit V, Messa P, Martin P. The impact of hepatitis C virus infection on survival in dialysis patients: meta-analysis of observational studies. *J Viral Hepat*. 2007;14(10):697-703.

Feld JJ, Liang TJ. Hepatitis C -- identifying patients with progressive liver injury. *Hepatology*. 2006;43(2 Suppl 1):S194-S206.

Feld JJ, Kowdley KV, Coakley E, et al. Treatment of HCV with ABT-450/r-ombitasvir and dasabuvir with ribavirin. *N Engl J Med*. 2014;370(17):1594-1603.

Ferenci P, Bernstein D, Lalezari J, et al. ABT-450/r-ombitasvir and dasabuvir with or without ribavirin for HCV. *N Engl J Med*. 2014;370(21):1983-1992.

Fierer DS, Dieterich DT, Fiel MI, et al. Rapid progression to decompensated cirrhosis, liver transplant, and death in HIV-infected men after primary hepatitis C virus infection. *Clin Infect Dis*. 2013;56(7):1038-1043.

Fierer DS, Uriel AJ, Carriero DC, et al. Liver fibrosis during an outbreak of acute hepatitis C virus infection in HIV-infected men: a prospective cohort study. *J Infect Dis*. 2008;198(5):683-686.

Finelli L, Miller JT, Tokars JI, Alter MJ, Arduino MJ. National surveillance of dialysis-associated diseases in the United States, 2002. *Semin Dial*. 2005;18(1):52-61.

Fissell RB, Bragg-Gresham JL, Woods JD, et al. Patterns of hepatitis C prevalence and seroconversion in hemodialysis units from three continents: the DOPPS. *Kidney Int.* 2004;65(6):2335-2342.

Flamm SL, Everson GT, Charlton M et al. Ledipasvir/sofosbuvir with ribavirin for the treatment of HCV in patients with decompensated cirrhosis: preliminary results of a prospective, multicenter study. 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 1-5, 2014; Boston, MA.

Fleischer R, Boxwell D, Sherman KE. Nucleoside analogues and mitochondrial toxicity. *Clin Infect Dis.* 2004;38:e79-e80.

Fontana RJ, Sanyal AJ, Ghany MG, et al. Factors that determine the development and progression of gastroesophageal varices in patients with chronic hepatitis C. *Gastroenterology.* 2010;138(7):2321-31, 2331.

Forman LM, Lewis JD, Berlin JA, Feldman HI, Lucey MR. The association between hepatitis C infection and survival after orthotopic liver transplantation. *Gastroenterology.* 2002;122:889-896.

Forns X, Fontana RJ, Moonka D, et al. Initial Evaluation of the Sofosbuvir Compassionate Use Program for Patients with Severe Recurrent HCV Following Liver Transplantation. 64th Annual Meeting of the American Association for the Study of Liver Diseases: The Liver Meeting 2013. Epub Oct 1, 2013, 10-1-2013a;732A.

Forns X, Lawitz E, Zeuzem S, et al. Simeprevir (TMC435) with peg-interferon a-2a/ribavirin for treatment of chronic HCV genotype 1 infection in patients who relapsed after previous interferon-based therapy: efficacy and safety in patient sub-populations in the PROMISE phase III trial [Abstract 1092]. 64th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 1-5, 2013b; Washington, DC.

Forns X, Fontana RJ, Moonka D, McHutchison JG, Symonds WT. Initial evaluation of the sofosbuvir compassionate use program for patients with severe recurrent HCV following liver transplantation. *Hepatology.* 2013c;58(S1):730A-760A.

Forns X, Navasa M, Rodes J. Treatment of HCV infection in patients with advanced cirrhosis. *Hepatology.* 2004;40(2):498.

Foster GR, Goldin RD, Thomas HC. Chronic hepatitis C virus infection causes a significant reduction in quality of life in the absence of cirrhosis. *Hepatology.* 1998;27(1):209-212.

Foster GR, Hezode C, Bronowicki JP, et al. Telaprevir alone or with peginterferon and ribavirin reduces HCV RNA in patients with chronic genotype 2 but not genotype 3 infections. *Gastroenterology.* 2011;141(3):881-889.

Gane EJ, Stedman CA, Hyland RH, et al. Nucleotide polymerase inhibitor sofosbuvir plus ribavirin for hepatitis C. *N Engl J Med.* 2013a;368(1):34-44.

Gane EJ, Stedman CA, Hyland RH, et al. Once daily sofosbuvir plus ribavirin for 12 and 24 weeks in treatment-naive patients with HCV infection: the QUANTUM study. [Abstract Abstract 14.]

Program and abstracts of the 48th Annual Meeting of the European Association for the Study of the Liver. April 24-28, 2013b; Amsterdam, the Netherlands.

Gane EJ, Hyland RH, An D et al. High efficacy of LDV/SOF regimens for 12 weeks for patients with HCV genotype 3 or 6 infection. [Abstract LB11.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.

Garcia-Bengoechea M, Basaras M, Barrio J, et al. Late disappearance of hepatitis C virus RNA from peripheral blood mononuclear cells in patients with chronic hepatitis C in sustained response after alpha-interferon therapy. *Am J Gastroenterol*. 1999;94(7):1902-1905.

Garcia-Tsao G, Sanyal AJ, Grace ND, Carey W. Prevention and management of gastroesophageal varices and variceal hemorrhage in cirrhosis. *Hepatology*. 2007;46(3):922-938.

Garraffo R, Poizot-Martin I, Piroth L et al. Pharmacokinetic (PK) interactions between Boceprevir (BOC) and Atazanavir/r (ATV/r) or Raltegravir (RAL) in HIV/HCV coinfecting patients (pts). 14th International Workshop on Clinical Pharmacology on HIV Therapy. April 22-24, 2013; Amsterdam, the Netherlands.

Genentech Inc. COPEGUS (ribavirin) package insert. 2014a.
<http://dailymed.nlm.nih.gov/dailymed/lookup.cfm?setid=d370635f-5530-4d42-a019-d76b61639787>. Accessed on August 28, 2014.

Genentech Inc. PEGASYS (peginterferon alfa-2a) package Insert. 2014b.
<http://dailymed.nlm.nih.gov/dailymed/lookup.cfm?setid=de61685e-2b8c-4e22-84bb-869e13600440>. Accessed on August 28, 2014.

George SL, Bacon BR, Brunt EM, Mihindukulasuriya KL, Hoffmann J, Di Bisceglie AM. Clinical, virologic, histologic, and biochemical outcomes after successful HCV therapy: a 5-year follow-up of 150 patients. *Hepatology*. 2009;49(3):729-738.

German P, Pang P, West S et al. Drug interactions between direct acting anti-HCV antivirals sofosbuvir and ledipasvir and HIV antiretrovirals. [Abstract 06.] 15th International Workshop on Clinical Pharmacology of HIV and Hepatitis Therapy. May 19-21, 2014; Washington, DC.

Ghany MG, Kleiner DE, Alter H, et al. Progression of fibrosis in chronic hepatitis C. *Gastroenterology*. 2003;124(1):97-104.

Ghany MG, Nelson DR, Strader DB, Thomas DL, Seeff LB. An update on treatment of genotype 1 chronic hepatitis C virus infection: 2011 practice guideline by the American Association for the Study of Liver Diseases. *Hepatology*. 2011;54(4):1433-1444.

Ghany MG, Strader DB, Thomas DL, Seeff LB. Diagnosis, management, and treatment of hepatitis C: an update. *Hepatology*. 2009;49(4):1335-1374.

Gilead Sciences, Inc. Sofosbuvir [package insert]. 2013. Foster City, CA, Gilead Sciences, Inc.

Gilead Sciences Inc. Harvoni (ledipasvir and sofosbuvir) [package insert]. 2014. Foster City, CA, Gilead Sciences, Inc.

Gisbert JP, Garcia-Buey L, Pajares JM, Moreno-Otero R. Prevalence of hepatitis C virus infection in porphyria cutanea tarda: systematic review and meta-analysis. *J Hepatol.* 2003;39(4):620-627.

Gisbert JP, Garcia-Buey L, Pajares JM, Moreno-Otero R. Systematic review: regression of lymphoproliferative disorders after treatment for hepatitis C infection. *Aliment Pharmacol Ther.* 2005;21(6):653-662.

Govindasamy D, Ford N, Kranzer K. Risk factors, barriers and facilitators for linkage to antiretroviral therapy care: a systematic review. *AIDS.* 2012;26(16):2059-2067.

Grady BP, Schinkel J, Thomas XV, Dalgard O. Hepatitis C virus reinfection following treatment among people who use drugs. *Clin Infect Dis.* 2013;57(Suppl 2):S105-S110.

Grebely J, Page K, Sacks-Davis R, et al. The effects of female sex, viral genotype, and IL28B genotype on spontaneous clearance of acute hepatitis C virus infection. *Hepatology.* 2014;59(1):109-120.

Gumber SC, Chopra S. Hepatitis C: a multifaceted disease. Review of extrahepatic manifestations. *Ann Intern Med.* 1995;123(8):615-620.

Gunthard HF, Aberg JA, Eron JJ, et al. Antiretroviral treatment of adult HIV infection: 2014 recommendations of the International Antiviral Society-USA panel. *JAMA.* 2014;312(4):410-425.

Hammond KP, Wolfe P, Burton JR, Jr., et al. Pharmacokinetic interaction between boceprevir and etravirine in HIV/HCV seronegative volunteers. *J Acquir Immune Defic Syndr.* 2013;62(1):67-73.

Harris DR, Gonin R, Alter HJ, et al. The relationship of acute transfusion-associated hepatitis to the development of cirrhosis in the presence of alcohol abuse. *Ann Intern Med.* 2001;134(2):120-124.

Hellard ME, Jenkinson R, Higgs P, et al. Modelling antiviral treatment to prevent hepatitis C infection among people who inject drugs in Victoria, Australia. *Med J Aust.* 2012;196(10):638-641.

Hermine O, Lefrere F, Bronowicki JP, et al. Regression of splenic lymphoma with villous lymphocytes after treatment of hepatitis C virus infection. *N Engl J Med.* 2002;347(2):89-94.

Holmberg SD, Spradling PR, Moorman AC, Denniston MM. Hepatitis C in the United States. *N Engl J Med.* 2013;368(20):1859-1861.

Holmes HM, Hayley DC, Alexander GC, Sachs GA. Reconsidering medication appropriateness for patients late in life. *Arch Intern Med.* 2006;166(6):605-609.

Hosein SR, Wilson DP. HIV, HCV, and drug use in men who have sex with men. *Lancet.* 2013;382(9898):1095-1096.

Hourigan LF, Macdonald GA, Purdie D, et al. Fibrosis in chronic hepatitis C correlates significantly with body mass index and steatosis. *Hepatology.* 1999;29(4):1215-1219.

Hsu L, Bowlus CL, Stewart SL, et al. Electronic messages increase hepatitis B screening in at-risk Asian American patients: a randomized, controlled trial. *Dig Dis Sci.* 2013;58(3):807-814.

Hsu YC, Lin JT, Ho HJ, et al. Antiviral treatment for hepatitis C virus infection is associated with improved renal and cardiovascular outcomes in diabetic patients. *Hepatology*. 2014;59(4):1293-1302.

Hulskotte EG, Feng HP, Xuan F, et al. Pharmacokinetic interactions between the hepatitis C virus protease inhibitor boceprevir and ritonavir-boosted HIV-1 protease inhibitors atazanavir, darunavir, and lopinavir. *Clin Infect Dis*. 2013;56(5):718-726.

Hung CH, Wang JH, Hu TH, et al. Insulin resistance is associated with hepatocellular carcinoma in chronic hepatitis C infection. *World J Gastroenterol*. 2010;16(18):2265-2271.

Islam MM, Topp L, Conigrave KM, et al. Linkage into specialist hepatitis C treatment services of injecting drug users attending a needle syringe program-based primary healthcare centre. *J Subst Abuse Treat*. 2012;43(4):440-445.

Jacobson IM, Dore GJ, Foster G, et al. Simeprevir (TMC435) with Peginterferon/ Ribavirin for Chronic HCV Genotype-1 Infection in Treatment-Naive Patients: Results From QUEST-1, a Phase III Trial. Digestive Disease Week. May 18-21, 2013a; Orlando, FL.

Jacobson IM, Ghalib RH, Rodriguez-Torres M, et al. SVR results of a once-daily regimen of simeprevir (TMC435) plus sofosbuvir (GS-7977) with or without ribavirin in cirrhotic and non-cirrhotic HCV genotype 1 treatment-naive and prior null responder patients: The COSMOS study. [Abstract LB3.] 64th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 1-5, 2013b; Washington, DC

Jacobson IM, Gordon SC, Kowdley KV, et al. Sofosbuvir for hepatitis C genotype 2 or 3 in patients without treatment options. *N Engl J Med*. 2013c;368(20):1867-1877.

Jadoul M, Cornu C, van Ypersele de SC. Universal precautions prevent hepatitis C virus transmission: a 54 month follow-up of the Belgian Multicenter Study. The Universitaires Cliniques St-Luc (UCL) Collaborative Group. *Kidney Int*. 1998;53(4):1022-1025.

Janssen Therapeutics. Simeprevir [package insert]. 2013. Titusville, NJ, Janssen Therapeutics.

Janssen R&D. A study of pharmacokinetics, efficacy, safety, tolerability, of the combination of simeprevir (TMC435), daclatasvir (BMS-790052), and ribavirin (RBV) in patients with recurrent chronic hepatitis C genotype 1b infection after orthotopic liver transplantation (posted 2013b). <http://www.clinicaltrials.gov/ct2/show/NCT01938625>. Accessed on September 25, 2014.

Jensen DM, O'Leary JG, Pockros P et al. Safety and efficacy of sofosbuvir-containing regimens for hepatitis C: real-world experience in a diverse, longitudinal observational cohort. [Abstract 45.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA

Johnson M, Borland J, Chen S-J et al. Dolutegravir, Boceprevir, Telaprevir PK: The effect of Boceprevir and Telaprevir on Dolutegravir Pharmacokinetics, in Healthy Adult Subjects. 14th International Workshop on Clinical Pharmacology of HIV Therapy. Apr 22-24, 2013, 2013; Amsterdam.

Johnson RJ, Gretch DR, Couser WG, et al. Hepatitis C virus-associated glomerulonephritis. Effect of alpha-interferon therapy. *Kidney Int.* 1994;46(6):1700-1704.

Johnson RJ, Gretch DR, Yamabe H, et al. Membranoproliferative glomerulonephritis associated with hepatitis C virus infection. *N Engl J Med.* 1993;328(7):465-470.

Kakuda T, Leopold L, Nijs S. Pharmacokinetic interaction between etravirine or rilpivirine and telaprevir in healthy volunteers: a randomised, two-way crossover trial [Abstract O-18]. 13th International Workshop on Clinical Pharmacology of HIV Therapy. April 16-18, 2012; Barcelona, Spain.

Kamal SM. Acute hepatitis C: a systematic review. *Am J Gastroenterol.* 2008;103(5):1283-1297.

Kanwal F, Kramer JR, Ilyas J, Duan Z, El-Serag HB. HCV genotype 3 is associated with an increased risk of cirrhosis and hepatocellular cancer in a national sample of U.S. Veterans with HCV. *Hepatology.* 2014;60(1):98-105.

Kapoor R, Kohli A, Sidharthan S et al. All oral treatment for genotype 4 chronic hepatitis C infection with sofosbuvir and ledipasvir: interim results from the NIAID SYNERGY trial. [Abstract 240.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.

Kasserra C, Hughes E, Treitel M, Gupta S, O'Mara E. Clinical Pharmacology of BOC: Metabolism, Excretion, and Drug-Drug Interactions. [Abstract 118.] 18th Conference on Retroviruses and Opportunistic Infections (CROI): February 27-March 2. 2011; Boston, MA.

KDIGO. KDIGO clinical practice guidelines for the prevention, diagnosis, evaluation, and treatment of hepatitis C in chronic kidney disease. *Kidney Int Suppl.* 2008;(109):S1-99.

Khatri A, Wang T, Wang H et al. Drug-drug interactions of the direct-acting antiviral regimen of ABT-450/r, ombitasvir, and dasabuvir with HIV protease inhibitors. [Abstract 484.] 54th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC). September 5-9, 2014a; Washington, DC.

Khatri A, Wang T, Wang H et al. Drug-drug interactions of the direct-acting antiviral regimen of ABT-450/r, ombitasvir, and dasabuvir with emtricitabine + tenofovir, raltegravir, rilpivirine, and efavirenz. [Abstract 483.] 54th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC). September 5-9, 2014b; Washington, DC.

Khokhar OS, Lewis JH. Reasons why patients infected with chronic hepatitis C virus choose to defer treatment: do they alter their decision with time? *Dig Dis Sci.* 2007;52(5):1168-1176.

Kim AY, Nagami EH, Birch CE, Bowen MJ, Lauer GM, McGovern BH. A simple strategy to identify acute hepatitis C virus infection among newly incarcerated injection drug users. *Hepatology.* 2013;57(3):944-952.

Kirby B, Mathias A, Rossi S, et al. No clinically significant pharmacokinetic drug interactions between sofosbuvir (GS-7977) and HIV antiretrovirals atripla, rilpivirine, darunavir/ritonavir, or raltegravir in healthy volunteers. 63rd Annual Meeting of the American Association of the Study of

Liver Diseases (AASLD). November 9-11, 2012; Boston, MA.

Kirk GD, Mehta SH, Astemborski J, et al. HIV, age, and the severity of hepatitis C virus-related liver disease: a cohort study. *Ann Intern Med.* 2013;158(9):658-666.

Kiser JJ, Burton JR, Jr., Everson GT. Drug-drug interactions during antiviral therapy for chronic hepatitis C. *Nat Rev Gastroenterol Hepatol.* 2013;10(10):596-606.

Kleiner DE. The liver biopsy in chronic hepatitis C: a view from the other side of the microscope. *Semin Liver Dis.* 2005;25(1):52-64.

Kohler JJ, Nettles JH, Amblard F, et al. Approaches to hepatitis C treatment and cure using NS5A inhibitors. *Infect Drug Resist.* 2014;7:41-56.

Kowdley KV, Gordon SC, Reddy KR, et al. Ledipasvir and sofosbuvir for 8 or 12 weeks for chronic HCV without cirrhosis. *N Engl J Med.* 2014;370(20):1879-1888.

Konerman MA, Mehta SH, Sutcliffe CG, et al. Fibrosis progression in human immunodeficiency virus/hepatitis C virus coinfecting adults: prospective analysis of 435 liver biopsy pairs. *Hepatology.* 2014;59(3):767-775.

Lai JC, Kahn JG, Tavakol M, Peters MG, Roberts JP. Reducing infection transmission in solid organ transplantation through donor nucleic Acid testing: a cost-effectiveness analysis. *Am J Transplant.* 2013;13(10):2611-2618.

Lalezari L, Nelson DR, Hyland RH, et al. Once daily sofosbuvir plus ribavirin for 12 and 24 weeks in treatment-naive patients with HCV infection: the QUANTUM study. Program and abstracts of the 48th Annual Meeting of the European Association for the Study of the Liver. April 24-28, 2013, 2013; Amsterdam, The Netherlands.

Lambers FA, Prins M, Thomas X, et al. Alarming incidence of hepatitis C virus re-infection after treatment of sexually acquired acute hepatitis C virus infection in HIV-infected MSM. *AIDS.* 2011;25(17):F21-F27.

Landau DA, Scerra S, Sene D, Resche-Rigon M, Saadoun D, Cacoub P. Causes and predictive factors of mortality in a cohort of patients with hepatitis C virus-related cryoglobulinemic vasculitis treated with antiviral therapy. *J Rheumatol.* 2010;37(3):615-621.

Larney S, Kopinski H, Beckwith CG, et al. Incidence and prevalence of hepatitis C in prisons and other closed settings: Results of a systematic review and meta-analysis. *Hepatology.* 2013;58(4):1215-1224.

Lawitz E, Mangia A, Wyles D, et al. Sofosbuvir for previously untreated chronic hepatitis C infection. *N Engl J Med.* 2013a;368(20):1878-1887.

Lawitz E, Gane EJ. Sofosbuvir for previously untreated chronic hepatitis C infection. *N Engl J Med.* 2013b;369(7):678-679.

Lawitz E, Poordad FF, Pang PS, et al. Sofosbuvir and ledipasvir fixed-dose combination with and without ribavirin in treatment-naïve and previously treated patients with genotype 1 hepatitis C virus infection (LONESTAR): an open-label, randomised, phase 2 trial. *Lancet*. 2014a;383(9916):515-523.

Lawitz E, Sulkowski MS, Ghalib R, et al. Simeprevir plus sofosbuvir, with or without ribavirin, to treat chronic infection with hepatitis C virus genotype 1 in non-responders to pegylated interferon and ribavirin and treatment-naïve patients: the COSMOS randomised study. *Lancet*. 2014b;384(9956):1756-1765.

Lee SR, Kardos KW, Schiff E, et al. Evaluation of a new, rapid test for detecting HCV infection, suitable for use with blood or oral fluid. *J Virol Methods*. 2011;172(1-2):27-31.

Lewis JH, Mortensen ME, Zweig S, Fusco MJ, Medoff JR, Belder R. Efficacy and safety of high-dose pravastatin in hypercholesterolemic patients with well-compensated chronic liver disease: Results of a prospective, randomized, double-blind, placebo-controlled, multicenter trial. *Hepatology*. 2007;46(5):1453-1463.

Limketkai BN, Mehta SH, Sutcliffe CG, et al. Relationship of liver disease stage and antiviral therapy with liver-related events and death in adults coinfecting with HIV/HCV. *JAMA*. 2012;308(4):370-378.

Linas BP, Wong AY, Schackman BR, Kim AY, Freedberg KA. Cost-effective screening for acute hepatitis C virus infection in HIV-infected men who have sex with men. *Clin Infect Dis*. 2012;55(2):279-290.

Litwin AH, Harris KA, Jr., Nahvi S, et al. Successful treatment of chronic hepatitis C with pegylated interferon in combination with ribavirin in a methadone maintenance treatment program. *J Subst Abuse Treat*. 2009;37(1):32-40.

Litwin AH, Smith BD, Drainoni ML, et al. Primary care-based interventions are associated with increases in hepatitis C virus testing for patients at risk. *Dig Liver Dis*. 2012;44(6):497-503.

Lo Re V, Kallan MJ, Tate JP, et al. Hepatic decompensation in antiretroviral-treated patients co-infected with HIV and hepatitis C virus compared with hepatitis C virus-monoinfected patients: a cohort study. *Ann Intern Med*. 2014;160(6):369-379.

Lok AS, McMahon BJ. Chronic hepatitis B: update 2009. AASLD practice guideline update. *Hepatology*. 2009;50(3):661-662.

Louie KS, St LS, Forssen UM, Mundy LM, Pimenta JM. The high comorbidity burden of the hepatitis C virus infected population in the United States. *BMC Infect Dis*. 2012;12:86.

Lucas GM, Ross MJ, Stock PG, et al. Executive Summary: Clinical Practice Guideline for the Management of Chronic Kidney Disease in Patients Infected With HIV: 2014 Update by the HIV Medicine Association of the Infectious Diseases Society of America. *Clin Infect Dis*. 2014;59(9):1203-1207.

Macias J, Berenguer J, Japon MA, et al. Fast fibrosis progression between repeated liver biopsies

in patients coinfecting with human immunodeficiency virus/hepatitis C virus. *Hepatology*. 2009;50(4):1056-1063.

Maddison AR, Fisher J, Johnston G. Preventive medication use among persons with limited life expectancy. *Prog Palliat Care*. 2011;19(1):15-21.

Mahajan R, Liu SJ, Klevens RM, Holmberg SD. Indications for testing among reported cases of HCV infection from enhanced hepatitis surveillance sites in the United States, 2004-2010. *Am J Public Health*. 2013;103(8):1445-1449.

Mahale P, Kontoyiannis DP, Chemaly RF, et al. Acute exacerbation and reactivation of chronic hepatitis C virus infection in cancer patients. *J Hepatol*. 2012;57(6):1177-1185.

Mangia A, Santoro R, Minerva N, et al. Peginterferon alfa-2b and ribavirin for 12 vs. 24 weeks in HCV genotype 2 or 3. *N Engl J Med*. 2005;352(25):2609-2617.

Manns MP, Pockros PJ, Norkrans G, et al. Long-term clearance of hepatitis C virus following interferon alpha-2b or peginterferon alpha-2b, alone or in combination with ribavirin. *J Viral Hepat*. 2013;20(8):524-529.

Mantry PS, Kwo PY, Coakley E et al. High sustained virologic response rates in liver transplant recipients with recurrent HCV genotype 1 infection receiving ABT-450/r/ombitasvir+dasabuvir plus ribavirin. [Abstract 198.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.

Marcellin P, Boyer N, Gervais A, et al. Long-term histologic improvement and loss of detectable intrahepatic HCV RNA in patients with chronic hepatitis C and sustained response to interferon-alpha therapy. *Ann Intern Med*. 1997;127(10):875-881.

Martin NK, Hickman M, Hutchinson SJ, Goldberg DJ, Vickerman P. Combination interventions to prevent HCV transmission among people who inject drugs: modeling the impact of antiviral treatment, needle and syringe programs, and opiate substitution therapy. *Clin Infect Dis*. 2013a;57(Suppl 2):S39-S45.

Martin NK, Vickerman P, Grebely J, et al. Hepatitis C virus treatment for prevention among people who inject drugs: Modeling treatment scale-up in the age of direct-acting antivirals. *Hepatology*. 2013b;58(5):1598-1609.

Mazzaro C, Little D, Pozzato G. Regression of splenic lymphoma after treatment of hepatitis C virus infection. *N Engl J Med*. 2002;347(26):2168-2170.

McGovern BH, Birch CE, Bowen MJ, et al. Improving the diagnosis of acute hepatitis C virus infection with expanded viral load criteria. *Clin Infect Dis*. 2009;49(7):1051-1060.

McGowan CE, Monis A, Bacon BR, et al. A global view of hepatitis C: physician knowledge, opinions, and perceived barriers to care. *Hepatology*. 2013;57(4):1325-1332.

Mehta SH, Brancati FL, Sulkowski MS, Strathdee SA, Szklo M, Thomas DL. Prevalence of type 2 diabetes mellitus among persons with hepatitis C virus infection in the United States. *Ann Intern*

Med. 2000;133(8):592-599.

Mehta SH, Lucas GM, Mirel LB, et al. Limited effectiveness of antiviral treatment for hepatitis C in an urban HIV clinic. *AIDS*. 2006a;20(18):2361-2369.

Mehta SD, Ghanem KG, Rompalo AM, Erbedding EJ. HIV seroconversion among public sexually transmitted disease clinic patients: analysis of risks to facilitate early identification. *JAIDS*. 2006b;42:116-122.

Merchante N, Giron-Gonzalez JA, Gonzalez-Serrano M, et al. Survival and prognostic factors of HIV-infected patients with HCV-related end-stage liver disease. *AIDS*. 2006;20(1):49-57.

Miller L, Fluker SA, Osborn M, Liu X, Strawder A. Improving access to hepatitis C care for urban, underserved patients using a primary care-based hepatitis C clinic. *J Natl Med Assoc*. 2012;104(5-6):244-250.

Mira JA, Rivero-Juárez A, López-Cortes LF, et al. Benefits from sustained virologic response to pegylated interferon plus ribavirin in HIV/hepatitis C virus-coinfected patients with compensated cirrhosis. *Clin Infect Dis*. 2013;56(11):1646-1653.

Molina JM, Orkin C, Iser DM et al. All-oral therapy with sofosbuvir plus ribavirin for the treatment of HCV genotypes 1, 2, 3 and 4 infection in patients co-infected with HIV (PHOTON-2). [Abstract MOAB0105LB.] 20th International AIDS Conference. July 20-25, 2014; Melbourne, Australia

Molina JM, Orkin C, Iser DM, et al. Sofosbuvir plus ribavirin for treatment of hepatitis C virus in patients co-infected with HIV (PHOTON-2): a multicentre, open-label, non-randomised, phase 3 study. *Lancet*. 2015. [Epub ahead of print]

Moreno C, Berg T, Tanwandee T, et al. Antiviral activity of TMC435 monotherapy in patients infected with HCV genotypes 2-6: TMC435-C202, a phase IIa, open-label study. *J Hepatol*. 2012;56(6):1247-1253.

Moreno C, Hezode C, Marcellin P et al. Simeprevir with peginterferon/ribavirin for treatment of chronic HCV genotype 4 infection in treatment-naïve or -experienced patients: interim results of a phase III trial. [Abstract 60.] Hep DAART. December 8-12, 2013a; Big Island, Hawaii

Moreno C, Herzode C, Marcellin P, et al. Simeprevir with peginterferon/ribavirin in treatment-naïve or experienced patients with chronic HCV genotype 4 infection: Interim results of a Phase III trial. 14th European AIDS conference Brussels Belgium Oct 2013. Oct 16-19, 2013b; Brussels, Belgium

Morgan RL, Baack B, Smith BD, Yartel A, Pitasi M, Falck-Ytter Y. Eradication of hepatitis C virus infection and the development of hepatocellular carcinoma: a meta-analysis of observational studies. *Ann Intern Med*. 2013;158(5 Pt 1):329-337.

Morgan TR, Ghany MG, Kim HY, et al. Outcome of sustained virological responders with histologically advanced chronic hepatitis C. *Hepatology*. 2010;52(3):833-844.

Morisco F, Granata R, Stroffolini T, et al. Sustained virological response: a milestone in the treatment of chronic hepatitis C. *World J Gastroenterol*. 2013;19(18):2793-2798.

- Morrill JA, Shrestha M, Grant RW. Barriers to the treatment of hepatitis C. Patient, provider, and system factors. *J Gen Intern Med.* 2005;20(8):754-758.
- Mosley JW, Operskalski EA, Tobler LH, et al. The course of hepatitis C viraemia in transfusion recipients prior to availability of antiviral therapy. *J Viral Hepat.* 2008;15(2):120-128.
- Moyer VA. Screening for HIV: U.S. Preventive Services Task Force Recommendation Statement. *Ann Intern Med.* 2013;159(1):51-60.
- Musso G, Gambino R, Cassader M, Pagano G. A meta-analysis of randomized trials for the treatment of nonalcoholic fatty liver disease. *Hepatology.* 2010;52(1):79-104.
- Naggie S, Cooper C, Saag M et al. Ledipasvir/sofosbuvir for 12 weeks in patients coinfectd with HCV and HIV-1. Conference on Retroviruses and Opportunistic Infections (CROI) 2015. February 23-26, 2015; Seattle, WA.
- Nakamoto S, Kanda T, Wu S, Shirasawa H, Yokosuka O. Hepatitis C virus NS5A inhibitors and drug resistance mutations. *World J Gastroenterol.* 2014;20(11):2902-2912.
- Nakano Y, Kiyosawa K, Sodeyama T, et al. Acute hepatitis C transmitted by needlestick accident despite short duration interferon treatment. *J Gastroenterol Hepatol.* 1995;10(5):609-611.
- Neary MP, Cort S, Bayliss MS, Ware JE, Jr. Sustained virologic response is associated with improved health-related quality of life in relapsed chronic hepatitis C patients. *Semin Liver Dis.* 1999;19(Suppl 1):77-85.
- Nelson PK, Mathers BM, Cowie B, et al. Global epidemiology of hepatitis B and hepatitis C in people who inject drugs: results of systematic reviews. *Lancet.* 2011;378(9791):571-583.
- Nelson DR, Cooper JN, Lalezari JP et al. All-oral 12-week combination treatment with daclatasvir (DCV) and sofosbuvir (SOF) in patients infected with HCV genotype (GT) 3: ALLY-3 phase 3 study. [Abstract LB-3.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.
- Neumann UP, Berg T, Bahra M, et al. Fibrosis progression after liver transplantation in patients with recurrent hepatitis C. *J Hepatol.* 2004;41(5):830-836.
- Noda K, Yoshihara H, Suzuki K, et al. Progression of type C chronic hepatitis to liver cirrhosis and hepatocellular carcinoma--its relationship to alcohol drinking and the age of transfusion. *Alcohol Clin Exp Res.* 1996;20(1 Suppl):95A-100A.
- Nutt AK, Hassan HA, Lindsey J, Lamps LW, Raufman JP. Liver biopsy in the evaluation of patients with chronic hepatitis C who have repeatedly normal or near-normal serum alanine aminotransferase levels. *Am J Med.* 2000;109(1):62-64.
- Ortiz V, Berenguer M, Rayon JM, Carrasco D, Berenguer J. Contribution of obesity to hepatitis C-related fibrosis progression. *Am J Gastroenterol.* 2002;97(9):2408-2414.
- Osinusi A, Meissner EG, Lee YJ, et al. Sofosbuvir and ribavirin for hepatitis C genotype 1 in

patients with unfavorable treatment characteristics: a randomized clinical trial. *JAMA*. 2013;310(8):804-811.

Osinusi A, Marti M, Kohli A et al. Sofosbuvir/ledipasvir in retreatment of HCV genotype-1 patients who previously failed sofosbuvir/ribavirin therapy. [Abstract 011.] 49th Annual Meeting of the European Association for the Study of the Liver (EASL). April 9-13, 2014a; London, United Kingdom.

Osinusi A, Kohli A, Marti MM, et al. Re-treatment of chronic hepatitis C virus genotype 1 infection after relapse: an open-label pilot study. *Ann Intern Med*. 2014b;161(9):634-638.

Osinusi A, Townsend K, Kohli A, et al. Virologic Response Following Combined Ledipasvir and Sofosbuvir Administration in Patients With HCV Genotype 1 and HIV Co-infection. *JAMA*. 2015.

Ouwerkerk-Mahadevan S, Sekar V, Peeters M, Beumont-Mauviel M. The pharmacokinetic interactions of HCV protease inhibitor TMC435 with RPV, TDF, EFV, or RAL in health volunteers [Abstract 49]. 19th Conference on Retroviruses and Opportunistic Infections (CROI). March 5-8, 2012; Seattle, Washington.

Panel on Antiretroviral Guidelines for Adults and Adolescents. Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents. Department of Health and Human Services. May 1, 2014; 1-284. <http://www.aidsinfo.nih.gov/ContentFiles/AdultandAdolescentGL.pdf>. Accessed on July 24, 2014

Pawlotsky JM. Use and interpretation of virological tests for hepatitis C. *Hepatology*. 2002;36(5 Suppl 1):S65-S73.

Petta S, Camma C, Di M, V, et al. Insulin resistance and diabetes increase fibrosis in the liver of patients with genotype 1 HCV infection. *Am J Gastroenterol*. 2008;103(5):1136-1144.

Picciotto FP, Tritto G, Lanza AG, et al. Sustained virological response to antiviral therapy reduces mortality in HCV reinfection after liver transplantation. *J Hepatol*. 2007;46(3):459-465.

Pineda JA, Romero-Gómez M, Díaz-García F, et al. HIV coinfection shortens the survival of patients with hepatitis C virus-related decompensated cirrhosis. *Hepatology*. 2005;41:779-789.

Pol S, Reddy KR, Baykal T et al. Interferon-free regimens of ombitasvir and ABT-450/r with or without ribavirin in patients with HCV genotype 4 infection: PEARL-I study results. [Abstract 1928.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.

Poordad F, Manns MP, Marcellin P, et al. Simeprevir (TMC435) with Peginterferon/Ribavirin for Treatment of Chronic HCV Genotype-1 Infection in Treatment-Naive Patients: Results From QUEST-2, a Phase III Trial. *Digestive Disease Week*. May 18-21, 2013, 2013; Orlando, FL.

Poordad F, Hezode C, Trinh R, et al. ABT-450/r-Ombitasvir and Dasabuvir with Ribavirin for Hepatitis C with Cirrhosis. *N Engl J Med*. 2014;[Epub ahead of print]

Post JJ, Arain A, Lloyd AR. Enhancing assessment and treatment of hepatitis C in the custodial

setting. *Clin Infect Dis*. 2013;57(Suppl 2):S70-S74.

Poynard T, Bedossa P, Opolon P. Natural history of liver fibrosis progression in patients with chronic hepatitis C. The OBSVIRC, METAVIR, CLINIVIR, and DOSVIRC groups. *Lancet*. 1997;349(9055):825-832.

Poynard T, Cacoub P, Ratziu V, et al. Fatigue in patients with chronic hepatitis C. *J Viral Hepat*. 2002a;9(4):295-303.

Poynard T, McHutchison J, Manns M, et al. Impact of pegylated interferon alfa-2b and ribavirin on liver fibrosis in patients with chronic hepatitis C. *Gastroenterology*. 2002b;122(5):1303-1313.

Poynard T, Ratziu V, Charlotte F, Goodman Z, McHutchison J, Albrecht J. Rates and risk factors of liver fibrosis progression in patients with chronic hepatitis c. *J Hepatol*. 2001;34(5):730-739.

Pradat P, Alberti A, Poynard T, et al. Predictive value of ALT levels for histologic findings in chronic hepatitis C: a European collaborative study. *Hepatology*. 2002;36(4 Pt 1):973-977.

Proeschold-Bell RJ, Patkar AA, Naggie S, et al. An integrated alcohol abuse and medical treatment model for patients with hepatitis C. *Dig Dis Sci*. 2012;57(4):1083-1091.

Pungpapong S, Werner KT, Aqel B et al. Multicenter experience using sofosbuvir and simeprevir with/without ribavirin to treat HCV genotype 1 after liver transplantation. [Abstract 9.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.

Reddy KR, Everson GT, Flamm SL et al. Ledipasvir/sofosbuvir with ribavirin for the treatment of HCV in patients with post transplant recurrence: preliminary results of a prospective, multicenter study. [Abstract 8.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.

Regev A, Berho M, Jeffers LJ, et al. Sampling error and intraobserver variation in liver biopsy in patients with chronic HCV infection. *Am J Gastroenterol*. 2002;97(10):2614-2618.

Reilley B, Leston J, Redd JT, Geiger R. Lack of Access to Treatment as a Barrier to HCV Screening: A Facility-Based Assessment in the Indian Health Service. *J Public Health Manag Pract*. 2013;

Rockey DC, Bissell DM. Noninvasive measures of liver fibrosis. *Hepatology*. 2006;43(2 Suppl 1):S113-S120.

Rockstroh JK, Puoti M, Rodriguez-Torres M et al. Sofosbuvir and ribavirin therapy for the treatment of HIV/HCV coinfecting patients with HCV GT1-4 infection: the PHOTON-1 and -2 trials. [Abstract 195.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014; Boston, MA.

Rodriguez-Torres M, Rodriguez-Orengo JF, Gaggar A, et al. Sofosbuvir and Peginterferon Alfa-2a/Ribavirin for Treatment-Naïve Genotype 1-4 HCV-Infected Patients Who Are Coinfected With HIV. [Abstract #714.] 53rd ICAAC 2013. Sept 10-13, 2013; Denver, CO.

Rossaro L, Torruellas C, Dhaliwal S, et al. Clinical Outcomes of Hepatitis C Treated with Pegylated Interferon and Ribavirin via Telemedicine Consultation in Northern California. *Dig Dis Sci*. 2013;58(12):3620-3625.

Ruane P, Ain D, Meshrekey R, Stryker R. Sofosbuvir Plus Ribavirin in the Treatment of Chronic HCV Genotype 4 Infection in Patients of Egyptian Ancestry. 64th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 1-5, 2013; Washington DC.

Ruane PJ, Ain D, Stryker R, et al. Sofosbuvir plus ribavirin for the treatment of chronic genotype 4 hepatitis C virus infection in patients of Egyptian ancestry. *J Hepatol*. 2014.

Saadoun D, Resche RM, Thibault V, et al. Peg-IFN α /ribavirin/protease inhibitor combination in hepatitis C virus associated mixed cryoglobulinemia vasculitis: results at week 24. *Ann Rheum Dis*. 2014;73(5):831-837.

Safdar K, Schiff ER. Alcohol and hepatitis C. *Semin Liver Dis*. 2004;24(3):305-315.

Sangiovanni A, Prati GM, Fasani P, et al. The natural history of compensated cirrhosis due to hepatitis C virus: a 17-year cohort study of 214 patients. *Hepatology*. 2006;43(6):1303-1310.

Sarkar S, Jiang Z, Evon DM, Wahed AS, Hoofnagle JH. Fatigue before, during and after antiviral therapy of chronic hepatitis C: results from the Virahep-C study. *J Hepatol*. 2012;57(5):946-952.

Schmidt AJ, Falcató L, Zahno B, et al. Prevalence of hepatitis C in a Swiss sample of men who have sex with men: whom to screen for HCV infection? *BMC Public Health*. 2014;14(1):3.

Schneider MD, Sarrazin C. Antiviral therapy of hepatitis C in 2014: do we need resistance testing? *Antiviral Res*. 2014;105:64-71.

Sebastiani G, Halfon P, Castera L, et al. SAFE biopsy: a validated method for large-scale staging of liver fibrosis in chronic hepatitis C. *Hepatology*. 2009;49(6):1821-1827.

Seem DL, Lee I, Umscheid CA, Kuehnert MJ. Excerpt from PHS guideline for reducing HIV, HBV and HCV transmission through organ transplantation. *Am J Transplant*. 2013;13(8):1953-1962.

Selph, S. and Chou, R. Impact of contacting study authors on systematic review conclusions: diagnostic tests for hepatic fibrosis. <http://www.ncbi.nlm.nih.gov/books/NBK198806/>. Accessed on July 11, 2014.

Shaw K, Gennat H, O'Rourke P, Del MC. Exercise for overweight or obesity. *Cochrane Database Syst Rev*. 2006;(4):CD003817.

Shiffman ML, Suter F, Bacon BR, et al. Peginterferon alfa-2a and ribavirin for 16 or 24 weeks in HCV genotype 2 or 3. *N Engl J Med*. 2007;357:124-134.

Shiffman RN, Shekelle P, Overhage JM, Slutsky J, Grimshaw J, Deshpande AM. Standardized reporting of clinical practice guidelines: a proposal from the Conference on Guideline Standardization. *Ann Intern Med*. 2003;139(6):493-498.

Singal AG, Volk ML, Jensen D, Di Bisceglie AM, Schoenfeld PS. A sustained viral response is associated with reduced liver-related morbidity and mortality in patients with hepatitis C virus. *Clin Gastroenterol Hepatol*. 2010;8(3):280-8, 288.

Smith BD, Morgan RL, Beckett GA, et al. Recommendations for the identification of chronic hepatitis C virus infection among persons born during 1945-1965. *MMWR Recomm Rep*. 2012;61(RR-4):1-32.

Smith BD, Jewett A, Burt RD, Zibbell JE, Yartel AK, DiNenno E. "To share or not to share?" Serosorting by hepatitis C status in the sharing of drug injection equipment among NHBS-IDU2 participants. *J Infect Dis*. 2013;208(12):1934-1942.

Spaulding AC, Weinbaum CM, Lau DT, et al. A framework for management of hepatitis C in prisons. *Ann Intern Med*. 2006;144(10):762-769.

Stein MR, Soloway IJ, Jefferson KS, Roose RJ, Arnsten JH, Litwin AH. Concurrent group treatment for hepatitis C: implementation and outcomes in a methadone maintenance treatment program. *J Subst Abuse Treat*. 2012;43(4):424-432.

Stone NJ, Robinson J, Lichtenstein AH, et al. 2013 ACC/AHA Guideline on the Treatment of Blood Cholesterol to Reduce Atherosclerotic Cardiovascular Risk in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2013.

Strategies for Management of Antiretroviral Therapy (SMART) Study Group, El-Sadr WM, Lundgren JD, et al. CD4+ count-guided interruption of antiretroviral treatment. *N Engl J Med*. 2006;355:2283-2296.

Strathdee SA, Latka M, Campbell J, et al. Factors associated with interest in initiating treatment for hepatitis C Virus (HCV) infection among young HCV-infected injection drug users. *Clin Infect Dis*. 2005;40(Suppl 5):S304-S312.

Sulkowski M, Pol S, Mallolas J, et al. Boceprevir versus placebo with pegylated interferon alfa-2b and ribavirin for treatment of hepatitis C virus genotype 1 in patients with HIV: a randomised, double-blind, controlled phase 2 trial. *Lancet Infect Dis*. 2013a;13(7):597-605.

Sulkowski M, Rodriguez-Torres M, Lalezari J, et al. All-Oral Therapy With Sofosbuvir Plus Ribavirin For the Treatment of HCV Genotype 1, 2, and 3 Infection in Patients Co-infected With HIV (PHOTON-1) . *Hepatology*. 2013b;58(Number 4 Suppl 1):313A.

Sulkowski M, Rodriguez-Torres M, Lalezari JP, et al. All-Oral Therapy with Sofosbuvir Plus Ribavirin for the treatment of HCV genotype 1,2 and 3 infection in patients coinfecting with HIV (PHOTON-1). 64th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 1-5, 2013c; Washington, DC.

Sulkowski MS, Sherman KE, Dieterich DT, et al. Combination therapy with telaprevir for chronic hepatitis C virus genotype 1 infection in patients with HIV: a randomized trial. *Ann Intern Med*. 2013d;159(2):86-96.

Sulkowski MS, Naggie S, Lalezari J, et al. Sofosbuvir and ribavirin for hepatitis C in patients with HIV coinfection. *JAMA*. 2014;312(4):353-361.

Sulkowski MS, Eron JJ, Wyles D, et al. Ombitasvir, Paritaprevir Co-dosed With Ritonavir, Dasabuvir, and Ribavirin for Hepatitis C in Patients Co-infected With HIV-1: A Randomized Trial. *JAMA*. 2015.

Svoboda J, Andreadis C, Downs LH, Miller Jr WT, Tsai DE, Schuster SJ. Regression of advanced non-splenic marginal zone lymphoma after treatment of hepatitis C virus infection. *Leuk Lymphoma*. 2005;46(9):1365-1368.

Swain MG, Lai MY, Shiffman ML, et al. A sustained virologic response is durable in patients with chronic hepatitis C treated with peginterferon alfa-2a and ribavirin. *Gastroenterology*. 2010;139(5):1593-1601.

Takahashi K, Nishida N, Kawabata H, Haga H, Chiba T. Regression of Hodgkin lymphoma in response to antiviral therapy for hepatitis C virus infection. *Intern Med*. 2012;51(19):2745-2747.

Takikawa H, Yamazaki R, Shoji S, Miyake K, Yamanaka M. Normalization of urinary porphyrin level and disappearance of skin lesions after successful interferon therapy in a case of chronic hepatitis C complicated with porphyria cutanea tarda. *J Hepatol*. 1995;22(2):249-250.

Tedaldi E, Peters L, Neuhaus J, et al. Opportunistic disease and mortality in patients coinfecting with hepatitis B or C virus in the strategic management of antiretroviral therapy (SMART) study. *Clin Infect Dis*. 2008;47(11):1468-1475.

Terrault NA, Roland ME, Schiano T, et al. Outcomes of liver transplant recipients with hepatitis C and human immunodeficiency virus coinfection. *Liver Transpl*. 2012;18(6):716-726.

Thein HH, Yi Q, Dore GJ, Krahn MD. Natural history of hepatitis C virus infection in HIV-infected individuals and the impact of HIV in the era of highly active antiretroviral therapy: a meta-analysis. *AIDS*. 2008a;22(15):1979-1991.

Thein HH, Yi Q, Dore GJ, Krahn MD. Estimation of stage-specific fibrosis progression rates in chronic hepatitis C virus infection: a meta-analysis and meta-regression. *Hepatology*. 2008b;48(2):418-431.

Thomas DL. The challenge of hepatitis C in the HIV-infected person. *Annu Rev Med*. 2008;59:473-485.

Thomas DL, Villano SA, Riester KA, et al. Perinatal transmission of hepatitis C virus from human immunodeficiency virus type 1-infected mothers. Women and Infants Transmission Study. *J Infect Dis*. 1998;177(6):1480-1488.

Thorpe J, Saeed S, Moodie EE, Klein MB. Antiretroviral treatment interruption leads to progression of liver fibrosis in HIV-hepatitis C virus co-infection. *AIDS*. 2011;25(7):967-975.

Tyson GL, Kramer JR, Duan Z, Davila JA, Richardson PA, El-Serag HB. Prevalence and predictors of hepatitis B virus coinfection in a United States cohort of hepatitis C virus-infected patients.

Hepatology. 2013;58(2):538-545.

Urbanus AT, van de Laar TJ, Stolte IG, et al. Hepatitis C virus infections among HIV-infected men who have sex with men: an expanding epidemic. *AIDS*. 2009;23(12):F1-F7.

US FDA. FDA Antiviral Drugs Advisory Committee Meeting October 25, 2013: Background Package for NDA 204671 Sofosbuvir (GS-7977).
<http://www.fda.gov/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/AntiviralDrugsAdvisoryCommittee/ucm371875.htm>. Accessed on November 15, 2013a.

US FDA. FDA Introductory Remarks: Sofosbuvir NDA 204671. Presented on October 25, 2013.
<http://www.fda.gov/downloads/AdvisoryCommittees/CommitteesMeetingMaterials/Drugs/AntiviralDrugsAdvisoryCommittee/UCM375285.pdf>. Accessed on December 6, 2013b.

US Preventive Services Task Force. Screening for hepatitis C virus infection in adults: US Preventive Services Task Force recommendation statement.
<http://www.uspreventiveservicestaskforce.org/uspstf/uspshcpc.htm>. Accessed on October 28, 2013.

van de Laar T, Pybus O, Bruisten S, et al. Evidence of a large, international network of HCV transmission in HIV-positive men who have sex with men. *Gastroenterology*. 2009;136(5):1609-1617.

van de Laar TJ, Matthews GV, Prins M, Danta M. Acute hepatitis C in HIV-infected men who have sex with men: an emerging sexually transmitted infection. *AIDS*. 2010;24(12):1799-1812.

van der Meer AJ, Veldt BJ, Feld JJ, et al. Association between sustained virological response and all-cause mortality among patients with chronic hepatitis C and advanced hepatic fibrosis. *JAMA*. 2012;308(24):2584-2593.

van Heeswijk R, Garg V, Vandevoorde A, Witek J, Dannemann B. The pharmacokinetic interaction between telaprevir and raltegravir in healthy volunteers. [Abstract A1-1738a.] 51st Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC). September 17-20, 2011a; Chicago, IL

van Heeswijk R, Vandevoorde A, Boogaerts G et al. Pharmacokinetic interactions between ARV agents and the investigational HCV protease inhibitor TVR in healthy volunteers [Abstract 119]. 18th Conference on Retroviruses and Opportunistic Infections (CROI). February 27-March 2, 2011b; Boston, MA.

Veldt BJ, Heathcote EJ, Wedemeyer H, et al. Sustained virologic response and clinical outcomes in patients with chronic hepatitis C and advanced fibrosis. *Ann Intern Med*. 2007;147(10):677-684.

Vermeersch P, Van RM, Lagrou K. Validation of a strategy for HCV antibody testing with two enzyme immunoassays in a routine clinical laboratory. *J Clin Virol*. 2008;42(4):394-398.

Villano SA, Vlahov D, Nelson KE, Cohn S, Thomas DL. Persistence of viremia and the importance of long-term follow-up after acute hepatitis C infection. *Hepatology*. 1999;29(3):908-914.

von Wagner WM, Huber M, Berg T, et al. Peginterferon-alpha-2a (40KD) and ribavirin for 16 or 24 weeks in patients with genotype 2 or 3 chronic hepatitis C. *Gastroenterology*. 2005;129(2):522-527.

Vourvahis M, Plotka A, Kantaridis C, Fang A, Heera J. The effect of boceprevir and telaprevir on the pharmacokinetics of maraviroc: an open-label, fixed-sequence study in healthy volunteers [Abstract O-17]. 14th International Workshop on Clinical Pharmacology of HIV Therapy. April 22-24, 2013; Amsterdam, Netherlands.

Wai CT, Greenson JK, Fontana RJ, et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology*. 2003;38(2):518-526.

Wandeler G, Gsponer T, Bregenzer A, et al. Hepatitis C virus infections in the Swiss HIV Cohort Study: a rapidly evolving epidemic. *Clin Infect Dis*. 2012;55(10):1408-1416.

Wang C, Jia L, O'Boyle DR, et al. Comparison of daclatasvir resistance barriers on NS5A from hepatitis C virus genotypes 1 to 6: implications for cross-genotype activity. *Antimicrob Agents Chemother*. 2014;58(9):5155-5163.

Ward JW. Testing for HCV: the first step in preventing disease transmission and improving health outcomes for HCV-infected individuals. *Antivir Ther*. 2012;17(7 Pt B):1397-1401.

Wedemeyer H, Duberg AS, Buti M, et al. Strategies to manage hepatitis C virus (HCV) disease burden. *J Viral Hepat*. 2014;21(Suppl 1):60-89.

Westin J, Lagging LM, Spak F, et al. Moderate alcohol intake increases fibrosis progression in untreated patients with hepatitis C virus infection. *J Viral Hepat*. 2002;9(3):235-241.

White DL, Ratziu V, El-Serag HB. Hepatitis C infection and risk of diabetes: a systematic review and meta-analysis. *J Hepatol*. 2008;49(5):831-844.

Whitlock EP, Polen MR, Green CA, Orleans T, Klein J. Behavioral counseling interventions in primary care to reduce risky/harmful alcohol use by adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med*. 2004;140(7):557-568.

Wiley TE, McCarthy M, Breidi L, McCarthy M, Layden TJ. Impact of alcohol on the histological and clinical progression of hepatitis C infection. *Hepatology*. 1998;28(3):805-809.

Williams IT, Bell BP, Kuhnert W, Alter MJ. Incidence and transmission patterns of acute hepatitis C in the United States, 1982-2006. *Arch Intern Med*. 2011;171(3):242-248.

Witt MD, Seaberg EC, Darilay A, et al. Incident hepatitis C virus infection in men who have sex with men: a prospective cohort analysis, 1984-2011. *Clin Infect Dis*. 2013;57(1):77-84.

Wong KA, Worth A, Martin R, et al. Characterization of Hepatitis C virus resistance from a multiple-dose clinical trial of the novel NS5A inhibitor GS-5885. *Antimicrob Agents Chemother*. 2013;57(12):6333-6340.

Wyles DL, Pockros PJ, Yang JC et al. Retreatment of patients who failed prior sofosbuvir-based regimens with all oral fixed-dose combination ledipasvir/sofosbuvir plus ribavirin for 12 weeks.

[Abstract 235.] 65th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 7-11, 2014b; Boston, MA

Wyles D, Ruane P, Sulkowski M et al. Daclatasvir in combination with sofosbuvir for HIV/HCV coinfection: ALLY-2 study. [Abstract 151LB.] Conference on Retroviruses and Opportunistic Infections (CROI) 2015. February 23-26, 2015; Seattle, WA.

Yoneda M, Saito S, Ikeda T, et al. Hepatitis C virus directly associates with insulin resistance independent of the visceral fat area in nonobese and nondiabetic patients. *J Viral Hepat.* 2007;14(9):600-607.

Younossi ZM, Stepanova M, Gerber L, Nader F, Frost S, Hunt SL. P717 Improvement of central fatigue is associated with sustained virologic response (SVR) following sofosbuvir (SOF) containing regimens. *J Hepatol.* 2014;60(1):S308.

Younossi ZM, Stepanova M, Henry L, et al. Effects of sofosbuvir-based treatment, with and without interferon, on outcome and productivity of patients with chronic hepatitis C. *Clin Gastroenterol Hepatol.* 2013; [Epub ahead of print]

Yu ML, Dai CY, Huang JF, et al. A randomised study of peginterferon and ribavirin for 16 versus 24 weeks in patients with genotype 2 chronic hepatitis C. *Gut.* 2007;56(4):553-559.

Zarski JP, Bohn B, Bastie A, et al. Characteristics of patients with dual infection by hepatitis B and C viruses. *J Hepatol.* 1998;28(1):27-33.

Zeuzem S, Andreone P, Pol S, et al. Telaprevir for retreatment of HCV infection. *N Engl J Med.* 2011;364(25):2417-2428.

Zeuzem S, Berg T, Gane E, et al. Simeprevir Increases Rate of Sustained Virologic Response Among Treatment-Experienced Patients with HCV Genotype-1 Infection: a Phase IIb Trial. *Gastroenterology.* 2013a;

Zeuzem S, Dusheiko GM, Salupere R. Sofosbuvir + ribavirin for 12 or 24 weeks for patients with HCV genotype 2 or 3: the VALENCE trial. [Abstract 1085.] 64th Annual Meeting of the American Association for the Study of Liver Diseases (AASLD). November 1-5, 2013b;58:733A-734A; Washington, DC

Zeuzem S, Jacobson IM, Baykal T, et al. Retreatment of HCV with ABT-450/r-ombitasvir and dasabuvir with ribavirin. *N Engl J Med.* 2014;370(17):1604-1614.

Ziol M, Handra-Luca A, Kettaneh A, et al. Noninvasive assessment of liver fibrosis by measurement of stiffness in patients with chronic hepatitis C. *Hepatology.* 2005;41(1):48-54.

WEBSITE POLICIES

Visitor's Agreement

Welcome to www.HCVguidelines.org, a service of the Guidance. Please read this Visitor's Agreement. By using this service, you agree to its terms.

By registering with HCVguidelines.org, or by using the service, you agree that we may use and disclose, as we see fit and as is consistent with applicable law, any information that you provide or we obtain concerning your use of the service. We may change the terms of this Agreement from time to time. By continuing to use the service after we post notice of any such changes, you agree to accept the new terms.

If you disagree with any material you find on our service, we invite you to bring to our attention the material you believe to be inaccurate. Please forward a copy of the material to us at HCVguidelines@iasusa.org, along with an explanation of your objection or disagreement.

Medical Information Disclaimer

The information contained in HCVguidelines.org is presented primarily for the purpose of providing information for and educating physicians on HCV treatment and disease management topics. Nothing contained at HCVguidelines.org is intended to constitute a specific medical diagnosis, treatment, or recommendation. The information should not be considered complete, nor should it be relied on to suggest a course of treatment for a particular individual. It should not be used in place of a visit, call, consultation, or the advice of a physician or other qualified health care provider. Information obtained at HCVguidelines.org is not exhaustive and does not cover all complications, diseases, ailments, physical conditions, or their treatment with respect to HCV or any other disease. Patients who have any health care-related questions should call or see a physician or other qualified health care provider promptly. You should never disregard medical advice or delay seeking it because of something you have read at HCVguidelines.org.

The information contained at HCVguidelines.org is prepared by or compiled from a variety of sources. HCVguidelines.org, information providers, and sources do not directly or indirectly practice medicine or dispense medical services as a part of HCVguidelines.org. HCVguidelines.org may contain information about the investigational uses of drugs or products that are not approved by the U.S. Food and Drug Administration. You should consult full prescribing information before using any information or product mentioned at HCVguidelines.org. By using HCVguidelines.org, you agree to the above terms.

Copyrights and Trademarks

HCVguidelines.org expects that you will not use the service to violate anyone's rights in copyrights, trademarks, or other intellectual property. By submitting material to HCVguidelines.org, you are either representing yourself as the owner of the material or making your submission with the explicit consent of the owner. Submitting material that is the property of another without the specific consent of its owner is not only a violation of this agreement, but may also subject you to legal liability for infringement of copyright, trademark or other intellectual property rights. The materials available through HCVguidelines.org are the property of the Guidance and its licensors, and are protected by copyright, trademark, and other intellectual property laws. You are welcome to display and print for your personal,

non-commercial use information you receive through HCVguidelines.org. However, you may not otherwise reproduce any of the materials without the prior written consent of the owner. You may not distribute copies of materials found on HCVguidelines.org in any form—including by e-mail or other electronic means—without prior written permission from the owner. Of course, you are free to encourage others to access the information themselves on HCVguidelines.org and to tell them how to find it. Requests for permission to reproduce or distribute materials found on HCVguidelines.org should be submitted to RightsLink@automated permissions service at [Get Permissions](#).

Trademarks: You may not use any trademark or service mark appearing on HCVguidelines.org without the prior written consent of the owner of the mark. HCVguidelines.org and the Guidance logo are trademarks of the Guidance.

We welcome links to our service. You are free to establish a hypertext link to this site so long as the link does not state or imply any sponsorship of your site by HCVguidelines.org or the Guidance.

No Framing: Without the prior written permission of the Guidance, you may not frame any of the content of HCVguidelines.org, or incorporate into another Web site or other service any intellectual property of HCVguidelines.org, the Guidance, or of their licensors. Requests for permission to frame our content may be sent to the Guidance either by e-mail or to the address above.

Disclaimer of Warranties and Liability

Please read this Disclaimer carefully before using HCVguidelines.org:

YOU AGREE THAT YOUR USE OF THIS SERVICE IS AT YOUR SOLE RISK. BECAUSE OF THE NUMBER OF POSSIBLE SOURCES OF INFORMATION AVAILABLE THROUGH THE SERVICE, AND THE INHERENT HAZARDS AND UNCERTAINTIES OF ELECTRONIC DISTRIBUTION, THERE MAY BE DELAYS, OMISSIONS, INACCURACIES OR OTHER PROBLEMS WITH SUCH INFORMATION. IF YOU RELY ON THIS SERVICE, OR ANY MATERIAL AVAILABLE THROUGH THIS SERVICE, YOU DO SO AT YOUR OWN RISK.

THIS SERVICE IS PROVIDED TO YOU "AS IS." HCVGUIDELINES.ORG, THE GUIDANCE, AND THEIR AFFILIATES, AGENTS, AND LICENSORS CANNOT AND DO NOT WARRANT THE ACCURACY, COMPLETENESS, NONINFRINGEMENT, MERCHANTABILITY, OR FITNESS FOR A PARTICULAR PURPOSE OF THE INFORMATION AVAILABLE THROUGH THE SERVICE, NOR DO THEY GUARANTEE THAT THE SERVICE WILL BE ERROR-FREE, OR CONTINUOUSLY AVAILABLE, OR THAT THE SERVICE WILL BE FREE OF VIRUSES OR OTHER HARMFUL COMPONENTS. UNDER NO CIRCUMSTANCES SHALL HCVGUIDELINES.ORG, THE GUIDANCE, OR THEIR AFFILIATES, AGENTS OR LICENSORS BE LIABLE TO YOU OR ANYONE ELSE FOR ANY DAMAGES OTHER THAN DIRECT DAMAGES. THIS PROVISION INCLUDES, WITHOUT LIMITATION, ANY LIABILITY FOR CONSEQUENTIAL, SPECIAL, INCIDENTAL, INDIRECT OR SIMILAR DAMAGES, EVEN IF WE ARE ADVISED BEFOREHAND OF THE POSSIBILITY OF SUCH DAMAGES. YOU AGREE THAT THE LIABILITY OF HCVGUIDELINES.ORG, THE GUIDANCE, AND THEIR AFFILIATES, AGENTS, AND LICENSORS, IF ANY, ARISING OUT OF ANY KIND OF LEGAL CLAIM IN ANY WAY CONNECTED TO THE SERVICE, SHALL NOT EXCEED THE AMOUNT YOU PAID TO THE SERVICE FOR THE USE OF THE SERVICE.

By using HCVguidelines.org, you agree to abide by the terms of this Visitor's Agreement. We hope you enjoy using HCVguidelines.org, and we welcome suggestions for improvements.

